

UNIVERSITAS NAHDLATUL ULAMA SURABAYA LEMBAGA PENELITIAN DAN PENGABDIAN KEPADA MASYARAKAT

Kampus A Wonokromo : Jl. SMEA No.57 Tlp. 031-8291920, 8284508 Fax. 031-8298582 – Surabaya 60243 Kampus B RSIJemursari : Jl. Jemursari NO.51-57 Tlp. 031-8479070 Fax. 031-8433670 – Surabaya 60237

Website: unusa.ac.id Email: info@unusa.ac.id

SURAT KETERANGAN

Nomor: 145-UNUSA-LPPM/Adm-I/II/2024

Lembaga Penelitian dan Pengabdin Kepada Masyarakat (LPPM) Universitas Nahdlatul Ulama Surabaya menerangkan telah selesai melakukan pemeriksaan duplikasi dengan membandingkan artikel-artikel lain menggunakan perangkat lunak **Turnitin** pada tanggal 05 Februari 2024

Judul : Decreasing Blood Pressure with Lavender Aromatherapy in Elderly

with Hypertension

Penulis : Nur Ainiyah, Dewi Sunarti, Erika Martining Wardani, Difran

NobelBistara, Yurike Septianingrum, Andikawati Fitriasari,

Chilyatiz Zahroh

No. Pemeriksaan : 2024.02.06.084

Dengan Hasil sebagai Berikut:

Tingkat Kesamaan diseluruh artikel (Similarity Index) yaitu 19%

Demikian surat keterangan ini dibuat untuk digunakan sebagaimana mestinya.

Surabaya, 6 Februari 2024

Ketua LPPM

Achmad Syafiuddin, Ph.D

NPP: 20071300

LPPM Universitas Nahdlatul Ulama Surabaya

Website : Ippm.unusa.ac.id Email : Ippm@unusa.ac.id Hotline : 0838.5706.3867

May_Decreasing blood pressure with lavender aromatherapy in elderly with hypertension

by Erika Martining Wardani

Submission date: 05-Feb-2024 12:10PM (UTC+0700)

Submission ID: 2286648877

File name: sure_with_lavender_aromatherapy_in_elderly_with_hypertension.pdf (742.56K)

Word count: 1931

Character count: 10839

RESEARCH ARTICLE | MAY 15 2023

Decreasing blood pressure with lavender aromatherapy in elderly with hypertension *⊙*

Nur Ainiyah ■; Dewi Sunarti; Erika Martining Wardani; ... et. al





CrossMark

Articles You May Be Interested In

Aromatherapy oils of lavender (Lavandula angustifolia) inhalance reduced norepinephrine levels of woman with postpartum blues based on Edinburgh postpartum depression scale

AIP Conference Proceedings (April 2020)

Toxicity assay of lavender (Lavandula angustifolia), peppermint (Mentha piperita), and eucalyptus (Eucalyptus globulus) essential oils by Brine Shrimp Lethality Test (BSLT) method

AIP Conference Proceedings (May 2023)

Contribution of traditional medicinal and aromatic plants species on elderly health care

AIP Conference Proceedings (April 2023)

Downloaded from http://pubs.aip.org/aip/acp/article-pdf/doi/10.1063/5.0123961/17570205/090003_1_5.0123961.pdf



Time to get excited. Lock-in Amplifiers – from DC to 8.5 GHz









Decreasing Blood Pressure with Lavender Aromatherapy in Elderly with Hypertension

Nur Ainiyah^{1, a)}, Dewi Sunarti¹, Erika Martining Wardani¹, Difran Nobel Bistara¹, Yurike Septianingrum¹, Andikawati Fitriasari¹, Chilyatiz Zahroh¹

Department of Nursing, Faculty of Nursing and Midwifery, Universitas Nahdlatul Ulama Surabaya, 602310 Surabaya, East Java, Indonesia

a)corresponding author: ainiyahannuri@unusa.ac.id

Abstract. Hypertension requires appropriate management to overcome the increased blood pressure through complementary therapy. Complementary therapy with lavender essential oil is an option for hypertension, but many people don't use it. This study aimed to analyze the effect of lavender aromatherapy on blood pressure changes in the elderly with hypertension. The research design used was a pre-experimental design with a one-group pre-posttest design. The population was all elderly with hypertension. The sample consisted of 36 elderly with hypertension determined by a simple random sampling technique. The independent variable was lavender aromatherapy, and the dependent variable was blood pressure changes. The instruments of this research were an observation sheet, a sphygmomanometer, a stethoscope, a diffuser, and lavender essential oil. The research data were analyzed using the Wilcoxon Signed Rank Test. The results showed that more elderlies (52.8%) were in the moderate hypertension category before receiving lavender aromatherapy. After being given the aromatherapy, 63.9% of the elderly were in the mild hypertension category. The Wilcoxon Signed-Rank Test obtained a value of 0.000. Therefore, lavender aromatherapy influenced blood pressure changes. Lavender aromatherapy decreased blood pressure.

INTRODUCTION

The Elderly is someone who has reached the age of 60 years. Increasing age is usually followed by decreased physiological function due to a degenerative process (aging). Therefore, many degenerative diseases are found in the elderly, such as hypertension, arthritis, stroke, chronic obstructive pulmonary disease (COPD), and diabetes mellitus [1]. Hypertension is a major risk factor for heart disease and stroke. Hypertension was often referred to as the "silent killer" because often, people with hypertension go for years without feeling any disturbances or symptoms [2].

According to the Indonesian Basic Health Research (2018), South Kalimantan Province has the highest prevalence of hypertension at 44.13%, Jest Java at 39.6%, and East Kalimantan at 39.3%. On the other hand, Japua Province has the lowest hypertension prevalence at 22.2 %. Related to the medication, most patients regularly take medication (54.4%), do not take medication regularly (32,27%), and never take medication 13.33% [3].

Factors that influence changes in blood pressure are heredity, age, gender, physical and psychological stress, obesity (obesity), unhealthy eating patterns, high salt consumption, lack of physical activity, alcohol consumption, caffeine consumption, other diseases, and smoking habit. If hypertension is not treated, it can trigger stroke, damage to blood vessels (arteriosclerosis), heart failure, and kidney failure [2].

There are two ways to treat hypertension, namely pharmacological and non-pharmacological treatment. The pharmacological treatment uses diuretics, beta-blockers, sympatholytics, central, alpha-blockers, arterial vasodilators, calcium channel blockers, ACE inhibitors, and type I angiotensin II receptor antagonists. Besides that, there is a laternative treatment (non-pharmacological therapy) which includes acupressure (acupuncture without needles), herbal medicine from China, juice therapy, herbal therapy, massage, yoga, breathing and relaxation, and aromatherapy [4]. One that can be used as a non-pharmacological therapy using aromatherapy is aromatherapy. Plants that function as aromatherapy are lavender flowers because they can be used for relaxation [5].

Aromatherapy is a healing paces used to use pure aromatic plant extracts. The goal is to improve phaceal, mental, and emotional health. One of the plants that have a function as aromatherapy is lavender. When aromatherapy is inhaled, volatile molecules will carry aromatic elements that stimulate memory and emotional responses that cause feelings of calm and relaxation and facilitate blood flow [6]. Lavender oil already has a lot of potentials, and there are several ingredients such as camphene, limonene, nerol, and most of them contain linalool with the amount of 30%-60% of the total weight of the oil, which is the main active ingredient in relaxation [7]

METHOD

General Background of Research

The research design used 3 as a one-group pre-post-test design. The population was all patients with hypertension. The independent variable in this study was blood pressure. The respondents measured their blood pressure levels before and before the treatments.

The Sample of the Research

This research involved 36 respondents determined using the purposive sampling technique. The populations involved were patients suffering from hypertension. The population in this study was those who had inclusion criteria of hypertension and did not have complications such as heart disease, heart failure, kidney failure, or stroke.

Instrument and Procedures

This study measured blood pressure using a sphygmomanometer before and after giving lavender aromatherapy.

Data Analysis

The data analysis to determine the blood pressure was Wilcoxon Signed Rank Test. The level of significance was set at P < 0.01.

RESULTS

TABLE 1. Distribution of Respondents by Age, Gender, Duration of Diabetes, Smoking Habit, Medication Adherence, and

Exercise					
Characteristics of Respondent	Frequency	(%)			
Gender					
Female	11	30,6			
Male	25	69,4			
Age					
45-59	7	19.5			
60-74	21	58.3			
75-90	8	22.2			
Having a Family with hypertension					
Yes	24	66.7			
No	12	33.3			
Medicine Consumption					
Regularly	14	38.9			
Irregularly	22	61.1			
Exercise					
Regularly	10	27.8			
Irregularly	26	72.2			
Coffee Consumption Habit					
Yes	29	80.6			

Characteristics of Respondent	Frequency	(%)
No	7	19.4
Smoking Habit		
Yes	30	83.3
No	6	16.7
High Salt Consumption		
Yes	20	55.6
No	16	44.4
Having stress		
Often	7	19.4
Rarely	27	75
No	2	5.6
Total	36	100,0

Table 1 describes that most respondents were male (69.4%), 60-74 years old (58.3%), had a family with hypertension (66.7%), consumed the medicine irregularly (61.1%), exercised irregularly (72.2%), had coffee consuming habit (80.6%), had a smoking habit (83.3%), had high salt consumption habit (55.6), and rarely experienced stress (75%).

TABLE 2. The results of the Wilcoxon Signed-Rank Test

Blood Pressure	Pre		Post	
	(F)	(%)	(F)	(%)
Normal	0	0	7	19.4
Mild HT	14	36.1	23	63.9
Moderate HT	19	52.8	6	16.7
Severe HT	4	11.1	0	0
Total	36	100	36	100
Wilcoxon	$\rho = 0,000$			

DISCUSSION

Based on Table 2, most of the respondents' blood pressures (52.8%) were moderate before giving lavender aromatherapy. Moderate hypertension was a person's blood pressure with a systolic value of 160-179 mmHg and a diasolic value of 100-109 mmHg [8]. Hypertension is influenced by several uncontrolled and controlled factors [9].

Risk factors that cannot be controlled are gender, history of hypertension, and unwillingness to take medication. Based on Table 1, out of the 36 respondents, the majorities (69.4%) of respondents were female, aged 60-74 years (58.3%), had a history of hereditary hypertension (66.7%), did not take anti-hypertensive drugs regularly (61.1%).

Females are at risk of suffering from hypertension [10]. Another research showed that women had a 1.4 times risk of suffering from hypertension than men [11]. Menopausal conditions can increase women's risk of hypertension because the estrogen hormone decreases. It causes organ functions to decrease, such as hardened atrial vessels, damaged endothelial cells, and causes plaque in the blood, thereby stimulating blood pressure.

Age is a risk factor for hypertension that cannot be prevented because, naturally, age continues to grow. This factor cannot be controlled. Control efforts that can be done are to implement a healthy lifestyle by changing a healthy diet and exercising.

Other trigger factors that cause an increase in blood pressure are lack of exercise, coffee drinking habits, smoking, consumption of salty or high-salt foods, and stress. Table 1 shows that 72.2% of the respondents did not exercise regularly, 80.6% did not consume coffee, 83.3% did not smoke, 55.6% consumed salty or high-salt foods, and 75% of the respondents often experienced stress.

The results of this study are in line with research by Liu et al (2017), who show a relationship between physical activity factors and the occurrence of hypertension [12]. Regular exercise has a risk to reduce the occurrence of atherosclerosis, while atherosclerosis is one of the risk factors for hypertension [14]. Hypertensive sufferers who take medication, if they exercise, will help the drugs they take work [13].

Based on the Wilcoxon Signed Rank Test with a significance value of 0.05, the obtained value was 0.000. Therefore, H_1 was accepted. It means that lavender aromatherapy affects blood pressure changes in the elderly.

Lavender aromatherapy can lower blood pressure because aromatherapy can help cure hypertension sufferers in controlling blood pressure by minimizing triggers from increasing high blood pressure, such as stress. Lavender aromatherapy can relax hypertensive sufferers, reduce anxiety levels, and overcome sleep disorders.

CONCLUSIONS

Lavender aromatherapy can decrease blood pressure for the elderly

ACKNOWLEDGEMENTS

Thanks to LPPM Universitas Nahdlatul Ulama Surabaya, which had funded this project research. Furthermore, the researchers' team who have been willing to spend their time and provide input.

REFERENCES

- J. Preußner, Y. Rudnik, H. Brehm, R. Völkl, and U. Glatzel, Int. J. Plast. 25, 973–994 (2009).
- R. Pazoki et al., Circulation. 137, 653–661 (2018).
- 3. Kemenkes RI, "Hasil Riset Kesehatan Dasar Tahun 2018," Kementrian Kesehat. RI, 53, 1689-1699 (2018).
- 4. I. Levy, S. Attias, E. Ben-Arye, B. Bloch, and E. Schiff, Int. J. Geriatr. Psychiatry. 32, 492–508 (2017).
- L. Franco et al., J. Clin. Anesth. 33, 243–249 (2016).
- 6. E. Matsubara, T. Morikawa, N. Kusumoto, K. Hashida, N. Matsui, and T. Ohira, Molecules. 26, 1-14 (2021).
- K. Bhardwaj et al. Phyther. Res. 34, 2889–2910 (2020).
- 8. R. M. Carey et al., Ann. Intern. Med. 168, 351–358 (2018).
- F. S. Sarfo et al., PLoS One. 13, 1–19 (2018).
- 10. J. F. Reckelhoff, Hypertension. 37, 1199–1208 (2001).
- 11. K. T. Mills, A. Stefanescu, and J. He, Nat. Rev. Nephrol. 16, 223–237 (2020).
- 12. X. Liu et al., Hypertension. 69, 813-820 (2017.
- N. Alshammari, S., Almazyed, O. M., Alshoweir, A. A., Alhelal, S. H., AbuHamra, N., Alsouaan, A. A., & Aljarboa, International Journal of Academic Scientific Research. 5, 207-222 (2017).
- 14. T. P. Utami, E. Isfandiary, and I. Ilyas, J Indon Med Assoc, 66, 574-578 (2016).

May_Decreasing blood pressure with lavender aromatherapy in elderly with hypertension

ORIGINA	ALITY REPORT			
SIMILA	9% ARITY INDEX	14% INTERNET SOURCES	12% PUBLICATIONS	13% STUDENT PAPERS
PRIMAR	Y SOURCES			
1	Submitt Surabay Student Pape		s Nahdlatul U	lama 4 _%
2	reposito Internet Sour	ory.pnb.ac.id		3%
3	jgrph.or Internet Sour			3%
4	Submitt Student Pape	ed to Universita	s Sumatera U	tara 2%
5	Syabari Lowerin	Alfin, Destiani Ra yah. "Lemon Ard ng Blood Pressur ension", Jurnal K	omatherapy or re in Elderly wi	ith
6	ejourna Internet Sour	l.poltekkes-smg	.ac.id	2%
7	Submitt Student Pape	ed to University	of Aberdeen	2%

Exclude quotes Off Exclude matches < 2%

Exclude bibliography Off