

## **SURAT KETERANGAN**

Nomor: 081/UNUSA-LPPM/Adm-I/I/2023

Lembaga Penelitian dan Pengabdian Kepada Masyarakat (LPPM) Universitas Nahdlatul Ulama Surabaya menerangkan telah selesai melakukan pemeriksaan duplikasi dengan membandingkan artikel-artikel lain menggunakan perangkat lunak **Turnitin** pada tanggal 16 Januari 2023

Judul : Cycling to Regulate Random Blood Glucose Levels in Individuals with Diabetes

Penulis : Difran Nobel Bistara, Susanti Susanti, Budhi Setianto, Erika Martining Wardani, Dyah Ika Krisnawati, Nantiya Pupuh Satiti

No. Pemeriksaan : 2023.01.17.053

Dengan Hasil sebagai Berikut:

**Tingkat Kesamaan diseluruh artikel (*Similarity Index*) yaitu 18%**

Demikian surat keterangan ini dibuat untuk digunakan sebagaimana mestinya.

Surabaya, 17 Januari 2023

Ketua LPPM



UNUSA  
LPPM

Achmad Syafiuddin, Ph.D

NPP: 20071300

**LPPM Universitas Nahdlatul Ulama Surabaya**

Website : [lppm.unusa.ac.id](http://lppm.unusa.ac.id)

Email : [lppm@unusa.ac.id](mailto:lppm@unusa.ac.id)

Hotline : 0838.5706.3867

# Paper 12

*by* Difran Nobel Bistara

---

**Submission date:** 16-Jan-2023 09:30AM (UTC+0700)

**Submission ID:** 1993300288

**File name:** Maret\_Open\_Access\_Macedonia\_Journal\_-\_Difran\_Nobel\_Bistara.pdf (273.61K)

**Word count:** 4003

**Character count:** 21578



## Cycling to Regulate Random Blood Glucose Levels in Individuals with Diabetes

Difran Nobel Bistara<sup>1</sup>, Susanti Susanti<sup>2\*</sup>, Budhi Setianto<sup>3</sup>, Erika Martining Wardani<sup>1</sup>, Dyah Ika Krisnawati<sup>4</sup>, Nantiya Pupuh Satiti<sup>5</sup>

<sup>1</sup>Department of Nursing, <sup>2</sup>Faculty of Nursing and Midwifery, Universitas Nahdlatul Ulama Surabaya, Surabaya, East Java, Indonesia; <sup>3</sup>Department of Nursing, Sekolah Tinggi Ilmu Kesehatan Adi Husada, Surabaya, East Java, Indonesia; <sup>4</sup>Department of Public Health, Faculty of Health, Universitas Nahdlatul Ulama Surabaya, Surabaya, East Java, Indonesia; <sup>5</sup>Department of Nursing, Akademi Keperawatan Dharma Husada, Kediri, East Java, Indonesia; <sup>6</sup>Department of Nursing, Akademi Kesehatan Rustida, Banyuwangi, East Java, Indonesia

**Edited by:** Ksenija Bogoeva-Kostovska  
**Citation:** Bistara DN, Susanti S, Setianto B, Wardani EM, Krisnawati DI, Satiti NP. Cycling to Regulate Random Blood Glucose Levels in Individuals with Diabetes. Open Access Maced J Med Sci. 2022 Mar 05; 10(T5):157-161. <https://doi.org/10.3889/oamjms.2022.7821>

**Keywords:** Cycling; Random blood glucose levels; Diabetes  
**\*Correspondence:** Nantiya Pupuh Satiti, Department of Nursing, Sekolah Tinggi Ilmu Kesehatan Adi Husada, Surabaya, East Java, Indonesia.  
 E-mail: [nobel@unusa.ac.id](mailto:nobel@unusa.ac.id)

**Received:** 05-Nov-2021  
**Revised:** 17-Feb-2022  
**Accepted:** 25-Feb-2022

**Copyright:** © 2022 Difran Nobel Bistara, Susanti Susanti, Budhi Setianto, Erika Martining Wardani, Dyah Ika Krisnawati, Nantiya Pupuh Satiti  
**Funding:** This study was supported by Universitas Nahdlatul Ulama Surabaya

**Competing Interests:** The authors have declared that no competing interests exist  
**Open Access:** This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

### Abstract

**BACKGROUND:** In Indonesia, the four pillars of diabetes management include health education, food planning, physical exercise, and drug adherence. However, the most common imprudence in those four pillars was ignoring physical activity. Cycling has become a new social activity and a lifestyle among the community during the COVID-19 pandemic. It is an aerobic exercise that increases insulin receptor sensitivity.

**AIM:** This study aims to analyze the effect of cycling on Random Blood Glucose (RBG) levels in individuals with diabetes.

**METHODOLOGY:** This paper used a quasi-experiment pre-post test design with the control group. It utilized total sampling with 60 respondents. The independent variable was cycling using a dynamic bicycle. Meanwhile, the dependent variable was RBG levels with a glucometer as the instrument. The procedure in the intervention group was cycling using a dynamic bicycle twice a week with a distance of 2–3 kilometers each session. The data analysis used a paired T-test and independent sample T-test.

**RESULTS:** After cycling, the independent T-test result was  $p = 0.00$  ( $p < 0.05$ ). Thus, there was a difference in the mean RBG levels between the intervention and control groups after cycling. There was a decrease in RBG levels in the intervention group ( $206.67 \pm 69.887$  in pre-test and  $114.60 \pm 30.395$  in post-test). In addition, the paired T-test resulted in  $p = 0.00$  ( $p < 0.05$ ). Thus, there was a difference in the intervention group's mean RBG levels before and after cycling.

**CONCLUSION:** Cycling can lower RBG levels in individuals with diabetes.

### Introduction

Various epidemiological studies indicate a trend increase in the incidence and prevalence of diabetes mellitus in the world [1]. The incidence of diabetes cases has increased every year. Theoretically, individuals with type 2 diabetes (T2D) are mostly the elderly. However, a previous study in 2013 showed that most of them were adults—approximately 382 million [2]. Epidemiological data states that between 2013 and 2035, the number of people with diabetes will reach 210 million [3], [4]. It is due to ignorance about the disease and non-adherence to dm management. In Indonesia, the four pillars of diabetes management include health education, food planning, physical exercise, and drug adherence [5]. However, the most common imprudence in those four pillars was ignoring physical activity [6].

Research on the effect of exercise on blood glucose levels has been widely carried out. A study evaluated the impact of aerobic exercise and walking using the continuous glucose measurement method.

The study revealed that acute moderate-intensity aerobic exercise before breakfast reduced the morning blood glucose rise in individuals with T2D. In addition, prior research also evaluated the effect of 12 weeks of moderate-intensity activity on blood glucose responses using the Generalized Estimating Equations. The study found that moderate-intensity exercise training for 12 weeks was safe for individuals with T2D [3], [6].

Those studies only focused on exercise, walking [7], and moderate-intensity exercise for 12 weeks [6]. Research on cycling to regulate RBG levels in individuals with diabetes is still minimal. Cycling has become a new social activity and a lifestyle among the community during the COVID-19 pandemic [8]. Cycling is an aerobic exercise that increases insulin receptor sensitivity to convert glucose into energy through metabolism. In addition, it impacts the endocrine system to regulate the pancreas gland in producing insulin to manage blood glucose levels [9].

Therefore, this study aims to analyze the effect of cycling on Random blood glucose (RBG) levels in

individuals with diabetes. This article can contribute as a policy recommendation for managing diabetes.

## Methods

### Study design

This article used a quasi-experiment pre-post test design with the control group [10]. The population was individuals with diabetes registered at Tanah Kali Kedinding Public Health Center (PHC), Surabaya. Inclusion criteria were (1) Individuals with diabetes without complications (2) Do routine checks at the Tanah Kali Kedinding PHC once a month (3) Diabetes management without using diabetes medication (4) Complying with the guidelines diet management as evidenced by recording the evaluation in the diabetic patient's diet guide [11]. This research used total sampling with 60 respondents, 30 respondents in the intervention group, and 30 in the control group. The independent variable was cycling using a dynamic bicycle. Meanwhile, the dependent variable was RBG levels. The authors measured RBG levels 30 min before the first cycling schedule and 30 min after the last cycling schedule with a glucometer as the instrument. We did this research from February to March 2021.

### Data collection

The cycling procedure in this study refers to previous research with slight modifications [12]. Before cycling (pre-test), the authors measured RBG levels in intervention and control groups. Then, the procedure in the intervention group was cycling using a dynamic bicycle carried out jointly by the researchers and the respondents. The cycling schedule was divided into three groups, and each group consisted of 10 respondents. The cycling schedule was twice a week with a distance of 2–3 kilometers each session. Respondent's speed was adapted to local conditions-for example, traffic jams. Meanwhile, the control group was not given any intervention during the study. However, after the pre-test,

we provided health education about the benefits of cycling to regulate blood sugar to fulfill the principle of fairness in ethics. 30 min after the last cycling (post-test), the researchers remeasured RBG levels in intervention and control groups [12]. The authors and respondents followed the COVID-19 prevention health protocol guidelines during the study. We used disposable medical masks or two layers of cotton cloth masks. In addition, we always kept hand hygiene by cleaning hands with hand sanitizer or using soap and running water for 20 s and avoiding touching the face area. Furthermore, we maintained a minimum distance of 1 meter.

### Data analysis

After collecting data, the data was processed with editing, coding, tabulating, and scoring. The data analysis used a paired T-test to compare RBG levels before and after cycling in intervention and control groups. In addition, an independent sample T-test to compare RBG levels of intervention and control groups before and after cycling. Furthermore, we did homogeneity analysis in the characteristics of respondents to control the confounding factors. The homogeneity analysis for age and educational levels variables used the Mann-Whitney test, while for gender and diabetes duration variable utilized the Chi-square test [13].

### Ethical clearance

This article was declared ethically feasible by the Health Research Ethics Committee of Universitas Nahdlatul Ulama Surabaya with certificate number 121.1/EC/KEPK/UNUSA/2021. Data collection began with informed consent and used anonymous names for respondents.

## Results

Table 1 shows the characteristics of respondents in this article.

**Table 1: Characteristics of respondents by age, gender, educational levels, and duration of diabetes**

Characteristics of respondents	Control group		Intervention group		Homogeneity test
	Frequency	Percentage	Frequency	Percentage	
Age (years)					p = 1.00
45-54	5	16.7	4	13.3	
55-65	16	53.3	18	60	
66-74	9	30	8	26.7	
Sex					p = 0.75
Male	9	30	11	36.7	
Female	21	70	19	63.3	
Educational Level					p = 0.87
Elementary School	4	13.3	5	16.7	
Junior High School	6	20	9	30	
Senior High School	18	60	15	50	
University	2	6.7	1	3.3	
Duration of diabetes (years)					p = 0.37
<5	21	70	27	90	
>5	9	30	3	10	
Total	30	100	30	100	

**Table 2: Random blood glucose (RBG) levels before and after cycling in the control and intervention groups**

Variable	Control group (Mean ± SD)	Intervention group (Mean ± SD)	mean Difference	95% CI	p*
RBG levels before cycling	176.57 ± 88.627	206.67 ± 69.887	-30.100	-71.349; 11.149	0.149
RBG levels after cycling	182.70 ± 81.518	114.60 ± 30.395	68.100	36.305; 99.895	0.00

\*p < 0.05 based on independent t-test, RBG: Random blood glucose, SD: Standard deviation.

Most respondents in the control group were 45–65 years old (53.3%) and female (70%). In addition, most of them graduated from Senior High School (60%), and their diabetes duration was <5 years (70%). Meanwhile, most respondents in the intervention group were 55–65 years old (60%) and female (63.3%). In addition, most of them graduated from Senior High School (50%), and their diabetes duration was <5 years (90%). All homogeneity tests resulted in  $p > 0.05$  ( $p = 1.00$  in age variable,  $p = 0.75$  in sex variable,  $p = 0.87$  in educational level variable and  $p = 0.37$  in diabetes duration variable). Thus, there were no differences in the characteristics of respondents between the control and intervention groups (Table 1).

Before cycling, the independent T-test result was  $p = 0.149$  ( $p > 0.05$ ). Thus, there was no significant difference in the mean RBG levels between the intervention and control groups before cycling ( $176.57 \pm 88.627$  in the control group and  $206.67 \pm 69.887$  in the intervention group). Meanwhile, after cycling, the independent T-test result was  $p = 0.00$  ( $p < 0.05$ ). Thus, there was a difference in the mean RBG levels between the intervention and control groups after cycling ( $182.70 \pm 81.518$  in the control group and  $114.60 \pm 30.395$  in the intervention group (Table 2).

There was an increase in the average RBG levels in the control group ( $176.57 \pm 88.627$  pre-test and  $182.70 \pm 81.518$  in post-test). In addition, the paired T-test resulted in  $p = 0.342$  ( $p > 0.05$ ). Thus, there was no difference in the mean RBG levels before and after the intervention in the control group. Meanwhile, there was a decrease in mean RBG levels in the intervention group ( $206.67 \pm 69.887$  in pre-test and  $114.60 \pm 30.395$  in post-test). In addition, the paired T-test resulted in  $p = 0.00$  ( $p < 0.05$ ). Thus, there was a difference in the intervention group's mean RBG levels before and after cycling (Table 3).

## Discussion

### Identification of RBG levels before cycling

There was no significant difference in the mean RBG levels between the intervention and control groups before cycling. Before cycling, the mean RBG levels in

the intervention group were 206.67 mg/dL (Table 1). It is categorized as hyperglycemia.

Most respondents in this study were 55–65 years old (Table 1). According to the World Health Organization (WHO), the age is classified as elderly. One of the risk factors for diabetes is age. The elderly have impaired glucose tolerance due to insulin resistance. So that there are changes in body composition and diet also decreased activity and neurohormonal function [14].

In addition, most respondents in this paper were females (Table 1). Sex was also a risk factor for diabetes. The incidence of diabetes was higher in females than males. It was due to decreased estrogen hormone levels after menopause. The changes in hormone levels would trigger fluctuations in RBG levels [15].

Furthermore, most respondents graduated from Senior High School (Table 1). According to the Law of the Republic of Indonesia on National Education System Number 20 the Year 2003, it is classified as secondary education. Another risk factor for diabetes was educational level. Education could affect a person's ability and knowledge to implement healthy living behaviors, especially diabetes management [16].

Moreover, the duration of diabetes in most respondents was <5 years (Table 1). A study showed an independent association between diabetes duration and microvascular events in T2D diabetics. The study reported that for each 5-year increase in diabetes duration, the risk of microvascular events was increased by 28% [17].

There were no differences in the characteristics of respondents between the control and intervention groups (Table 1). Characteristics of respondents were confounding factors in this paper. Thus, we controlled confounding factors by analyzing homogeneity in the characteristics of respondents.

### Identification of RBG levels after cycling

There was a difference in the mean RBG levels between the intervention and control groups after cycling, in the control group were 182.70 mg/dL while in the intervention group were 114.60 mg/dL (Table 2). Respondents who regularly cycled had lower average

**Table 3: Changes random blood glucose (RBG) levels before and after cycling in the control and intervention groups**

RBG levels	Pre-test (Mean ± SD)	Post-test (Mean ± SD)	95% CI	T	p*
Control group	176.57 ± 88.627	182.70 ± 81.518	-19.119; 6.853	0.966	0.342
Intervention group	206.67 ± 69.887	114.60 ± 30.395	72.701; 111.432	9.723	0.00

\*p < 0.05 based on paired t-test, RBG: Random blood glucose, SD: Standard deviation.

RBG levels than respondents who did not. It is in line with a previous study. The study revealed that more active individuals had lower RBG levels than less active ones [12].

Physical exercise can improve insulin sensitivity. A study revealed that physical activity must fulfill frequency 3–5 times per week, with light and moderate intensity, and duration 30–60 min per session [6]. Cycling is an aerobic exercise that can increase insulin receptor sensitivity to convert glucose into energy through metabolism. Cycling impacts the endocrine system because the endocrine system regulates the pancreas gland that produces insulin to regulate RBG levels [9]. Cycling affects the endocrine system because the endocrine system regulates the pancreas gland that produces insulin to regulate RBG levels). Insulin is a carbohydrate homeostatic protective hormone. It reduces RBG levels by promoting stored glucose utilization, storage, and metabolic conversion. In addition, it increases the sensitivity of insulin work. The glucose will be used as energy and can reduce glucose levels in the blood [18].

#### **The effect of cycling on RBG levels**

There was a difference in the intervention group's mean RBG levels before and after cycling. Before cycling, the mean RBG levels in the intervention group were 206.67 mg/dL-hyperglycemia. Meanwhile, after cycling, the mean RBG levels in the intervention group were 114.60 mg/dL-normoglycemia (Table 3). Cycling affected decreased RBG levels in individuals with diabetes. Respondents with hyperglycemia experienced lower RBG levels after regular cycling with the direction of the researchers. Both males and females who cycled experienced a decrease in RBG levels.

Meanwhile, there were increased RBG levels in pre-test and post-test in the control group. In pre-test were 176.57 mg/dL and in post-test were 182.70 mg/dL (Table 3). In the control group, respondents did not do the physical activity by cycling but only controlled their diet so that RBG levels tended to increase. Research showed that decreased physical activity could increase the incidence of diabetes [19]. During physical activity, muscles will increase the burning of glucose and cause reduced blood sugar levels.

A prior study showed that dynamic physical activities involving the main muscles would cause increased permeability in contracting muscles. So that insulin receptors will be more sensitive during exercise [20]. Cycling is an activity that depends on the availability of oxygen to help burn energy sources. It also depends on the organs' optimal work to transport oxygen so that the combustion process of energy sources can run perfectly. In cycling sports, there is the burning of stored carbohydrates, fats, and a small

portion ( $\pm 5\%$ ) of protein stored in the body to produce ATP (adenosine triphosphate). As a result, it causes RBG levels to return to normal [12].

The frequency of exercise in this study met the minimum standard of 2 times per week. According to the American Diabetes Association (ADA), people with diabetes should do at least 150 min of aerobic physical activity per week with moderate intensity. The ADA also encourages people with T2D without contraindications to exercise twice a week. Cycling is one of the physical activities that people with diabetes can do [21].

A combination of four pillars in diabetes management is crucial. It includes medication and diet adherence for people with diabetes. Respondents who always follow therapy regularly and adhere to diet will cause a decrease in RBG levels [11], [22]. Effective health education will create self-management and increased adherence to the recommendations of health workers [23]. The successful management of T2D needs patient acceptance and adaptation to changes due to diabetes disease. In addition, they must build commitment to comply with diabetes management. Optimal health increases the quality of life in people with diabetes [11].

#### **Limitations**

There may be some possible limitations in this study. First, the study does not analyze psychological status as confounding factors. Second, the small sample examined limits the generalizability of the finding in this study.

#### **Conclusion**

Cycling can lower RBG levels in individuals with diabetes. Further research should evaluate the use of the stationary bicycle for speed consistency in cycling. In addition, it must measure continuous glucose monitoring to analyze changes after physical activity by cycling.

#### **Acknowledgments**

The authors would like to thank Universitas Nahdlatul Ulama Surabaya for supporting and facilitating this paper. In addition, we deeply acknowledge a great collaborative effort from Stikes Adi Husada Surabaya, Akademi Keperawatan Dharma Husada Kediri and Akademi Kesehatan Rustida Banyuwangi.

## References

- Hajime M, Okada Y, Mori H, Otsuka T, Kawaguchi M, Miyazaki M, et al. Twenty-four-hour variations in blood glucose level in Japanese Type 2 diabetes patients based on continuous glucose monitoring. *J Diabetes Investig.* 2018;9(1):75-82. <https://doi.org/10.1111/jdi.12680>  
PMid:28418217
- Forouhi NG, Wareham NJ. Epidemiology of diabetes. *Medicine (Abingdon).* 2014;42(12):698-702. <https://doi.org/10.1016/j.mpmed.2014.09.007>  
PMid:25568613
- Wild S, Roglic G, Green A, Sicree R, King H. Global prevalence of diabetes: Estimates for the year 2000 and projections for 2030. *Diabetes Care.* 2004;27(5):1047-53. <https://doi.org/10.2337/diacare.27.5.1047>  
PMid:15111519
- IDF Diabetes Atlas Group. Update of mortality attributable to diabetes for the IDF diabetes atlas: Estimates for the year 2013. *Diabetes Res Clin Pract.* 2015;109(3):461-5. <https://doi.org/10.1016/j.diabres.2015.05.037>  
PMid:26119773
- Tritisari KP, Ariestingsih AD, Handayani D, Kusumastuty I. The relationship among four pillar of diabetes mellitus management with blood glucose levels and nutritional status in middle-aged diabetic adult. *Res J Life Sci.* 2018;5(7):23-33.
- Chiang S, Heitkemper MM, Hung Y, Tzeng W, Lee M, Lin C. Effects of a 12-week moderate-intensity exercise training on blood glucose response in patients with Type 2 diabetes: a prospective longitudinal study. *Medicine (Baltimore).* 2019;98(36):e16860. <http://doi.org/10.1097/MD.00000000000016860>  
PMid:31490370
- Zheng X, Qi Y, Bi L, Shi W, Zhang Y, Zhao D, et al. Effects of exercise on blood glucose and glycemic variability in Type 2 diabetic patients with dawn phenomenon. *Biomed Res Int.* 2020;2020:6408724. <https://doi.org/10.1155/2020/6408724>  
PMid:32149118
- Alvian MA, Priyono B. Cycling culture is a movement the new social community to maintain physical fitness when COVID-19 pandemic. *Indones J Phys Educ Sport.* 2021;2(1):6-45.
- Rojas-Rueda D. Health impacts of urban bicycling in Mexico. *Int J Environ Res Public Health.* 2021;18(5):2300. <https://doi.org/10.3390/ijerph18052300>  
PMid:33652688
- Nursalam N. Concept and Application of Nursing Research Methodology, Salemba Medika; 2016.
- Bistara DN, Susanti S, Wardani EM, Septianingrum Y, Ainiyah N, Fitriyani A, et al. Acceptance and commitment therapy (ACT) on increasing the compliance of management diabetes mellitus Type 2. *Int J Psychosoc Rehabil.* 2020;24(9):942-6. <https://doi.org/10.37200/IJPR/V24I9/PR290115>
- Yuliasih Y. Relationship of Body Weight with Decreased Glucose Levels. Relationship Between Body Weight and Decrease in RBG Levels after 30 Minutes of Cycling; 2012. p. 189-92.
- Ismail F. Statistics for Research in Education and Social Sciences. Jakarta: Prenamedia Group; 2018.
- Liu L, Liu J, Xu L, Ke W, Wan X, Li H, et al. Lower mean blood glucose during short-term intensive insulin therapy is associated with long-term glycemic remission in patients with newly diagnosed Type 2 diabetes: Evidence based recommendations for standardization. *J Diabetes Investig.* 2018;9(4):908-16. <https://doi.org/10.1111/jdi.12782>  
PMid:29193795
- Hu X, Yu W, Yang L, Pan W, Si Q, Chen X, et al. Inverse association between physical activity and blood glucose is independent of sex, menopause status and first-degree family history of diabetes. *J Diabetes Investig.* 2019;10(6):1502-9. <https://doi.org/10.1111/jdi.13062>  
PMid:31012524
- Clara H. The Relationship of Education and Knowledge with Type 2 Diabetes Mellitus Self-Management Behavior; 2020. Available from: <https://www.akper-pasarrebo.e-journal.id/nurs/article/view/44> [Last accessed on 2021 Sep 13].
- Zoungas S, Woodward M, Li Q, Cooper ME, Hamet P, Harrap S, et al. Impact of age, age at diagnosis and duration of diabetes on the risk of macrovascular and microvascular complications and death in Type 2 diabetes. *Diabetologia.* 2014;57(12):2465-74. <https://doi.org/10.1007/s00125-014-3369-7>  
PMid:25226881
- Triana L, Salim M. Differences in RBG levels 2 hours post prandial. *J Equator Lab.* 2017;1(1):51. <https://doi.org/10.30602/jlk.v1i1.97>
- Cicilia C. The relationship of physical activity with the incidence of diabetes mellitus in outpatients at the bitung city regional general hospital. *KESMAS J.* 2018;7(5):
- Seo DY, Ko JR, Jang JE, Kim TN, Youm JB, Kwak H, et al. Exercise as a potential therapeutic target for diabetic cardiomyopathy: Insight into the underlying mechanisms. *Int J Mol Sci.* 2019;20(24):6284. <https://doi.org/10.3390/ijms20246284>  
PMid:31842522
- Lewis S. Medical Surgical Nursing Assessment and Management of Clinical Problems. Amsterdam, Netherlands: Elsevier, Mosby; 2011.
- Susanti S, Bistara DN. The effect of coaching support on kidney function in chronic kidney disease patients. *Open Access Maced J Med Sci.* 2021;9(T4):106-10. Available from: <https://www.oamjms.eu/index.php/mjms/article/view/5820> [Last accessed on 2021 Jul 05].
- Susanti S, Bistara DN. Relationship between diet and RBG levels in diabetes mellitus patients. *J Vocat Health.* 2018;3(1):29-34.

# Paper 12

---

## ORIGINALITY REPORT

---

18%

SIMILARITY INDEX

16%

INTERNET SOURCES

13%

PUBLICATIONS

8%

STUDENT PAPERS

---

## PRIMARY SOURCES

---

1	<a href="http://journal2.unusa.ac.id">journal2.unusa.ac.id</a> Internet Source	5%
2	Siti Nur Hasina, Iis Noventi, P.H. Livana, Dodik Hartono. "Mindfulness Meditation Based on Spiritual Care to Reduce Community Anxiety due to the Impact of Pandemic Coronavirus Disease", Open Access Macedonian Journal of Medical Sciences, 2021 Publication	2%
3	<a href="http://pasca.uns.ac.id">pasca.uns.ac.id</a> Internet Source	1%
4	Submitted to Universitas Nahdlatul Ulama Surabaya Student Paper	1%
5	<a href="http://elearning.medistra.ac.id">elearning.medistra.ac.id</a> Internet Source	1%
6	<a href="http://eprints.poltekkesjogja.ac.id">eprints.poltekkesjogja.ac.id</a> Internet Source	1%
7	Submitted to University of Glasgow Student Paper	1%

---



8	garuda.ristekbrin.go.id Internet Source	1 %
9	balimedicaljournal.org Internet Source	1 %
10	Atik Badi'ah, Ni Ketut Mendri, Bondan Palestin, Amin Subargus et al. "Family Empowerment Psychoeducation on Family Support Caring of Children Diarrhea", Open Access Macedonian Journal of Medical Sciences, 2022 Publication	1 %
11	worldwidescience.org Internet Source	1 %
12	www.scitepress.org Internet Source	1 %
13	Izabela Michońska, Edyta Łuszczki, Magdalena Zielińska, Łukasz Oleksy, Artur Stolarczyk, Katarzyna Dereń. "Nutritional Programming: History, Hypotheses, and the Role of Prenatal Factors in the Prevention of Metabolic Diseases—A Narrative Review", Nutrients, 2022 Publication	1 %
14	Sani Nurlaela Fitriansyah, Rika Hartati, Irda Fidrianny. "Effect of Different Solvent on Phytochemical Content, Tyrosinase Inhibition and Antioxidant Activities of Campolay	1 %

(Pouteria campechiana kunth. [Baehni.]",  
Open Access Macedonian Journal of Medical  
Sciences, 2022

Publication

---

15 [www.tandfonline.com](http://www.tandfonline.com) 1 %  
Internet Source

---

16 Susanti Susanti, Difran Nobel Bistara. "The  
Effect of Coaching Support on Kidney  
Function in Chronic Kidney Disease Patients",  
Open Access Macedonian Journal of Medical  
Sciences, 2021 1 %  
Publication

---

17 [erepo.unud.ac.id](http://erepo.unud.ac.id) 1 %  
Internet Source

---

18 [repository.unusa.ac.id](http://repository.unusa.ac.id) 1 %  
Internet Source

---

Exclude quotes On

Exclude matches < 1%

Exclude bibliography On