

SURAT KETERANGAN

Nomor: 750/UNUSA/Adm-LPPM/VII/2020

Lembaga Penelitian dan Pengabdian Kepada Masyarakat (LPPM) Universitas Nahdlatul Ulama Surabaya menerangkan telah selesai melakukan pemeriksaan duplikasi dengan membandingkan artikel-artikel lain menggunakan perangkat lunak **Turnitin** pada tanggal 06 Juli 2020.

Judul : *Evaluation Of Erythrocyte Sedimentation Rate (Esr) Of Motorcycle Workshop Worker Exposed To Benzene*

Penulis : Moch. Sahri, Maharani Pertiwi Koentjoro, Shamsul Bahari
Shamsudin

No. Pemeriksaan : 2020.07.09.340

Dengan Hasil sebagai Berikut:

Tingkat Kesamaan diseluruh artikel (*Similarity Index*) yaitu 24%

Demikian surat keterangan ini dibuat untuk digunakan sebagaimana mestinya.

Surabaya, 09 Juli 2020

a.n. Ketua LPPM
Sekretaris LPPM



UNUSA
LPPM

Dr. Ubaidillah Zuhdi, S.T., M.Eng., M.S.M.
NPP: 18101208

LPPM Universitas Nahdlatul Ulama Surabaya

Website : lppm.unusa.ac.id

Email : lppm@unusa.ac.id

Hotline : 0838.5706.3867

2

by 2 2

Submission date: 18-Oct-2019 06:28PM (UTC+0700)

Submission ID: 1195425838

File name: 2_Moch_Sahri.pdf (710.76K)

Word count: 2329

Character count: 12752

Evaluation of Erythrocyte Sedimentation Rate (ESR) of Workshop Motorcycle Workshop Worker Exposed to Benzene

Moch. Sahri¹, Maharani Pertiwi Koentjoro², Shamsul Bahari Shamsudin³

¹Study Program of Occupational Health and Safety, Faculty of Health, Nahdlatul Ulama University of Surabaya, Indonesia

²Study Program of Medical Laboratory Technology, Faculty of Health, Nahdlatul Ulama University of Surabaya, Indonesia

³Departement of Community Health, Faculty of Medicine and Health Science, University Putra Malaysia, Malaysia

Correspondence:

³Maharani Pertiwi Koentjoro,
Jl. Jemursari No. 51-57, Surabaya,
East Java, Indonesia
Zip Code : 60237

Email: maharani@unusa.ac.id

Received: September 9, 2019

Revised: October 7, 2019

Accepted: October 12, 2019



Abstract

Occupational exposure to benzene from fuel oil in workshop motorcycle workshop causes several health effects, depending upon the level and duration of exposure. Analysis of blood is important to assess the status of worker's health. The study aimed to assess the gather basic information required for protecting workers' health and improve working conditions during works sites by investigating the ESR levels. A comparative cross-sectional study conducted in Surabaya City, East Java-Indonesia, which involved 100 workers. The occupational data collected using a structured questionnaire while blood analysis parameters measured with an automated hematology analyzer. The result showed that leukocytes, erythrocytes, hemoglobin (HGB), hematocrit values (HCT), platelet, MCV, RDW, eosinophil, basophils, neutrophil rods, neutrophil segment, lymphocyte, monocytes were in normal value. Whereas, MCH, MCHC, ESR has shown in abnormal value as average 24.5 pg, 29.18 g/dL and 8.21 mm/h, respectively. Those value, especially in ESR value indicated increasing concentration plasma viscosity. This is may cause by inflammation in the workers' body. To prevent the hazardous effect of benzene exposure, occupational health should be surveillance for workers in order to protect them from exposure to benzene.

Keywords

Workshop motorcycle workshop workers, ESR, Surabaya city, benzene

INTRODUCTION

The level of benzene exposure in the workshop motorcycle workshop during operations may occur. Mechanical workshop sites process various chemical products based

on highly volatile and flammable liquid hydrocarbons (1, 2), namely Premium, Pertamina, Solar which are the most widely used types of fuel oil. It presents many dangers associated with toxic chemical

materials, including carcinogens such as benzene. However, recent reports have indicated a link between occupational diseases such as cancer and benzene exposure in workshop site (2). Workers in sites exposure to benzene are more risk in occupational health (1, 3).

Erythrocyte Sedimentation Rate (ESR) is a blood test that measures a “sed rate”. This rate is depends on various factors, namely hemoglobin concentration, the ratio of plasma proteins, serum lipid concentration, and plasma pH. It has been a widely used marker for inflammation and indicates disease (5). In the occupational health of fuel oil, ESR determination is an important part of critical and emergent diagnoses such acute and chronic lymphocytic leukemia; non-Hodgkin’s lymphoma, multiple myeloma, anemia due to hematopoietic organs dysfunction, malignancy, temporal arteritis, renal disease, collagen vascular disease (6) (7). Furthermore, the role of ESR in the clinical management of various disease (6).

Routine tasks in the workshop motorcycle workshop include such as replacing spark plugs or changing the oil, repair engines and transmissions, removing dents from fenders. Thus, evaluation of the effect of benzene exposure is one of occupational health requirement at workshop motorcycle workshop (8). This study aims to gather basic information required for protecting workers’ health and improve

working conditions during works sites by investigating the hematological parameters, especially in the ESR levels.

MATERIALS AND METHODS

Study design

A comparative cross-sectional study of workers conducted at a workshop motorcycle workshop. Study subject involved 100 occupationally workshop motorcycle workers exposed fuel oil. All worker requiring hematological parameters evaluation, especially in ESR levels. All workers were asked to give informed consent.

Data collection

Demographic and occupational data were collected using a structured questionnaire, and subjects interviewed by the researcher. Before the blood collection, resting blood pressure was measured by using sphygmomanometer. Blood samples from worker respondents were taken at the end of the work shift. 4 mL blood samples were collected using venipuncture techniques into a Vacutainer tube containing potassium EDTA from the antecubital area of participants. The tubes were then placed in a holder or rack and placed/stored undisturbed in cold box. All samples were analyzed using an automated hematology analyzer Horiba Type ABX Micros 60 (KX-21N; Sysmex, Kobe, Japan).

Data processing and analysis

All the data were clean, double entered into Microsoft Excel spreadsheets and analyzed using Stata software version 13 (StataCorp, College Station, TX, USA).

Ethical considerations

The proposal in this study was approved by the Surabaya occupational safety and health technical implementation unit. All respondents in this study gave written agreements to participate in this study. All respondents were given an explanation of the purpose and results of this study. Laboratory test results were given to respondents in this study.

RESULTS

Table 2. Major Activities of The Workshop Motorcycle Workers

Job	Task
Mechanical engineer	1. Dismantle engines and repair or replace defective parts, such as magnetos, carburetors, and generators.
	2. Remove cylinder heads, grind valves, and scrape off carbon, and replace defective valves, pistons, cylinders and rings, using hand tools and power tools.
	3. Hammer out dents and bends in frames, weld tears and breaks; then reassemble frames and reinstall engines.
	4. Repair or replace other parts, such as headlights, horns, handlebar controls, gasoline and oil tanks, starters, and mufflers.
	5. Repair and adjust motorcycle subassemblies such as forks, transmissions, brakes, and drive chains, according to specifications.

¹ General characteristics

Personal characteristics of the workshop motorcycle workers are presented in Table 1. They were comparable with regard to age and sex. 91% of workers were males. The average age of the workers was 28 years. Protective measures were poorly followed during work in workshop.

Table 1. Personal Characteristics of Workshop Motorcycle Workers

Age Group	Range	Percentage (%)
Age-group (years)	18-27	55
	28-37	33
	38-53	12
Sex	Male	9
	Female	91



Fig 1. Partial view of working practice in of the workshop motorcycle. Workers are engaged in routine activities without proper personal protective equipment (mask and gloves)

Based on this study, the possible routes of chemical entry into the body through inhalation. It has shown in Figure 1 that the workers did not wear a mask during workpractice

Table 3. Blood Analysis and Hematological Parameters of Workshop Motorcycle Workers

Parameters	Average value	Normal (%)	Abnormal (%)
Leukocytes	8.53 x10 ³ /mm ³	80	20
Erythrocytes	5.07 x 10 ⁶ /mm ³	93	7
Hemoglobin (HGB)	12.51 g/dL	96	4
Hematocrit values (HCT)	43.26 %	98	2
Platelet	298.96	91	9
MCV	84.36 μm ³	86	14
MCH	24.5 pg	11	89
MCHC	29.18 g/dL	7	93
RDW	13.62 %	96	4
Eosinofil	1 %	100	0
Basophils	0 %	100	0
Natrofil rod	0.65 %	100	0
Neutrophil Segment	63.94 %	95	5
Lymphocyte	30.99 %	98	2
Monocytes	3.44 %	35	65
ESR	8.21 mm/h	8	92

MCV (Mean Corpuscular Volume); MCH (Mean Corpuscular Hemoglobin); MCHC (Mean Corpuscular Hemoglobin Concentration); RDW (RBC Distribution Width); ESR (Erythrocyte Sedimentation Rate).

DISCUSSION

The results presented here are based on the study in workshop motorcycle worker to evaluate the ESR value during benzene exposed in their worksites. Workshop motorcycle workers are exposed to different chemicals of fuel oil in their workshop. Studies have found that they can be exposed

to numerous types of heavy metals like lead, chromium, and cadmium, including carcinogens such as benzene (1, 9). Benzene was induced poisoning occurs almost entirely by inhalation of benzene vapors in the air.

Workshop motorcycle workers in our study exposed to benzene showed mean RBCs (average value is 5.07 10⁶/mm³) and

hemoglobin (12.5 g/dL) were in the normal range. RBC indicates MCV, MCH and MCHC which are calculated from hemoglobin, hematocrit, and erythrocytes count. Normal MCV indicates normocytic (normal average RBC size). Mean cell hemoglobin concentration (MCHC) refers to the average concentration of hemoglobin in the RBCs contained within the sample. Abnormally low MCHC in workshop motorcycle workers is called hypochromic. Mean cell hemoglobin (MCH) refers to the average weight of hemoglobin in the RBCs in the sample (units are picograms (pg/cell)).

Among workshop motorcycle worker, the ESR abnormal rate is as much as 92%. It is suggested that some inflammation reaction in the body (9). The ESR value indicated the increasing concentration of fibrinogen, the main clotting protein and alpha globulin (10). This is lead to plasma viscosity increased and red blood cell start to aggregate (10, 11).

We assumed that raised ESR is observed in workshop motorcycle workers caused by benzene in fuel oil exposure. It is well known that benzene causing haematological toxins, such as aplastic anemia and leukemia. Recent studies demonstrated that low-level benzene exposure (<1 ppm) has potential in hepatotoxicity (12).

Benzene is slightly soluble in water. Hence, this is easy to circulating blood enters the tissue and fat tissue. The primary route of benzene exposure is entering the body from

inhalation (13). This is caused by benzene very easy to evaporate into the air very quickly. Benzene will enter into the bloodstream, and travel into the entire organ. In temporally, benzene can be stored in the bone marrow and fat. The metabolism benzene includes in oxidations of the benzene rings by cytochrome P450 monooxygenase. This enzyme is primarily produced in liver cells. Therefore, accumulation of benzene in long periodic could affect liver function. Normally, benzene is degraded into phenol (14). Further, phenol is absorbed into the blood and distributed to other tissue.

Short term exposure to fuel oil, especially in benzene results in neurotoxic, irritatory, reproductive and developmental effects (15). Long-term exposure to benzene may cause a hazardous effect on health, such as hematopoietic cancer.

Evaluation of hematological parameters, including ESR is useful for monitoring the exposure of workshop motorcycle workers. This exposure carries a serious risk. Therefore, it is important to keel fuel oil securely from workshop motorcycle workers.

This study has some limitations. We evaluated the hematological parameters, namely ESR in workshop motorcycle workers exposed to fuel oil. We did not measure the specific chemicals to which the workshop motorcycle workers were exposed. It would be better to explore the effect of types of occupational chemical exposure on

health in future studies.

CONCLUSIONS

Benzene contamination in the air may cause reduced MCH and MCHC level and ESR. It was possible that the workers' exposure to benzene in fuel oil during work. The advice that can be given in this research is to reduce benzene exposure should install exhaust ventilation and workers using

personal protective equipment such as gas masks and gloves.

ACKNOWLEDGEMENTS

We would like to acknowledge all the workshop motorcycle workers who were willing to be respondents in this study.

3

CONFLICT OF INTEREST

There are no conflicts of interest.

REFERENCES

1. Chung E-K, Shin J-A, Lee B-K, Kwon J, Lee N, Chung K-J, et al. Characteristics of Occupational Exposure to Benzene during Turnaround in the Petrochemical Industries. *Saf Health Work*. 2010 Sep;1(1):51–60.
2. Kamal A, Rashid A. Benzene exposure among auto-repair workers from workplace ambience: A pioneer study from Pakistan. *Int J Occup Med Environ Health*. 2014 Oct;27(5):830–9.
3. Chaiklieng S, Suggaravetsiri P, Autrup H. Risk Assessment on Benzene Exposure among Gasoline Station Workers. *Int J Environ Res Public Health*. 2019 Jul 16;16(14).
4. Hashemi R, Majidi A, Motamed H, Amini A, Najari F, Tabatabaey A. Erythrocyte Sedimentation Rate Measurement Using as a Rapid Alternative to the Westergren Method. :5.
5. Siemons L, ten Klooster PM, Vonkeman HE, van Riel PL, Glas CA, van de Laar MA. How age and sex affect the erythrocyte sedimentation rate and C-reactive protein in early rheumatoid arthritis. *BMC Musculoskelet Disord*. 2014 Dec;15(1):368.
6. Bray C, Bell LN, Liang H, Haykal R, Kaikow F, Mazza JJ, et al. Erythrocyte Sedimentation Rate and C-reactive Protein Measurements and Their Relevance in Clinical Medicine. 2016;115(6):6.
7. Salehiforouz B, Vahdati A, Malekirad AA, Edalatmanesh MA. Evaluation of hematological indices of workers exposed to benzene. 2017;15:41–9.
8. Tualeka AR, Salesman F, Jalaluddin J, Wahyu A. Safe Concentration of Benzene Exposure in Work Environment at Motor Workshop. *Glob J Health Sci*. 2018 Oct 25;10(11):p188.
9. Salemcity A. Toxicological Assessment Of Consumption Of Benzene-Polluted Water On The Heart And Blood Of Albino Rats. [cited 2019 Aug 28]; Available from: https://www.academia.edu/6439376/Toxicological_Assessment_Of_Consumption_Of_Benzene-Polluted_Water_On_The_Heart_And_Blood_Of_Albino_Rats
10. Harrison M. Erythrocyte sedimentation rate and C-reactive protein. *Aust Prescr*. 2015 Jun;38(3):93–4.
11. Yin W, Xu Z, Sheng J, Xie X, Zhang C. Erythrocyte sedimentation rate and fibrinogen concentration of whole blood influences the cellular composition of platelet-rich plasma obtained from centrifugation methods. *Exp Ther Med*. 2017 Sep;14(3):1909–18.
12. Koh D-H, Jeon H-K, Lee S-G, Ryu H-W. The relationship between low-level benzene exposure and blood cell counts in Korean workers. *Occup Environ Med*. 2015 Jun;72(6):421–7.
13. Brekalo-Lazarevic S, Begic A, Ademovic Z, Horozic E. Determination Of Benzene Metabolite Phenol In The Urine And Analysis Of Blood Parameters Of Workers Exposed To Benzene. :6.
14. Brekalo-Lazarevic S, Begic A, Ademovic Z, Horozic E. Determination Of Benzene Metabolite Phenol In The Urine And Analysis Of Blood Parameters Of Workers Exposed To Benzene. :6.
15. Tuomi T, Veijalainen H, Santonen T. Managing Exposure to Benzene and Total Petroleum Hydrocarbons at Two Oil Refineries 1977-2014. *Int J Environ Res Public Health*. 2018 24;15(2).

ORIGINALITY REPORT

24%

SIMILARITY INDEX

24%

INTERNET SOURCES

18%

PUBLICATIONS

7%

STUDENT PAPERS

PRIMARY SOURCES

1**www.dovepress.com**

Internet Source

11%**2****www.ncbi.nlm.nih.gov**

Internet Source

5%**3****repository.unusa.ac.id**

Internet Source

4%**4****www.ajol.info**

Internet Source

3%Exclude quotes OnExclude bibliography OnExclude matches < 2%