

SURAT KETERANGAN

Nomor: 219/UNUSA/Adm-LPPM/III/2020

Lembaga Penelitian dan Pengabdian Kepada Masyarakat (LPPM) Universitas Nahdlatul Ulama Surabaya menerangkan telah selesai melakukan pemeriksaan duplikasi dengan membandingkan artikel-artikel lain menggunakan perangkat lunak **Turnitin** pada tanggal 27 Februari 2020.

Judul : Effect of puerperium intensive care on willingness of exclusive breast feeding
Penulis : Yunik Windarti dan Uke Maharani
Identitas : International Conference Public Health Solo, Oktober 2019
No. Pemeriksaan : 2020.03.03.124

Dengan Hasil sebagai Berikut:

Tingkat Kesamaan diseluruh artikel (*Similarity Index*) yaitu 17%

Demikian surat keterangan ini dibuat untuk digunakan sebagaimana mestinya

Surabaya, 03 Maret 2020

Ketua LPPM,



Dr. Istas Pratomo, S.T., M.T.

NPP. 16081074

LPPM Universitas Nahdlatul Ulama Surabaya

Website : lppm.unusa.ac.id

Email : lppm@unusa.ac.id

Hotline : 0838.5706.3867

Effect of Puerperium Intensive Care

by Yunik Windarti

Submission date: 19-Mar-2020 02:00PM (UTC+0700)

Submission ID: 1278025344

File name: e_20care_20on_20willingness_20of_20exclusive_20breastfeeding.pdf (147.75K)

Word count: 376

Character count: 2109

EFFECT OF PUERPERIUM INTENSIVE CARE ON THE WILLINGNESS TO PROVIDE EXCLUSIVE BREASTFEEDING

Yunik Windarti,² Uke Maharani Dewi

Faculty of Nursing and Midwifery, Universitas Nahdlatul Ulama, Surabaya

ABSTRACT

³**Background:** The puerperium period refers to the time after delivery when maternal physiological changes related to pregnancy return to the nonpregnant state. In puerperal period the nursing staff has an important educational role to the care that the woman in puerperium must have with herself and with the baby. There are factors involved in the difficulty of implementing infant care, particularly exclusive breastfeeding (EBF) and early interruption of breastfeeding. This study aimed to determine the effect ¹ puerperium intensive care on the willingness to provide EBF.

Subjects and Method: This was a randomized controlled trial conducted in Wono-kromo, Surabaya, East Java. A sample of 80 postpartum mothers was selected for this study, and randomly allocated into 2 groups: 40 mothers in the puerperium intensive care group and 40 mothers in the control group. The dependent variable was the willingness to provide EBF. The independent variable (intervention) was puerperium intensive care. Willingness to provide EBF was measured by a questionnaire. Percent difference in the willingness to provide the EBF between the two groups was tested by chi-square, with risk ratio (RR) as the measure of effect.

Results: Before the intervention, the two groups were comparable in term of percent of mothers willing to provide EBF, indicating the success of randomization. After the intervention, percent of mothers willing to provide EBF in the intervention group (57.5%) was higher than the control group (45%) with RR=1.28, and it was statistically significant ($p=0.019$). RR=1.28 means that postpartum mothers who received the puerperium intensive care were 1.28 times as likely to provide EBF than those who did not receive the intervention.

Conclusion: Puerperium intensive care is effective to enhance the willingness to provide EBF among postpartum mothers.

Keywords: puerperium intensive care, breastfeeding

²Correspondence:

Yunik Windarti. Faculty of Nursing and Midwifery, Universitas Nahdlatul Ulama, Surabaya, East Java. Email: yunikwinda@unusa.ac.id. Mobile: 08133033090.

Effect of Puerperium Intensive Care

ORIGINALITY REPORT

17 %	11 %	0 %	11 %
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

PRIMARY SOURCES

1	media.neliti.com Internet Source	7 %
2	Submitted to iGroup Student Paper	6 %
3	Submitted to Melbourne Institute of Business and Technology Student Paper	4 %

Exclude quotes On

Exclude bibliography On

Exclude matches < 5 words