

## **SURAT KETERANGAN**

Nomor: 549/UNUSA/Adm-LPPM/VI/2020

Lembaga Penelitian dan Pengabdian Kepada Masyarakat (LPPM) Universitas Nahdlatul Ulama Surabaya menerangkan telah selesai melakukan pemeriksaan duplikasi dengan membandingkan artikel-artikel lain menggunakan perangkat lunak **Turnitin** pada tanggal 14 Juni 2020

Judul : Pharmacologic Aspect of Neuropathic Pain  
Penulis : Endang Mutiawati, Imai Indra, Syahrul, Mulyadi  
No. Pemeriksaan : 2020.06.14.285

Dengan Hasil sebagai Berikut:

**Tingkat Kesamaan diseluruh artikel (*Similarity Index*) yaitu 25%**

Demikian surat keterangan ini dibuat untuk digunakan sebagaimana mestinya

Surabaya, 14 Juni 2020

Ketua LPPM,



Dr. Istas Pratomo, S.T., M.T.

NPP. 16081074

**LPPM Universitas Nahdlatul Ulama Surabaya**

Website : [lppm.unusa.ac.id](http://lppm.unusa.ac.id)

Email : [lppm@unusa.ac.id](mailto:lppm@unusa.ac.id)

Hotline : 0838.5706.3867

# Paper 7

*by* Mulyadi 7

---

**Submission date:** 14-Jun-2020 01:41PM (UTC+0700)

**Submission ID:** 1343379029

**File name:** c.\_Pharmacologic-aspect-of-neuropathic-pain.pdf (387.37K)

**Word count:** 3303

**Character count:** 19900

## Pharmacologic Aspect of Neuropathic Pain

<sup>1</sup>Endang Mutiawati, <sup>2</sup>Imai Indra, <sup>3</sup>Syahrul, <sup>4</sup>\*Mulyadi

22

<sup>1</sup>Department of Neurology, Faculty of Medicine, Syiah Kuala University, Darussalam, Banda Aceh 23111, Indonesia;

<sup>2</sup>Department of anesthesiology, Medical Faculty, Syiah Kuala University, Banda Aceh 23111, Indonesia;

<sup>3</sup>Faculty of Medicine, Syiah Kuala University, Darussalam, Banda Aceh 23111, Indonesia;

<sup>4</sup>Department of Pulmonology and Respiratory Medicine, Faculty of Medicine, Syiah Kuala University, Darussalam, Banda Aceh 23111, Indonesia;

Corresponding author : mul.0862@gmail.com

### Abstract

Neuropathic pain is pain arising from nerve damage to the conductive pathways of pain (ranging from nociceptors to post central gyrus). Neuropathic pain can be caused by 1) Carcinomas, 2) Trauma/compressive, 3) Congenital, 4) Immunomediated, 5) Infection, 6) Metabolic disorders, 7) nutritional deficiency, 8) Toxin, 9) Lesion, 10) Vasculitis, 11) Connective tissue disorders. To date, the pathophysiology of neuropathic pain can not be explained thoroughly, this problem leads to the treatment which has not given satisfactory results as expected. There are many types of drugs has been used for the treatment of neuropathic pain, and they are generally aimed to stop the flow of impulses in the nervous system which was activated as a result of ectopic generators in areas experiencing nerve injury. These drugs work in several locations such as: drugs that works on 'sodium channel voltage gate' (i.e Carbamazepine group), drugs that works on 'calcium channel' (i.e Gabapentin and Pregabalin), and also drugs that works on 'the synapses gap' (i.e Tricyclic class). Besides drugs that inhibit pain impulses propagation, the treatment of neuropathic pain also include drugs that have the ability of nervous system regeneration such as methylcobalamin group. The rationale of the use of this kind of drugs is that this drug expected to regenerate the damage of the nervous system damage which is lead to decrease the ectopic generator activity, the end result is the reduction of neuropathic pain experienced by patients.

**Key words:** neuropathic pain, pharmacologic, regeneratif

### Introduction

Neuropathic pain is common, greatly impairs quality of life and has a high economic impact on society: the Institute of Medicine reports that at least 116 million American adults suffer from chronic pain, and estimates for people suffering from neuropathic pain are as high as 17.9%. Co-morbidities such as poor sleep, depression and anxiety are common in neuropathic pain patients, leading to unresolved arguments about whether pain causes mood and sleep changes or whether individuals with mood and sleep disorders are at a higher risk of developing pain (Hehn *et al.*, 2012). Neuropathic pain (NP) is estimated to afflict as high as 7–8% of the general population in Europe (Attal *et al.*, 2010), an estimated prevalence of 5% to 7% in France, compared with 20% to 31% for chronic pain (Delorme *et al.*, 2011) and It is a common condition with an overall prevalence between 0.9 and 8.0% (Athanasakis *et al.*, 2013).

17

The International Association for the Study of Pain (IASP) defines neuropathic pain (NP) as pain "initiated or caused by a primary lesion or dysfunction in the nervous system". It is estimated to afflict millions of people worldwide, although precise figures are not available. Neuropathic pain can be caused by a number of different diseases (e.g., diabetes mellitus, herpes zoster, human immunodeficiency virus [HIV] infection), medical interventions (e.g., chemotherapy, surgery), and injuries (e.g., brachial plexus avulsion) causing NP by producing lesions in somatosensory pathways in the peripheral or central nervous system (Dworkin *et al.*, 2007; O'Connor & Dworkin, 2009). The management of patients with NP is complex and response to existing treatments is often inadequate. Even with well-established NP medications, effectiveness is unpredictable, dosing can be complicated, analgesic onset is delayed, and side effects are common (Dworkin *et al.*, 2007; Attal *et al.*, 2010).

### Materials and Methods

This article review form articles related with the pharmacologic aspect of neuropathic pain. All data contained in this article is the result of a number of articles search.

### Results and Discussion

In 2006, the European Federation of Neurological Societies (EFNS) produced the first guidelines on pharmacological treatment of NP. Since 2006, new randomized controlled trials (RCTs) have appeared in various NP conditions, justifying an update (Dworkin *et al.*, 2007; Attal *et al.*, 2010).

Pharmacological treatment is the mainstream in post-herpetic neuralgia (PHN), diabetic peripheral neuropathic pain (DPNP), central post-stroke pain (CPSP), trigeminal neuralgia (TN), complex regional pain syndrome (CRPS), cancer pain, failed back syndrome etc, while polypharmacy is still the major prescriptions facing such kind of miserable patients. The tricyclic antidepressants (TCA), gamma-aminobutyric acid (GABA), voltage-dependent calcium channel blockers, selective norepinephrine reuptake inhibitor (SNRI), opioid or morphine etc, are still evidence-based medicines (EBM) but with different outcome for individuals (Yang *et al.*, 2012). Use of antiepileptics with demonstrated efficacy as first-line therapy has increased in neuropathic pain (Hall *et al.*, 2013).

16

Table 1. Stepwise Pharmacologic Management of Neuropathic Pain (O'Connor & Dworkin, 2009)

- Step 1
- Assess pain and establish the diagnosis of NP, if uncertain about the diagnosis, refer to a pain specialist or neurologist
  - Establish and treat the cause of NP; if uncertain about availability of treatment addressing NP etiology, refer to appropriate specialist
  - Identify relevant comorbidities (e.g., cardiac, renal, or hepatic disease, depression, gait instability) that might be relieved or exacerbated by NP treatment, or that might require dosage adjustment or additional monitoring of therapy
  - Explain the diagnosis and treatment plan to the patient, and establish realistic expectations
- Step 2
- Initiate therapy of the disease causing NP, if applicable
  - Initiate symptom treatment with one or more of the following:
    - ✓ Antidepressant medication: either secondary amine TCA (nortriptyline, desipramin) or SSNRI (duloxetine, venlafaxine)
    - ✓ Calcium channel  $\alpha_2\delta$  ligand: either gabapentin or pregabalin
    - ✓ For patients with localized peripheral NP: topical lidocaine used alone or in combination with 1 of the other first-line therapies
    - ✓ For patients with acute NP, neuropathic cancer pain, or episodic exacerbations of severe pain, and when prompt pain relief during titration of a first-line medication to an efficacious dosage is required, opioid analgesics or tramadol may be used alone or in combination with 1 of the first-line therapies.
  - Evaluate patient for nonpharmacologic treatments, and initiate if appropriate
- Step 3
- Reassess pain and health-related quality of life frequently
  - If substantial pain relief (e.g., average pain reduced to NRS  $\leq 3/10$ ) and tolerable side effects, continue treatment.
  - If partial pain relief (e.g., average pain remains  $\geq 4/10$ ) after an adequate trial, add 1 of the other first-line medications
  - If no or inadequate pain relief (e.g.,  $<30\%$  reduction) at target dosage after an adequate trial, switch to an alternative first-line medication
- Step 4
- If trials of first-line medications alone and in combination fail, consider second-line medications or referral to a pain specialist or multidisciplinary pain Center

NP-neuropathic pain; NRS-numeric rating scale; SSNRI-selective serotonin and norepinephrine reuptake inhibitor; TCA-tricyclic antidepressant. Reprinted with permission from Pain.

20

Three classes of medications were recommended as firstline treatments: antidepressants with both norepinephrine and serotonin reuptake inhibition (SSNRIs) and selective serotonin and norepinephrine reuptake inhibitors [SSNRIs]), calcium channel 2- ligands (gabapentin and pregabalin), and topical lidocaine (lidocaine patch 5%). Opioids and tramadol were recommended as generally second-line treatments, except in certain specific clinical situations in which it was recommended that first-line use could be considered. A number of medications were considered third-line choices (O'Connor & Dworkin, 2009). Based on efficacy and safety, pregabalin is considered a first-line drug together with gabapentin in the treatment of central pain (Finnerup & Jensen, 2007).

The guidelines acknowledge that a combination of medications with efficacy for neuropathic pain may provide greater analgesia than use of individual medications as monotherapy, although such combination therapy will often be associated with increased side effects, inconvenience, risk of drug interactions, and cost. Nevertheless, because 50% of patients in Neuropathic Pain trials of efficacious medications typically achieve satisfactory pain relief, many patients in clinical practice will require treatment with a combination of medications. Such combination therapy was incorporated into a

stepwise management strategy for patients with partial responses to treatment with first-line medications (O'Connor & Dworkin, 2009).

Table 2. Summary of the results of Randomized Clinical Trials Involving First and Second-Line Medications for Patients With Neuropathic Pain (O'Connor & Dworkin, 2009)

	TCA	Dul oxet ine	Venl afaxi ne	Gaba pen tin	Preg aba lin	Topical lidocaine 5% ic	Opioid analges dol	Tra ma
Peripheral NP								
- Painful DPN	(+)	(+)	(+)	Both	Both	-	(+)	(+)
- PHN	(+)	-	(-)	(+)	Both	(+)	(+)	(+)
- Painful polyneuropathy	(+)	-	(+)	(+)	-	(+)	(+)	(+)
- Phantom limb pain	(-)	-	-	Both	-	-	(+)	(+)
- Postmastectomy pain	(+)	-	(-)	-	-	-	-	-
- GBS	-	-	-	(+)	-	-	-	-
- Neuropathic cancer pain	(-)	-	-	(+)	-	-	-	-
- CPRS type I	-	-	-	(-)	-	-	-	-
- Chronic lumbal root pain	(-)	-	-	-	-	-	(-)	-
- Chemotherapy induced neuropathy	(-)	-	-	(-)	-	-	-	-
- HIV Neuropathy	(-)	-	-	(-)	-	-	-	-
Central NP								
- Central poststroke pain	(+)	-	-	-	(+)	-	-	-
- Spinal cord injury pain	(-)	-	(+)	(+)	-	-	-	-

The NeuPSIG guidelines note that few medications have been found to be efficacious in neuropathic pain originating from a lesion in the central nervous system. RCTs have demonstrated efficacy for TCAs in central poststroke pain, and for calcium channel  $\alpha_2$ -ligands in spinal cord injury and poststroke central neuropathic pain. The Canadian Pain Society created 4 levels of recommendation, with first- and second-line medications differentiated by "the quality of evidence and the evidence of efficacy" based on NNTs. Medications were classified as third-line treatments if they have good evidence of efficacy, but require specialized monitoring and follow-up not required of drugs at the other levels. Fourth-line medications were described as having "at least 1 positive RCT, but required further study" (O'Connor & Dworkin, 2009).

As with the NeuPSIG and Canadian Pain Society guidelines, the EFNS guidelines grade the level of evidence for different available treatments. However, unlike the other 2 sets of guidelines, separate recommendations were made for the treatment of patients with painful polyneuropathies (including painful diabetic peripheral neuropathy), postherpetic neuralgia, trigeminal neuralgia, and central neuropathic pain. Consistent with the NeuPSIG and Canadian Pain Society guidelines, the EFNS guidelines recommended gabapentin, pregabalin, and TCAs as first-line treatments for painful polyneuropathies, postherpetic neuralgia, and central neuropathic pain (Moulin *et al.*, 2007; O'Connor & Dworkin, 2009; Rhodes, 2011).

Table 3. Comparison of Neuropathic Pain Treatment Guidelines, Excluding Trigeminal Neuralgia (O'Connor & Dworkin, 2009).

Medication Class	NeuPSIG Guidelines	CPS Guidelines	EFNS Guidelines
Tricyclic antidepressants	First line	First line	First line for PPN, PHN, and CP
Calcium channel $\alpha_2$ -8 ligands (gabapentin and pregabalin)	First line	First line	First line for PPN, PHN, and CP
SSNRIs (duloxetine and venlafaxine)	First line	Second line	Second line for PPN
Topical lidocaine	First line for localized peripheral NP	Second line for localized peripheral NP	First line for PHN if small area of pain/allodynia
Opioid analgesics	Second line except in selected circumstances <sup>†</sup>	Third line	Second-third-line for PPN, PHN, and CP
Tramadol	Second line except in selected circumstances <sup>†</sup>	Third line	Second-third-line for PPN and PHN

CP = central pain; CPS = Canadian Pain Society; EFNS = European Federation of Neurological Societies; NeuPSIG = Neuropathic Pain Special Interest Group; NP = neuropathic pain; PHN = postherpetic neuralgia; PPN = painful polyneuropathy; SSNRIs = selective serotonin and norepinephrine reuptake inhibitors.

\*Only medications considered first or second-line in 1 of the guidelines are presented.

†Opioid analgesics and tramadol were considered first-line options in the following circumstances: for the treatment of acute NP, episodic exacerbations of severe NP, neuropathic cancer pain, and during titration of a first-line medication in patients with substantial pain.

Other EFNS recommendations for painful polyneuropathies were duloxetine and venlafaxine as second-line treatment ("because of moderate efficacy"), and opioids, tramadol, and lamotrigine as "second-/thirdline therapy." Additional recommendations for postherpetic neuralgia were topical lidocaine as a first-line treatment for patients with localized pain and allodynia, and opioids, tramadol, capsaicin, and valproic acid as second-line treatment options. "Second-/third-line" treatment options for patients with central neuropathic pain were lamotrigine, opioids, and cannabinoids (Moulin *et al.*, 2007; O'Connor & Dworkin, 2009; Rhodes, 2011).

Two drugs have been approved for neuropathic pain in the US – pregabalin and duloxetine – but neither of these afford complete relief, even when used in combination (Vinik & Cassellini, 2012). In addition, several novel drug treatments such as botulinum toxin, capsaicin patch and lacosamide have been also used for neuropathic pain therapy. Table 5 summarizes the major pharmacological treatments for neuropathic pain and their analgesia mechanisms (Xu *et al.*, 2012).

Table 4. Major pharmacological treatment for neuropathic pain and their basic mechanisms (Xu *et al.*, 2012)

Compound	Mode of action
<b>Antidepressant</b>	
- Nortriptyline	Inhibition of both serotonin and norepinephrine reuptake
- Desipramine	Inhibition of both serotonin and norepinephrine reuptake
- Duloxetine	Inhibition of both serotonin and norepinephrine reuptake
- Venlafaxine	Inhibition of both serotonin and norepinephrine reuptake
<b>Anticonvulsants</b>	
- Gabapentin	Decreases release of glutamate, norepinephrine and substance P, with ligands on $\alpha 2\text{-}\delta$ subunit of voltage
- Pregabalin	Decreases release of glutamate, norepinephrine and substance P, with ligands on $\alpha 2\text{-}\delta$ subunit of voltage
- Lacosamide	Decreases release of presynaptic transmitter inhibition of voltage-gated sodium channel
<b>Opioid agonists</b>	
- Morphine	$\mu$ -receptor agonism
- Oxycodone	$\mu$ -receptor agonism
- Methadone	$\mu$ -receptor agonism, $\kappa$ -receptor antagonism
- Levorphanol	$\mu$ -receptor agonism
- Tramadol	$\mu$ -receptor agonism, inhibition of norepinephrine and serotonin reuptake
<b>Topical therapy</b>	
- 5% lidocaine patch	Block of sodium channel
- High dose capsaicin patch	Damage of nociceptive sensory axons, a highly selective activating ligand for TRPV1
- Botulinum toxin	Inhibition of both the exocytosis of acetylcholine and some other neurotransmitter

Pregabalin reduces the enhanced noxious stimulus-induced spinal release of glutamate seen in neuropathic rats (Kumar *et al.*, 2010). Intranasal or intrathecal pregabalin relieves neuropathic pain behaviours, perhaps due to pregabalin's effect upon anterograde CaV $\alpha 2\delta$ -1 protein trafficking from the DRG to the dorsal horn. Intranasal delivery of agents such as pregabalin may be an attractive alternative to systemic therapy for management of neuropathic pain states (Martinez *et al.*, 2012). The anti-allodynic effects of gabapentin may be caused by upregulation of IL-10 expression in the spinal cord, which leads to inhibition of the expression of pro-inflammatory cytokines in the spinal cords (Lee *et al.*, 2013).

Capsaicin is a transient receptor potential vanilloid-1 agonist, which increases the intracellular calcium ion concentration. This triggers calcium-dependent protease enzymes causing cytoskeletal breakdown and leads to the loss of cellular integrity and 'defunctionalization' of nociceptor fibres. Efficacy and therapeutic effect has been shown in several clinical studies of PHN and HIV-DSP. The high-concentration capsaicin patch and its practical application are different from low-concentration creams; one application can last for up to 3 months (Irving *et al.*, 2011; Webster *et al.*, 2012; Baranidharan *et al.*, 2013). A single 30-minute application of NGX-4010 (Capsaicin 8% patch) provides significant pain relief for at least 12 weeks in patients with HIV-DSP and is well tolerated (Simpson *et al.*, 2008; Brown *et al.*, 2013).

Activation of TRPV1 by capsaicin results in sensory neuronal depolarization, and can induce local sensitization to activation by heat, acidosis, and endogenous agonists. Topical exposure to capsaicin leads to the sensations of heat, burning, stinging, or itching. High concentrations of capsaicin or repeated applications can produce a persistent local effect on cutaneous nociceptors, which is best

described as defunctionalization and constituted by reduced spontaneous activity and a loss of responsiveness to a wide range of sensory stimuli (Anand & Bley, 2011; Bley, 2013).

### Conclusions

All guidelines recommend TCAs, gabapentin, and pregabalin as first-line treatment options for patients with neuropathic pain (excluding trigeminal neuralgia). The NeuPSIG guidelines recommend duloxetine and venlafaxine as first-line treatment options, but the Canadian Pain Society and EFNS guidelines recommend these SSNRIs as second-line options for patients with painful polyneuropathies. Clinicians need to consider the advantage and disadvantage of these managements to avoid ineffective treatments, maximize curing proven beneficial in clinical trials, and minimize the side effect of therapies. To improve the current management of patients with neuropathic pain, evidence-based basic studies should be made for pharmacological approaches to guide the managements in the future.

10

### References

- Anand, P. & Bley, K., 2011. Topical Capsaicin for pain management: therapeutic potential and mechanisms of action of the new high-concentration capsaicin 8% patch. *British Journal of Anaesthesia*, 107(4): 490-502.
- Athanasakis, K., Petrakis, L., Karampli, E., Vitsou, E., Lyras, L., Kyriopoulos, J., 2013. Pregabalin versus Gabapentin in the Management of Peripheral Neuropathic Pain Associated with Post-herpetic Neuralgia and diabetic neuropathy: a cost effectiveness analysis for the Greek Healthcare setting. *BMC Neurology*, 13:56.
- Attal, N., Cruccu, G., Baron, R., Haanpaa, M., Hansson, P., Jensen, T.S., Nurmikko, T., 2010. EFNS Guidelines on the pharmacological treatment of neuropathic pain: 2010 revision. *European Journal of Neurology*, 17: 1113-1123.
- Baranidharan, G., Das, S., Bhaskar, A., 2013. A review of the high-concentration capsaicin patch and experience in its use in the management of neuropathic pain. *Therapeutic Advances in Neurological Disorders*, 6(5): 287-297.
- Bley, K., 2013. Effects of Topical Capsaicin on Cutaneous Innervation: Implications on Pain Management. *The Open Pain Journals*, 6: 81-94.
- Brown, S., Simpson, D.M., Moyle, G., Brew, B.J., Schifitto, G., Larbalestier, N., et al., 2013. NGX-4010, a Capsaicin 8% patch, for the treatment of the painful HIV-associated distal sensory polyneuropathy: integrated analysis of two phase III, randomised controlled trial. *AIDS Research and Therapy*, 10: 5.
- Delorme, C., Navez, M.L., Legout, V., Deleens, R., Moysse, D., 2011. Treatment of Neuropathic Pain with 5% lidocaine-medicated plaster: five years of clinical experience. *Pain Res Manage*, Vol 16, No 4.
- Dworkin, R.H., O'Connor, A.B., Backonja, M., Farrar, J.T., Finnerup, N.B., Jensen, T.S., et al., 2007. Pharmacologic Management of Neuropathic pain: Evidence-based Recommendations. *Pain*, 132: 237-251.
- Finnerup, N.B. & Jensen, T.S., 2007. Clinical use of pregabalin in the management of central neuropathic pain. *Neuropsychiatric Disease & Treatment*, 3(6): 885-891.
- Hall, G.C., Morant, S.V., Carroll, D., Gabriel, Z.L., McQuay, H.J., 2013. An Observational descriptive study of the epidemiology and treatment of neuropathic pain in a UK general population. *BMC Family Practice*, 14: 28.
- Hehn, C.A., Baron, R., Woolf, C.J., 2012. Deconstructing the Neuropathic pain Phenotype to Reveal Neural Mechanism. *Neuron*, 73(4): 638-652.
- Irving, G.A., Backonja, M., Rauck, R., Webster, L.R., Tobias, J.K., Vanhove, G.F., 2011. NGX-4010, a Capsaicin 8% dermal patch, administered alone or in combination with systemic neuropathic pain medications, reduces pain in patients with postherpetic neuralgia. *Clinical Journal Pain*.
- Kumar, N., Laferriere, A., Yu, J.S.C., Leavit, A., Coderre, T.J., 2010. Evidence that pregabalin reduces neuropathic pain by inhibiting the spinal release of glutamate. *Journal of Neurochemistry*, 113: 552-561.
- Lee, B., Jun, I., Kim, S., Park, J.Y., 2013. Intrathecal Gabapentin increases Interleukin-10 Expression and Inhibits pro-inflammatory Cytokines in a rat model of neuropathic pain. *J Korean Med Sci*, 28: 308-314.
- Martinez, J.A., Kasamatsu, M., Rosalez-Hernandez, A., Hanson, L.R., Frey, W.H., Toth, C.C., 2012. Comparison of central versus peripheral delivery of the pregabalin in neuropathic pain states. *Molecular Pain*, 8: 3.
- Moulin, D.E., Clark, A.J., Gilron, I., Ware, M.A., Watson, C.P.N., Sessle, B.J., et al., 2007. Pharmacological Management of chronic Neuropathic Pain – Consensus Statement and guidelines from the Canadian Pain Society. *Pain Res Manage*, Vol 12: No 1.
- O'Connor, A.B. & Dworkin, R.H., 2009. Treatment of Neuropathic Pain: An Overview of recent Guidelines. *The American Journal of Medicine*, 122: S22-S32.
- Presdes, C., 2011. Update on Therapies for Neuropathic Pain. *The Pain Practitioner*, Vol 21: No 3.
- Simpson, D.M., Brown, S., Tobias, J., 2008. Controlled trial of high-concentration capsaicin patch for the treatment of painful HIV neuropathy. *Neurology*, 70: 2305.
- Vinik, A.I. & Casellini, C.M., 2013. Guidelines in the management of diabetic nerve pain: clinical utility of pregabalin. *Diabetics, Metabolics syndrome and Obesity: Target and therapy*, 6: 57-78.
- Vranken, J.H., 2009. Mechanisms and Treatment of Neuropathic Pain. *Central Nervous System Agents in Medicinal Chemistry*, 9: 71-78.
- Webster, L.R., Peppin, J.F., Murphy, F.T., Tobias, J.K., Vanhove, G.F., 2012. Tolerability of NGX-4010, a capsaicin 8% patch, in conjunction with three topical anesthetic formulations for the treatment of neuropathic pain. *Journal of Pain Research*, 5: 7-13.
- Xu, B., Descalzi, G., Ye, H., Zhuo, M., Wang, Y., 2012. Translational investigation and treatment of neuropathic pain. *Molecular Pain*, 8: 15.
- Yang, C.M., Chen, N.C., Shen, H.C., Chou, C.H., Yeh, P.S., Lin, H.J., et al., 2012. Guideline of Neuropathic Pain Treatment and Dilemma from Neurological Point of View. *Acta Neurologica Taiwanica*, Vol 21: No 3.

# Paper 7

---

## ORIGINALITY REPORT

---

**25%**

SIMILARITY INDEX

**20%**

INTERNET SOURCES

**21%**

PUBLICATIONS

**15%**

STUDENT PAPERS

---

## PRIMARY SOURCES

---

- 1** Yang Zhang, Jinlin Wang, Li-Juan Ji, Lin Li, Meng Wei, Su Zhen, Cheng-Cai Wen. "Identification of Key Gene Modules of Neuropathic Pain by Co-Expression Analysis", *Journal of Cellular Biochemistry*, 2017  
Publication **1%**
  - 2** [scholarspace.manoa.hawaii.edu](http://scholarspace.manoa.hawaii.edu)  
Internet Source **1%**
  - 3** Submitted to Leeds Beckett University  
Student Paper **1%**
  - 4** Nisha S. Bhatia, Felicia C. Chow. "Neurologic Complications in Treated HIV-1 Infection", *Current Neurology and Neuroscience Reports*, 2016  
Publication **1%**
  - 5** [hepmp.med.bg.ac.rs](http://hepmp.med.bg.ac.rs)  
Internet Source **1%**
  - 6** Submitted to Nevada State College  
Student Paper **1%**
-



7	Submitted to University of Durham Student Paper	1%
8	m.koreamed.org Internet Source	1%
9	fjfsdata01prod.blob.core.windows.net Internet Source	1%
10	kiwiimporter.com Internet Source	1%
11	www.consultantlive.com Internet Source	1%
12	www.pnhelp.org Internet Source	1%
13	R Benoliel. "Peripheral painful traumatic trigeminal neuropathies : Peripheral painful traumatic trigeminal neuropathies", Oral Diseases, 05/2012 Publication	1%
14	Wiffen, Philip J, Sheena Derry, R Andrew Moore, Eija A Kalso, and Philip J Wiffen. "Carbamazepine for chronic neuropathic pain and fibromyalgia in adults", Cochrane Database of Systematic Reviews, 2014. Publication	1%
15	Widerstrom-Noga, Eva, Elizabeth Roy Felix, James P. Adcock, Maydelis Escalona, and	1%

Jacqueline Tibbett. "Multidimensional neuropathic pain phenotypes after spinal cord injury", Journal of Neurotrauma, 2015.

Publication

16

[www.ukessays.com](http://www.ukessays.com)

Internet Source

1%

17

Rafael Gálvez, Carlos Marsal, Javier Vidal, Manuel Ruiz, Javier Rejas. "Cross-sectional evaluation of patient functioning and health-related quality of life in patients with neuropathic pain under standard care conditions", European Journal of Pain, 2007

Publication

1%

18

"Pain and depression : The janus factor of human suffering", An Introduction to Pain and its Relation to Nervous System Disorders, 2016.

Publication

1%

19

[www.molecularpain.com](http://www.molecularpain.com)

Internet Source

1%

20

[whywoundcare.s3.amazonaws.com](http://whywoundcare.s3.amazonaws.com)

Internet Source

1%

21

[bpspubs.pericles-prod.literatumonline.com](http://bpspubs.pericles-prod.literatumonline.com)

Internet Source

1%

22

Submitted to Syiah Kuala University

Student Paper

1%

23	<a href="http://ir.uiowa.edu">ir.uiowa.edu</a> Internet Source	1%
24	<a href="http://www.researchsquare.com">www.researchsquare.com</a> Internet Source	1%
25	Submitted to Whitireia Community Polytechnic Student Paper	1%
26	Submitted to October University for Modern Sciences and Arts (MSA) Student Paper	1%
27	J D Markman. "Bundle Up: It's Painful Out There —The Case for Opioid–Acetaminophen Combinations", <i>Clinical Pharmacology &amp; Therapeutics</i> , 09/2010 Publication	1%
28	Pandhare, Ramdas B, B Sangameswaran, Popat B Mohite, and Shantaram G Khanage. "Attenuating effect of seeds of <i>Adenantha pavonina</i> aqueous extract in neuropathic pain in streptozotocin-induced diabetic rats: an evidence of neuroprotective effects", <i>Revista Brasileira de Farmacognosia</i> , 2012. Publication	1%
29	<a href="http://www.location-privee.eu">www.location-privee.eu</a> Internet Source	1%

---

Exclude quotes      On

Exclude bibliography      Off

Exclude matches      < 1%