

SURAT KETERANGAN

Nomor: 764/UNUSA/Adm-LPPM/VII/2020

Lembaga Penelitian dan Pengabdian Kepada Masyarakat (LPPM) Universitas Nahdlatul Ulama Surabaya menerangkan telah selesai melakukan pemeriksaan duplikasi dengan membandingkan artikel-artikel lain menggunakan perangkat lunak **Turnitin** pada tanggal 08 Juli 2020.

Judul : *Enzymatic Modification of Cotton Fiber For Promising Smart Medical Based Material*

Penulis : Maharani Pertiwi Koentjoro, Marisa Fitriana, Isdiantoni, Endry Nugroho Prasetyo

No. Pemeriksaan : 2020.07.14.349

Dengan Hasil sebagai Berikut:

Tingkat Kesamaan diseluruh artikel (*Similarity Index*) yaitu 14%

Demikian surat keterangan ini dibuat untuk digunakan sebagaimana mestinya.

Surabaya, 14 Juli 2020

a.n. Ketua LPPM

Sekretaris LPPM



UNUSA
LPPM

Dr. Ubaidillah Zuhdi, S.T., M.Eng., M.S.M.

NPP: 18101208

Enzymatic modification of cotton fiber for promising smart medical based material

by Maharani Koentjoro

Submission date: 07-Jul-2020 07:38AM (UTC+0700)

Submission ID: 1354332853

File name: n_of_cotton_fiber_for_promising_smart_medical_based_material.pdf (473.68K)

Word count: 4860

Character count: 26562

Enzymatic modification of cotton fiber for promising smart medical based material

Maharani Pertiwi Koentjoro
Study Program of Medical Laboratory
Technology
University of Nahdlatul Ulama
Surabaya
Surabaya, Indonesia
maharani@unusa.ac.id

Marisa Fitriana
Departement of Biology
Institute Technology of Sepuluh
Nopember
Surabaya, Indonesia
marisa.fitriana03@gmail.com

Isdiantoni
Faculty of Agriculture
University of Wiraraja
Sumenep, Indonesia
antonie_isd@yahoo.co.id

Endry Nugroho Prasetyo
Departement of Biology
Institute Technology of Sepuluh
Nopember
Surabaya, Indonesia
endry@bio.its.ac.id

Abstract — Color removal (bleaching) is an oxidation process whereby the undesirable inherent coloring component removal in organic material. Hydrogen peroxide (H_2O_2) is a fundamental stage of using bleaching agent in industrial and the medical field. As an oxidizing agent, H_2O_2 has merited a significant attention since it is able to degrade reactive dyes intensively. However, the H_2O_2 hydrolysis remain undesirable residues which leads a negative influence on fabric quality. Therefore, other alternatives method for bleaching is necessary by employing degradable materials like enzyme. Laccase is enzyme that considerable as a bio-bleaching agent for promising a smart medical based material. Therefore, in this study laccase from *Trametes versicolor* was used as bio-bleaching agents. The purpose of this study was to determine the potential of laccase for cotton fibers bio-bleaching. It was used H_2O_2 as control comparison meanwhile the laccase and laccase-ABTS (mediator) as treated groups. The brightness level and the presence of functional groups of cotton fiber were characterized by spectrophotometer and FTIR respectively. The highest brightness level of cotton fiber was obtained from laccase treatment as 85.68%. Laccase bio-bleaching showed aromatic functional groups attack which was confirmed by IR spectra. The result indicated that the cotton fibers bio-bleaching using laccase is an environmental friendly process with zero remaining toxicant oxidizing agent which is very promising for medical purposes.

Keywords—Bio-bleaching, cotton fibers, laccase, *Trametes versicolor*

I. INTRODUCTION

Bleaching is an oxidizing process for removing dyes in organic materials [1, 2]. The bleaching process on textile fibers needs to be done to make the fiber color more uniform in the coloring process until the finishing process in the textile industry [1]. The most common natural fiber is cotton, which are consist of oils, fats, waxes, minerals and natural dyes [3]. Natural fibers and fabrics that have undergone a process of scouring still contain natural dyes that cause the color change to yellowish or brownish [4]. A change of yellowish or brownish color on cotton fibers is caused by flavonoids oxidation [5, 6].

Hydrogen peroxide as a conventional bleaching agent has applied under conditions alkaline [7]. The conventional bleaching process using hydrogen peroxide requires large amounts of water to neutralize the reaction [8]. However, conventional bleaching does not eliminate the yellowish color of cotton fibers perfectly [9]. The residual hydrogen peroxide still present in the cotton fibers also causes the process of cotton fibers staining to produce poor quality as a result [10].

Bio-bleaching process is an enzymatic technique to reduce the use of alkaline and more time-saving process [3, 11]. The enzymes that are widely used in textile bio-bleaching processes for instance laccase, manganese oxidase, cellobiose dehydrogenase, and catalase [2]. Laccase (benzendiol: oxygen oxidoreductase, EC 1.10.3.2) is an oxidoreductase-class extracellular enzyme containing copper metal and capable of oxidizing some aromatic (diphenol, monophenol, aromatic amine) and non-aromatic compounds via radical catalyst-reaction mechanisms [12]. Laccase uses molecular oxygen to oxidize aromatic and non-aromatic compounds as a substrate and reduces it into water [13]. The good thing about laccase is that in the bio-bleaching process, it only requires molecular oxygen as acceptor electron meanwhile other oxidoreductases are also required other form acceptor electron, such as manganese and hydrogen peroxide. In the nature laccase can be found in insects, plants and microbes [14]. However, the fungi laccase has higher redox potential than other organisms (up to +800 mV) make this source of laccase is the most promising for industrial applications [15]. The fungi which are able to produce laccase comes from genus *Trametes* [12] with a wide area of applications such as the decolorization and detoxification of dyes and paper bleaching, phenol removal in wine processing, biosensors and synthesis of any organic compound [16].

Cotton fiber is mainly a natural polymer composed of cellulose [17]. The newly harvested cotton fibers have a high impurity content of natural pigments [5]. The removal of natural pigment impurities is usually done by conventional bleaching techniques that could damage cotton fibers molecular structure and requires high operational costs [2]. Therefore, in this study demonstrates the novel strategy of

cotton fiber using laccase produced by *T. versicolor* to remove natural pigment impurities in moderate treatment conditions in order to avoid damage to the main polymer of cotton fiber. Accordingly, as a result the laccase treated cotton fiber could be a promising smart medical based material.

II. MATERIAL AND METHOD

T. versicolor was obtain from the Indonesian Culture Collection (InaCC), Center for Biological Research Institute of Science Indonesia. The strains have been tested for laccase producing ability through a specific fermentation medium. The performance of laccase to cotton fiber phenolic oxidation was analyzed using spectrophotometric analysis, FTIR spectroscopy and cotton brightness test analysis.

A. Fungal maintenance and fermentation medium

Stock cultures of *T. Versicolor* were maintained by storing at 26°C in a cool cabinet and regularly weekly sub culturing on Potato Dextrose Agar (PDA) medium for 5–6 days at room temperature [7, 13]. Prior using for laccase production, the fungi cultures were then inoculated on GDP medium and incubated for 6–7 days at room temperature with shaking at 130 rpm.

B. Laccase production and enzyme preparation

The enzyme production was performed using a fermentation medium formulated by Fukushima and Kirk [18] and Nyanhongo [13]. *T. Versicolor* starter culture was [29] as much as 15 ml to fermentation medium and incubated at room temperature while shaking at 130 rpm. The laccase produced under fermentation medium were harvested based on the growth curve profile of *T. Versicolor* after the fungus enters the start of the stationary phase (day 9) [19]. After incubation, the liquid culture was filtered through Whatman No. 1 and centrifuged at 5,000 rpm for 15 min. The supernatants were carefully collected and used as crude enzyme extract for laccase assay (activity test, protein content and isoelectric point) [14, 20, 21].

C. Protein concentration assay, Laccase activity, and Isoelectric point

The concentration of protein was measured using the Bradford protein assay method [22] at 595 nm and Bovine Serum Albumin (BSA) as a standard.

The laccase activity test was performed using 2,2'-azino-bis (3-ethylbenzothiazoline-6-sulfonate) (ABTS) as the substrate. The laccase activity test was conducted using the modified method [23]. Determination of laccase activity using ABTS were carried out by dissolving 0.5 ml of AB [3] 0.1 mM, 0.01 ml of crude laccase extract, and 2.4 ml of 50 mM citrate buffer at pH 4.5. The reaction mixture was incubated at 35°C for 10 min. The ABTS standard curve was used for enzyme activity determination [20].

The isoelectric point (pI) of the laccase was determined by preparing 6 test tubes containing 1ml of enzyme solution and 1 ml of phosphate buffer with vary level of pH namely 3, 4, 5, 6, 7 and 8. The mixtures were vortex vigorously for 3 seconds. The time required for coagulating were recorded. Precipitation in the respective mixtures were used as isoelectric point value.

D. Cotton fiber bio-bleaching

For bleaching experiment, it was used five treatments, namely control (without treatment), H₂O₂ (30% commercial stock solution), laccase (1% v/v) laccase (2%v/v) and laccase (1%v/v) with mediator (ABTS).

The bleaching with H₂O₂ was carried out using the method reported by Osluoglu, and G. Arabaci [3] and Diller and Yang [24] with slight modification. Briefly, the cotton fibers were placed into the six Erlenmeyer flask containing 5%(v/v) of H₂O₂, 3%(v/v) of NaOH 0.2 M, and 3%(wt/v) of sodium silicate. The ratio between cotton fiber to the solution was set into 1:20. The bleaching process were carried out at 80–95°C for 30 minutes. After bleaching process, the cotton fibers were neutralized with 10% of acetic acid solution and rewashed one time with distilled water.

Bio-bleaching of cotton fibers using laccase was performed with 2 different laccase concentrations of 1%(v/v) and 2%(v/v). Bio-bleaching of cotton fibers was done according to Pereira et al [7] with slight modification. The experiments [4] are initiated by heating the cotton fibers in boiling water for 15 minutes. Furthermore, the cotton fibers were washed with distilled water and let air dry for 7 h. The bio-bleaching reaction was started by putting 1 g [24] cotton fiber in each of two Erlenmeyer flasks containing 10 ml of citrate buffer pH 5.0, 5 ml of laccase 1% and laccase 2%, respectively. Meanwhile, the bio-bleaching of cotton fibers using laccase with ABTS were carried out the same way as bio-bleaching of cotton fibers using laccase above. One gram of cotton fibers was placed in the Erlenmeyer flask containing 5 ml of laccase, 2.5 ml of ABTS 0.1 mM and 10 ml of citrate [23] buffer pH 5.0. Both of the bio-bleaching experiments were incubated at 37°C for 45 min. Finally, in order to terminate the enzyme reaction, the mixtures were added 20 ml of boiling water and let stand for 5 min. The treated cotton fibers then were washed 3 times with distilled water and again let it air dry for 7 h and stored in room temperature until used for the next analysis.

E. Analysis of residual pigments on enzymatic modified cotton fiber, FTIR (Fourier Transform Infrared Spectroscopy) analysis, and Cotton Fiber Brightness analysis

The estimation of cotton fibers pigment residue was measured according to the absorbance maxima at $\lambda=510$ nm using UV-Vis Spectrophotometer [25, 26]. The fibers were soaked in acetone prior analysis.

Cotton fibers before and after the bio-bleaching process were analyzed using FTIR Spectroscopy at wavelengths 400–4000 cm⁻¹ to identify functional groups [27].

Further, the cotton fibers brightness was measured using Lorentzen & Wettre Elrepho Spectrophotometer Testing tool at a local paper print company PT Adiprima Suraprinta, Gresik-Indonesia.

III. RESULT AND DISCUSSION

A. The Production and characterization of *T. versicolor* laccase

Laccase production was carried out by growing the fungi *Trametes versicolor* in medium fermentation formulated by Nyanhongo et al [13] and Fukushima and Kirk [18]. The

fermentation medium was modified by the addition of copper (8) sulfate (CuSO_4) as inducer [14, 28]. CuSO_4 is the constituent of the catalytic center of laccase, hence it is important for the synthesis of the enzyme.

Characterization of *T. versicolor* laccase includes the activity, protein concentration and isoelectric point. Laccase activities (10) achieved 1,400.07 U/ml. using ABTS as a substrate. ABTS is a heterocyclic non-phenolic compound that can be oxidized to a stable cation radical (ABTS^+) and ABTS dication (ABTS^{2+}) [28, 29]. The blue-green color of ABTS^+ cationic radical intensity is correlated to the laccase activity [29]. Protein concentrations in crude laccase were observed using Bradford reagents using BSA as a standard. The protein concentration at the end of laccase fermentation was achieved at 0.172 mg/ml of BSA equivalent.

(22) The surface of a protein has a net charge and would be on zero at a certain pH called an isoelectric point [31]. In general, amino acids in proteins have different isoelectric points. Based on the isoelectric assay, laccase w (19) precipitate at pH 4.0 and the results were in accordance with a study conducted by Bourbonnais et al (5) [31] which stated that laccase produced by *T. versicolor* has an isoelectric point at pH 4.0. At the isoelectric points, there is no electrostatic force, so the enzyme has solubility lowest at the isoelectric point and coagulation occurs [31, 32]. Characterization of laccase demonstrates the use of rice straw and CuSO_4 in the fermentation medium serves as laccase inducer. Rice straw mainly consisted of lignin and *T. versicolor* uses as carbon source. The good thing about Laccase is that it uses only dioxygen to oxidize substrates as electron acceptor while other oxidoreductases (i.e. manganese peroxidases) required additional electron acceptor such as hydrogen peroxide for its oxidation [14].

B. Cotton fiber bio-bleaching

For comparison purposes, bleaching of cotton fibers was conducted using different bleaching agents, namely laccase, laccase with mediators ABTS and H_2O_2 . The result of cotton fibers color after bleaching process by each bleaching treatment shows in Figure 1. The color change in cotton fibers is evident in the treatment of H_2O_2 . The bleaching process occurs when the bond of the chromophore (dyestuff) is disconnected or destroyed. The oxidation process of flavonoid compounds in cotton fibers by H_2O_2 occurs at a pH between 10–12. In alkaline conditions, H_2O_2 dissociates to form ion hydroxyl. The mechanism of free radical formation in aromatic (18) ups (chromophores) is carried out by the ion hydroxyl. Free radicals are formed due to the transfer of electrons to peroxide and then demobilization of electrons in double bonds lead decolorization [33].

As shown in Figure 1, the color of cotton fibers treated using laccase 2% is slightly brighter than the laccase-ABTS and laccase 1% treatments, whereas the color of cotton fibers with laccase-ABTS treatment is slightly brighter than the color in the 1% laccase treatment. This suggests that the laccase produced by *T. versicolor* is capable of oxidizing pigments in cotton fibers and in accordance with the statement of Pereira et al [7]. In (17) her word, laccase produced by *T. versicolor* could oxidize the phenolic compounds of lignin present in the pulp and the decolorization of flavonoids in cotton. The brightness level

analysis of cotton bleaching which are determini (16) by the difference of brightness in each treatment is shown Figure 2.




Figure 1. Bleaching cotton fiber visualization: (A) untreated cotton fiber (negative controls); (B) Cotton fiber bleached with H_2O_2 (positive control) (C); Cotton fiber bleached with Laccase 1% (D); Cotton fiber bleached with Laccase 2%; (E) Cotton fiber bleached with Laccase and ABTS.

Cotton fibers naturally contain color pigments of flavonoids that include phenolic compounds [35]. In (15) bio-bleaching process using laccase, the phenolic are oxidized to phenoxyl free radicals by attack the hydroxyl group of aromatic compounds. The phenoxyl free radicals is unstable and then re-oxidized again by laccase (e.g. forming quinone from phenol) or undergoing non-enzymatic reactions (polymerization) [37].

The use of laccase-ABTS as a bio-bleaching agent can also oxidize phenolic compounds in cotton fibers. The addition of a mediator (ABTS) aims to improve the oxidation process [36]. The addition of ABTS is to have higher oxidation of non-phenolic compounds [16], since laccase is mostly only oxidation of phenolic compounds [38]. The amount of oxidized substrate is increased with the addition of mediator [37]. Oxidation of compounds using laccase-ABTS is called the Laccase Mediator System (LMS) [39]. According to Jong-Rok et al [40], the oxidation capacity of phenol compounds in the bio-bleaching process using LMS is greater than using laccase. However, in this study, the brightness level of cotton fibers resulting from bio-bleaching using laccase-ABTS was lower than that of bio-bleaching using 2% laccase. This most probably because of some ABTS were still stick on the fibers.




Figure 2. Brightness level of cotton fiber in the different treatment of bleaching agent.

The purpose of bleaching is to removed impurities from the fiber through removing the chromophore components that absorb light in the fiber and increase the whiteness level of fabric. The bleaching process can improve the optical properties of the pulp against light absorption, light scattering and reflections expressed in white [34]. Figure 2 shows that the highest brightness level is in the treatment using H₂O₂ (positive control) that is equal to 91.92%. The brightness level on laccase treatment 1%, laccase 2% and laccase-ABTS are 79,33%; 85,68% and 81,57%, respectively. The brightness level of cotton fibers resulting from bio-bleaching is higher when compared with treatment result using buffer (negative control). The use of laccase 2% as bio-bleaching agent yields cotton fibers with higher brightness level than using laccase-ABTS and laccase 1%. Brightness level shows the less-absorbing chromophore component. Based on research conducted by Pereira et al [7] pre-treatment of cotton fiber before bleaching process stated that the brightness level has increased significantly that is equal to 8,5% in cotton fiber with pre-treatment using laccase.

Spectrophotometric analysis was performed to determine the absorbance change of solution used before

and after bleaching process. The results of the spectrophotometric analysis are shown in Figure 3. indicated that there is a change in the absorbance of the solution used in the bleaching process. The absorbance ratio of the solution shows that after the bleaching process there is an increase in absorbance caused by the presence of a compound derived from the cotton fiber oxidation process. The result indicated that the absorbance ratio of bleaching solution does not related with bleaching result. In bleaching process using H₂O₂ which is a positive control, the absorbance ratio value is lower when compared with the treatment using laccase. The low absorbance ratio in H₂O₂ treatment can occur because during the bleaching process the H₂O₂ is dissociates into H⁺ and HO₂⁻ [33].




Figure 3. The ratio of the absorption residual cotton pigment after bleaching comparing to the control ($\lambda = 510 \text{ nm}$)

C. FTIR Analysis of Cotton Fiber Bleaching Results

FTIR spectra of cotton fiber functional groups is seen in Figure 4.




Figure 4. FTIR spectra of cotton fibers resulting in bio-bleaching using different agents: untreated cotton fiber (negative controls); (B) Cotton fiber bleached with H₂O₂ (positive control) (C); Cotton fiber bleached with Laccase 1% (D); Cotton fiber bleached with Laccase 2%; (E) Cotton fiber bleached with Laccase and ABTS.

The range of bleaching cotton fiber absorbance in FTIR spectra is at 400–4,000 cm^{-1} wavelength. The absorbance range at the wavelength of 3,000–3,700 cm^{-1} shows the group vibration (O–H) contained in the cotton fibers. The absorption of clusters (O–H) on the control cotton fibers has an absorption intensity at a wavelength of 3,272.72 cm^{-1} . In cotton fibers treated laccase 1%, laccase 2%, laccase–ABTS and H_2O_2 peak shift with higher absorption intensity than control, at wavelength 3,288.52 cm^{-1} , 3,288.52 cm^{-1} , 3,284.5 cm^{-1} , and 3,280.1 cm^{-1} respectively. According to Nugroho Prasetyo et al [23], the uptake of the (O–H) group becomes higher due to increased purity of the cellulose chain after the bleaching process. The hydrogen bond between the cellulose chains becomes stronger than the hydrogen bonds on the unbleached cotton.

The characteristic band at 700–900 cm^{-1} indicates the missing peak, the vibration of the C–H group of aromatic compounds which constitute the dyestuffs in the cotton fibers, namely on bleaching using H_2O_2 , laccase or laccase–ABTS.

These results implicated that the biotechnological potential of laccase as bio-bleaching agents was assessed, confirming the capacity to decolorize of cotton fiber and to enhance undesirable inherent coloring component. By the enzymatic process, dyes decoloring is environmentally-friendly and minimums remain residues. In addition, cotton fiber produced from modifying by laccase is less residual chemical, therefore its appropriate and suitable for alternatively to medical based materials. Cotton fiber materials resulting bio-bleaching has potential for healthcare and medical product ranges from simple gauze, scaffolds for tissue culturing, bandage materials to a variety of prosthesis for permanent body implants and much more [41, 42].

Medical materials are primarily made of biology inert polymers, commonly used in fabrics and other materials that are in contact with skin. As example, in diapers, specialized polymer materials including cellulose, polypropylene, polyester, and polyethylene arranged in different layers used to provide optimal absorption of urine and feces. As such understanding diaper materials and addressing myths about chlorine, latex, dyes, and other chemical additives in diapers is critical [43].

CONCLUSIONS

The conclusions of this study confirm that laccase produced by *T. versicolor* possessed a strong ability for to oxidize cotton fibers. The parameters including the brightness values were analyzed. Under optimal conditions, 2% laccase treatment has very efficient for bio-bleaching as the highest brightness level as 85.68%. FTIR spectra confirmed that bio-bleaching of cotton fibers using laccase destroys cation fiber functional groups. The effect of enzymes treatment could be explained in terms of selective oxidize of coloring component by the laccase used and better preservation of fibers under these peroxide bleaching conditions. Further studies will be carried out to evaluate the uptake fiber as alternatively to medical based materials.

ACKNOWLEDGMENT

This work was supported by Ministry of Research, Technology and Higher Education (RISTEKDIKT) Indonesia through the Penelitian Unggulan Perguruan

Tinggi (PUPT) Grant (No.603/PKS/ITS/2017) and University of Nahdlatul Ulama Surabaya–Indonesia.

REFERENCES

- [1] T. L. Vigo, "Textile Processing and Properties: Preparation, Dyeing, Finishing and Performance", Amsterdam: Elsevier Science B.V, 1994.
- [2] S.M.F. Kabir, M. I. Iqbal, P.P. Sikdar, M. M. Rahman and S. Akhter, "Optimization of Parameters of Cotton Fabric Whiteness," Eur. Sci. J, 2014, vol. 10, pp. 200–210.
- [3] A. Osluoglu, and G. Arabaci, "Bleaching of Cotton / Polyamide Fabrics with Enzyme and Peracetic Acid", Asia-Pacific J. Chem. Eng, 2014, vol. 9, pp. 364–367.
- [4] S. B. Abdul and G. Narendra, "Accelerated bleaching of cotton material with Hydrogen peroxide", J. Textile. Sci. Eng, 2013, vol 3, 140. doi: 10.4172/2165-8064.1000140.
- [5] A.A. Alghamdi, E.S. Abdel-Halim and Z.A. Al-Othman, "Low temperature bleaching of cotton cellulosa using an ultrasound-assisted tetraacetylenediamine/hydrogen peroxide/triethanolamine system", Bioresources, 2016, vol. 11, pp. 2784–2796.
- [6] J. Tan, M. Wang, L. Tu, Y. Nie, Y. Lin and X. Zhang, "The flavonoid pathway regulates the petal colors of cotton flower", Plos-One, 2013, vol. 8, doi: 10.1371/journal.pone.0072364.
- [7] L. Pereira, C. Bastos, T. Tzanov, A. Cavaco-Paulo and G.M. Guebitz, "Environmentally friendly bleaching of cotton using laccases", Environ. Chem. Lett., 2005, vol. 3, pp. 66–69.
- [8] K. D. Mojsov, "Trends in bio-Processing of textile: A review", Adv. Techno, 2014, vol. 3, pp. 135–138.
- [9] M.N. Miljkovic, M. M. urenovic, M. K. Novakovic dan S.S. Radelovic, "Influence of the fluorescent brightener BA on the degree of whiteness of the knitted cotton fabric". Hemijska Industrija, 2011, vol. 65, pp. 61–66.
- [10] A.M. Amorim, M.D.G. Gasque, J. Andreus and M. Charf, "The Application of Catalase for The Elimination of Hydrogen Peroxide Residues After Bleaching of Cotton Fabrics", An. Acad. Bras. Cienc, 2002, vol. 74, pp. 433–436.
- [11] A. Osluoglu and G. Arabaci, "Bio-bleaching of cotton/plyaide fabric with different enzyme system at low temperature", Multi. Eng. Sci. Tech. J., 2015, vol. 2, pp.3280–3284.
- [12] I. Stoilova, A. Krastanova and V. Stanchev, "Properties of crude laccase from *Trametes versicolor* produces by solid-substrate fermentation", Adv. Biosci. Biotechnol, 2010, vol. 1, pp. 208–215.
- [13] G.S. Nyanhongo, J. Gomes, G. Guebitz, R. Zvauya, J.S. Read and W. Steiner, "Production of laccase by a newly isolated strain of *Trametes modesta*", Bioresouce Technol, 2002, vol 84, pp. 259–263.
- [14] A.B. Vantamuri and B. B. Kaliwal, "Production and optimization of laccase by *Marasmius* sp. BBKAV79 in submerged fermentation", Int. J. Curr. Res, 2015, vol. 7, pp. 18308–18314.
- [15] S. Kaur and V. Nigam, "Production and application of laccase enzyme in pulp and paper industry", IMPACT: Int. J. Res. Appl. Nat. Soc. Sci, 2014, vol. 2, pp. 153–158.
- [16] A. Kunameni, F. J. Plou, Ballesterol A and A. Miguel, "Laccases and their applications: a patent review", Rec. Pat. Biotechnol, 2008, vol 2, pp. 10–24.
- [17] A. Kljun, H.M. El-Dessouky, T.A.S. Benians, F. Goubet, F. Meulewaeter, J.P. Knox and R. S. Blackburn, "Analysis of the physical properties of developing cotton fibres", J. Eur. Pol, 2014, vol. 51, pp. 51–57.
- [18] Y. Fukushima and T. K. Kirk, "Laccase Component of the *Ceriporiopsis subvernisporea* Lignin-Degrading System", Appl. Environ. Microbiol, 1995, vol. 61, pp. 872–876.
- [19] A. Fauzi and E. N. Prasetyo, "Biocycling Limbah Batik sebagai Sumber Karbon dalam Produksi Laccase oleh *Trametes versicolor*", Thesis Report, Institut of Sepuluh Nopember Technology, 2015.
- [20] M. Irshad, M. Asgher, M. A. Sheikh and H. Nawaz, "Purification and Characterization of Laccase Produced by *Schizophyllum commune* IBL-06 in Solid State Culture of Banana Stalks", Bioresources, 2011, vol. 6, pp. 2861–2673.
- [21] J. Cilerdzic, M. Stajic and J. Vukojevic, "Activity of Mn-Oxidizing Peroxidases of *Ganoderma lucidum* Depending on Cultivation Conditions", Bioresources, 2016, vol. 11, pp. 95–104.

- [22] M. M. Bradford, "A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding", *Analy. Biochem.*, 1976, vol. 72, pp. 248-254.
- [23] E. Nugroho Prasetyo, T. Kudanga, L. Ostergaard, J. Rencoret, A.C. Gutierrez, J. del Rio, J. I. Santos, L. Nieto, J. J. Barbero, A. T. Martinez, J. Li, G. Gellerstedt, S. Lepifre, C. Silva, S. Y. Kim, A. C. all author., "Polymerization of Lignosulfonates by the Laccase-HBT (1-Hydroxybenzotriazole) system improves dispersibility. *Biores. Technol.*, 2010, vol. 101, pp. 5054-5062.
- [24] G. B. Diller and X. D. Yang, "Enzymatic bleaching of cotton fabric with glucose oxidase", *J. Textile Res.*, 2001, vol. 71, pp. 388-394.
- [25] H. Feng, Y. Li, S. Wang, L. Zhang, Y. Liu, F. Xue, Y. Sun and J. Sun, "Molecular analysis of proanthocyanidins related to pigmentation in brown cotton fibre (*Gossypium hirsutum* L.)", *J. Exp. Bot.*, 2014, vol. 20, pp. 5759-5769.
- [26] S. H. Qian, L. Hong, M. Xu, Y. P. Cai, Y. Lin and J. S. Gao, "Cellulose synthesis in coloured cotton", *Science Asia*, 2015, vol. 41, pp. 180-186.
- [27] D. Saravanan, "Bleaching of cotton fabrics using hydrogen peroxide produced by glucose oxidase", *Indian J. Fibre Text. Res.*, 2010, vol. 35, pp. 281-283.
- [28] P. J. Collins and A. D. W. Dobson, "Regulation of laccase gene transcription in *Trametes versicolor*", *Appl. Environ. Microbiol.*, 1997, vol. 63, pp. 3444-3450.
- [29] L. Airong, Z. Yue, L. Xu, W. Zhu and X. Tian, "Comparative study on the determination of assay for laccase of *Trametes* sp.", *Afr. J. Biochem. Res.*, 2008, vol. 2, pp. 181-183.
- [30] J. Kirkwood, D. Hargreaves, S. O'Keefe and J. Wilson, "Using isoelectric point to determine the pH for initial protein crystallization trials", *Bioinformatics*, 2015, vol. 31, pp. 1444-1451.
- [31] R. Bourbonnais, M. G. Paice, I. D. Reid, P. Lanthier and M. Yaguchi, "Lignin oxidation by laccase isozymes from *Trametes versicolor* and role of the mediator 2,2'-azinobis(3-ethylbenzthiazoline-6-sulfonate) inkraft lignin depolymerization", *Appl. Environ. Microbiol.*, 1995, vol. 61, pp. 1876-1880.
- [32] N. K. Prasad, "Enzyme Technology : Pacemaker of Biotechnology", New Delhi : PHI Learning Private Limited, 2011.
- [33] S. H. Zeronian and M. K. Inglesby, "Bleaching of cellulose by hydrogen peroxide", *Cellulose*, 1995, vol. 2, pp. 265-272.
- [34] P. Coniwanti, M.N.P. Anka and C. Sanders, "Pengaruh konsentrasi waktu dan temperatur terhadap kandungan lignin pada proses pemutihan bubur kertas bekas", *Jurnal Teknik Kimia*, 2015, vol. Vol. 3.
- [35] S. Kim, D. Moldes and A. C. Paulo, "Laccases for enzymatic colouration of unbleached cotton. *Enzy. Microbial. Technol.*, 2007, vol. 40, pp. 1788-1793.
- [36] B. Viswanath, M.S. Chandra, H. Pallavi and B. Rajasekhar-Reddy, "Screening and assessment of laccase producing fungi isolated from different environmental samples", *Afr. J. Biotechnol.*, 2008, vol. 7, pp. 1129-1133.
- [37] O. V. Morozova, G. P. Shumakovich, S. V. Shleev and Y. A. Yaropolov, "Laccase mediator system and their applications: a review. *Appl. Biochem. Microbiol.*, 2007, vol. 43, pp. 523-535.
- [38] V. K. Gochev and A. I. Krastanov, "Fungal laccases", *Bulgarian J. Agri. Sci.*, 2017, vol. 13, pp. 75-83.
- [39] P. Widsten and A. Kandelcauer, "Laccase application in the forest products industry: a review", *Enzy. Micro. Technol.*, 2007, vol. 42, pp. 293-307.
- [40] J. Jong-Rok, P. Baldrian, K. Murugesan and C. Yoon-Seok, "Laccase-catalysed oxidations of naturally occurring phenols: from in vivo biosynthetic pathway to green synthetic applications. *Microb. Technol.*, 2012, vol. 5, pp. 318-332.
- [41] P. Shende and A. Desai, "Impact and scope of intelligent textiles in health care", *J. Bioequiv Availab*, 2017, vol. 9, pp. Doi: 10.4172/jbb.1000364.
- [42] P. B. Wijesingunawardana and B. G. K. Perera, "Development of a cotton smart textile with medicinal properties using lime oil microcapsules", *Acta. Chim. Slov*, 2018, vol 65, pp. 150-159.
- [43] J. Counts, A. Weisbrod and S. Yin, "Common diaper ingredient question: modern disposable diaper materials are safe and extensively tested", *Clin. Ped.*, 2017, vol 56, pp. 23-27.

Enzymatic modification of cotton fiber for promising smart medical based material

ORIGINALITY REPORT

14%

SIMILARITY INDEX

10%

INTERNET SOURCES

10%

PUBLICATIONS

7%

STUDENT PAPERS

PRIMARY SOURCES

1

journals.sagepub.com

Internet Source

2%

2

Asih Setiarini, Hanum Arrosida, Basuki Winarno. "Wireless Ankle Rehabilitation for Post Stroke Recovery based on Calf Muscle Strength", 2018 1st International Conference on Bioinformatics, Biotechnology, and Biomedical Engineering - Bioinformatics and Biomedical Engineering, 2018

Publication

2%

3

ijpsr.com

Internet Source

1%

4

Muhammad Taufiq Hidayat, Maharani Pertiwi Koentjoro, Isdiantoni, Ida Ekawati, Endry Nugroho Prasetyo. "Immobilization of Bacillus sp. SLII-1 on chitosan-alginate hybrid material for promising feedstock supplement", AIP Publishing, 2020

Publication

1%

5	www.ugsas.gifu-u.ac.jp Internet Source	1 %
6	www.omicsonline.org Internet Source	1 %
7	Muhammad Nasir Iqbal, Hafiz, and Muhammad Asgher. "Decolorization applicability of sol–gel matrix immobilized manganese peroxidase produced from an indigenous white rot fungal strain <i>Ganoderma lucidum</i> ", BMC Biotechnology, 2013. Publication	1 %
8	eprints.covenantuniversity.edu.ng Internet Source	<1 %
9	Cinzia Pezzella, Simona Giacobbe, Valerio Guido Giacobelli, Lucia Guarino et al. "Green routes towards industrial textile dyeing: A laccase based approach", Journal of Molecular Catalysis B: Enzymatic, 2016 Publication	<1 %
10	www.tandfonline.com Internet Source	<1 %
11	Rodríguez-Couto, Susana. "Fungal laccase in the textile industry", Fungal Biomolecules, 2015. Publication	<1 %
12	jurnal.permi.or.id Internet Source	<1 %

13

S. Anuradha Jabasingh, Pallavi Sahu, Abubeker Yimam. "Enviro-friendly biofinishing of cotton fibers using *Aspergillus nidulans* AJSU04 cellulases for enhanced uptake of Myrobalan dye from *Terminalia chebula*", *Dyes and Pigments*, 2016

Publication

<1 %

14

iraj.in

Internet Source

<1 %

15

Jong-Rok Jeon. "Laccase-catalysed oxidations of naturally occurring phenols: from in vivo biosynthetic pathways to green synthetic applications : Laccases and natural phenols for organic synthesis", *Microbial Biotechnology*, 05/2012

Publication

<1 %

16

www.eurorad.org

Internet Source

<1 %

17

S.S. Desai, C. Nityanand. "Microbial Laccases and their Applications: A Review", *Asian Journal of Biotechnology*, 2011

Publication

<1 %

18

Submitted to De Montfort University

Student Paper

<1 %

19

Submitted to Universiti Teknologi MARA

Student Paper

<1 %

20	S. Ben Younes. "Purification and characterization of the laccase secreted by the white rot fungus <i>Perenniporia tephropora</i> and its role in the decolourization of synthetic dyes", <i>Journal of Applied Microbiology</i> , 9/19/2006 Publication	<1 %
21	Honghua Jia, Chao Zhong, Fan Huang, Chunming Wang, Lisha Jia, Hua Zhou, Ping Wei. "The Preparation and Characterization of a Laccase Nanogel and Its Application in Naphthoquinone Synthesis", <i>ChemPlusChem</i> , 2013 Publication	<1 %
22	Submitted to University of Central Florida Student Paper	<1 %
23	aem.asm.org Internet Source	<1 %
24	www.freepatentsonline.com Internet Source	<1 %
25	proteinsoc.thai.net Internet Source	<1 %
26	biopublisher.ca Internet Source	<1 %
27	Kun Rohmatan Nazilah, Maharani Pertiwi Koentjoro, Isdiantoni, Ida Ekawati, Wirdhatul	<1 %

Muslihatin, Endry Nugroho Prasetyo. "Effect of laccase oxidation pre-treatment on coffee (Coffea arabica) bean processing waste for composting substrate", AIP Publishing, 2020

Publication

28

Koschorreck, K.. "Comparative characterization of four laccases from *Trametes versicolor* concerning phenolic C-C coupling and oxidation of PAHs", Archives of Biochemistry and Biophysics, 20080601

Publication

<1 %

29

Submitted to Surabaya University

Student Paper

<1 %

30

Spinelli, Daniele, Enrico Fatarella, Angelo Di Michele, and Rebecca Pogni. "Immobilization of fungal (*Trametes versicolor*) laccase onto Amberlite IR-120 H beads: Optimization and characterization", Process Biochemistry, 2012.

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On