

## **SURAT KETERANGAN**

Nomor: 748/UNUSA/Adm-LPPM/VII/2020

Lembaga Penelitian dan Pengabdian Kepada Masyarakat (LPPM) Universitas Nahdlatul Ulama Surabaya menerangkan telah selesai melakukan pemeriksaan duplikasi dengan membandingkan artikel-artikel lain menggunakan perangkat lunak **Turnitin** pada tanggal 06 Juli 2020.

Judul : *Blood Lead Concentrations and The Neuropsychology Scores of Pregnant Women in Klang Valley, Malaysia*

Penulis : Shamsul Baharai Shamsudin, Jamal Hisham Hashim, Nik Nasri  
Nik Ismail, A.Jamal.A Rahman, Maharani Pertiwi Koentjoro

No. Pemeriksaan : 2020.07.09.338

Dengan Hasil sebagai Berikut:

**Tingkat Kesamaan diseluruh artikel (*Similarity Index*) yaitu 24%**

Demikian surat keterangan ini dibuat untuk digunakan sebagaimana mestinya.

Surabaya, 09 Juli 2020

a.n. Ketua LPPM  
Sekretaris LPPM



UNUSA  
LPPM

Dr. Ubaidillah Zuhdi, S.T., M.Eng., M.S.M.  
NPP: 18101208

**LPPM Universitas Nahdlatul Ulama Surabaya**

Website : lppm.unusa.ac.id

Email : lppm@unusa.ac.id

Hotline : 0838.5706.3867

# Blood Lead Concentrations and The Neuropsychology Scores of Pregnant Women in Klang Valley, Malaysia

*by* Shamsul Shamsudin

---

**Submission date:** 06-Jul-2020 07:39AM (UTC+0700)

**Submission ID:** 1353835346

**File name:** Artikel\_Asli\_Blood\_Lead.pdf (919.9K)

**Word count:** 3731

**Character count:** 18575

## Blood Lead Concentrations and The Neuropsychology Scores of Pregnant Women in Klang Valley, Malaysia

Shamsul Bahari Shamsudin<sup>1</sup>, Jamal Hisham Hashim<sup>2</sup>, Nik Nasri Nik Ismail<sup>3</sup>,  
A. Jamal A. Rahman<sup>3</sup>, Maharani Pertiwi Koentjoro<sup>4</sup>

<sup>1</sup>Department of Community and Family Medicine, Faculty of Medicine & Health Sciences, Universiti Malaysia Sabah, Sabah, Malaysia

<sup>2</sup>United Nations University – International Institute for Global Health (UNU-IIGH) Building, UKM Medical Centre, Kuala Lumpur, Malaysia

<sup>3</sup>Universiti Kebangsaan Malaysia Medical Centre, Kuala Lumpur, Wilayah Persekutuan, Malaysia

<sup>4</sup>Department of Medical Laboratory Technology, Faculty of Health, Universitas Nahdlatul Ulama Surabaya, Surabaya, Indonesia

Correspondence:  
Shamsul Bahari Shamsudin,  
Faculty of Medical and Health Sciences, Universiti Malaysia Sabah, Kota Kinabalu, Sabah, Malaysia  
Zip Code : 88400

Email: shamsul@ums.edu.my

Received: March 13, 2020

Revised: March 16, 2020

Accepted: April 4, 2020



### Abstract

Pregnant women with high blood lead posed high risk to their fetus as placental transfer can occur to the fetus. The objective of this study was to identify the relationship between blood lead and the neuropsychological score of women who were in their 3<sup>rd</sup> trimester of pregnancy. These respondents were undergoing a routine antenatal checkup at a teaching hospital located in Klang Valley areas. Blood lead concentrations were analyzed using graphite furnace Atomic Absorption Spectrophotometer (AAS). The neuropsychological scores were measured with WHO Neurobehavioral Core Test battery (NCTB). The test consists of 7 items, which made up of the Digit Symbol, Trail Making, Digit Span, Benton Visual Retention Test, Pursuit Aiming, Santa Ana Manual Dexterity, Reaction Time and Movement Time tests. The mean blood lead was  $7.78 \pm 4.77 \mu\text{g/dL}$ . The mean score for the total NCTB test was  $50.00 \pm 5.24$ . Statistical analysis showed blood lead concentrations were inversely correlated with the total NCTB score ( $r = -0.462$ ,  $p \leq 0.01$ ). The correlation was about 21.3%. The General Linear Model (GLM) showed that age ( $\beta = -0.15$ ,  $p = 0.017$ ), weight ( $\beta = 2.67$ ,  $p = 0.05$ ) and height ( $\beta = -1.97$ ,  $p = 0.05$ ) also influence the total neuropsychological scores. In conclusion, blood lead reduces the total neuropsychological scores. The scores for each of the 7 items were inversely and significantly correlated with blood lead concentrations except for the Trail Making and Santa Ana Manual Dexterity tests.

### Keywords

Neuropsychological scores, blood lead concentrations, pregnant mothers

## INTRODUCTION

Lead could be produced by human activities such as industrial, mining and agricultural activities that can spread by air, water and soil (1). Lead route of entry to human body through the inhalation, ingestion and skin contact. Lead has no function in human body. Previous study suggested lead has a negative effect on human health even though at very low level. Lead is a neurotoxin and its toxicity affect the function of the central nervous system and peripheral nerve. For adults, the main effects are peripheral neuropathy whereby the impulse conduction in the nerve is slowed down. Peripheral neuropathies influence the motor and sensory nerve. The effects are the wrist and the ankle drop that are the result of the defect in the radial and peroneal nerves (1).

Pregnant woman is at very high risk to the lead toxicity because pregnant mothers need high level of nutrient such as calcium as well as iron to support the growth and the development of the fetus. Lead could replace the function of calcium and iron. Tests on the biochemical and physiological process found that the concentration of calcium and iron in the blood are inversely proportional to the lead concentrations (2).

Study shows blood lead levels (BLLs) in pregnant women at 20.8  $\mu\text{g/dL}$  can cause maternal disorders, when BLLs reaches 2.57  $\mu\text{g/dL}$ , it can cause stress and fatigue levels in pregnant women (4). BLLs  $<5 \mu\text{g/dL}$  may

cause pregnant women to have pre-eclampsia and endanger the mother's kidney, while BLLs  $<10 \mu\text{g/dL}$  can increase blood pressure or hypertension (4). Another research done by Bayat et al, 2016 showed there was an increase in systolic blood pressure (0.014 mmHg) and diastolic (0.013 mmHg) following the increase of 1  $\mu\text{g/dL}$  of BLLs ( $p = 0.04$ ) (5).

This study aims to investigate the association of BLLs with neuropsychology of pregnant women in Klang Valley, Malaysia. Klang Valley area was an ex-mining area in Peninsular Malaysia. Active mining activity was in 1950-1970. The residue of lead and others heavy metal was still high in air, water and soil. The Neurobehavioral Core Test battery (NCTB) can detect early signs of disturbance or failure on the functions of nervous system. WHO was introduced NCTB identifying the nervous dysfunction among workers who are working with the neurotoxin chemicals (3).

## MATERIALS AND METHODS

The study population is women in their 3<sup>rd</sup> trimester of pregnancy who reside in the Klang Valley areas. These pregnant women were attending their routine antenatal checkup at a hospital in Kuala Lumpur. Total 202 respondents were selected through purposive sampling based on specific criteria, that she must be Malaysian citizen, in the 3<sup>rd</sup> trimester pregnancy and have

signed consent to participate. Their pregnancies were categorized as “high-risk” because they were referral cases from private or government clinics within the areas. These women were either in their 1<sup>st</sup> or 5<sup>th</sup> pregnancies, had hypertension, diabetes, anemia or a history of previous delivery complication.

Questionnaire interviews were carried out to collect information on the demographic as well as socioeconomic background and their health status. The questionnaire consists of the questions on their age, ethnicity, number of children, educational level, occupation and areas of residential. While the questions on health status include of number of deliveries, medical history, smoking habit, alcohol intake and medication during pregnancy.

### Blood Sampling and Analysis

Venous blood was sampled during or after the 28 weeks of pregnancy. Five (5) ml of the blood samples were put into a special vacuum container tube, which contained heparin (anticoagulant). All sample of blood lead were analyzed using Graphite Furnace Atomic Absorption Spectrophotometer model HITACHI Z5700 with polarized Zeeman. Blood sample were diluted with matrix modifier solution in a ratio of 1:5. The modified matrix solution also acts as antifreeze agent during storage. The modified matrix solution consists of 10 mL Triton-X, 10% 0.3 g ethylene diamine tetra acetic acid

(EDTA) and 5 g ammonium dehydrogenate phosphate in 1-liter deionizer distilled water. Lypocheck sample were used as a reference sample for quality control procedure during the analysis.

### Neuropsychological Assessment

A neuropsychological test conducted in this study were Neuropsychological Core Test Battery (NCTB) and it is commonly used by WHO (3). It was an instrument used to detect early nervous systems failure because of the exposure of neurotoxin agents. This instrument is very sensitive and can detect early sign of nerve disturbance at a very low level of exposure. It consists of the Benton Visual Retention, Digit Span, Digit Symbol, Pursuit Aiming, Reaction Time, Santa Ana Manual Dexterity and Profile of Mood States (POMS) tests. However, the POMS test was not be conducted in this study, instead it was replaced by two more tests, Movement Time and Trail Making tests. The description and functional domain of the tests are shown in Table 1.

### Standard Score Calculation

The raw score obtained from the neurobehavioral tests, was transformed into standard scores. From these transformed data or standard scores, the mean value is 50 and the standard deviation is 10. The calculation for the standard score is shown:

Standard score = [(raw data – mean) / standard deviation] x 10 + 50.

**Table 1.** Descriptions of Neuropsychological (NCTB) Tests

| Test                       | Functional domain tested            | Description   |
|----------------------------|-------------------------------------|---|
| Reaction/Movement time     | Attention/Response & movement speed | It measures how fast a person reacts/moves. It requires sustained attention by the subject.   |
| Digit span                 | Auditory memory                     | It is a test of immediate (short-term) auditory memory that requires focused attention.   |
| Santa Ana Manual Dexterity | Manual dexterity                    | It requires rapid eye-hand coordination movements.  |
| Digit symbol               | Perceptual-motor speed              | It is a test of perceptual motor speed that also requires learning of associations.   |
| Benton visual retention    | Visual perception/memory            | It is a short-term visual memory that requires focused attention. It also measures the ability to organize geometrical patterns in space and memorize them. |
| Pursuit aiming             | Motor steadiness                    | It measures the ability to make quick and accurate movements with the hand.   |
| Trail making               | Motor tracking                      | It measures visual motor tracking and visual scanning. It requires an attention.  |

Source: WHO. 1983. Operational Guide for The WHO Neurobehavioral Core Test Battery. World Health Organization, Geneva. 15 Februari (14).

### Ethical Clearance


This research was approved by the Research Ethics Committee of Hospital Universiti Kebangsaan Malaysia (UKM/EC711002105465-1). Involvement of respondents is based on a written agreement with filling a consent form. Respondents may withdraw at any time if they do not agree or are not satisfied with any study procedures.

### RESULTS

Table 2 shows the demography and socioeconomic background information of the respondents. The mean age for mothers was 29.6 years old and their mean gestation

period was 31.6 weeks. The ethnic and religious distributions of the respondents are shown in Table 2. The Malays made up 58.4% of the respondents. About 35.6% respondents are full time housewives; while others work as general worker, operate a business, self-employed or part-time wage earner. The average household income is RM 2884.7 (Table 2). The mean blood lead level was 7.78 with standard deviation of 4.77  $\mu\text{g/dL}$ .

Figure 1 shows the data distribution of blood lead for the respondents. Kolmogorov-Smirnov Normality Test shows that the distribution was normal ( $p > 0.05$ ).



**Fig 1.** Distributions of Blood Lead Concentrations ( $\mu\text{g/dL}$ )

**Table 2.** Respondents' Demographic and Socioeconomic Background


| Variable                    | Range       | Mean   | Std. Dev. |
|-----------------------------|-------------|--------|-----------|
| Mother age (Year)           | 19 – 44     | 29.6   | 5.57      |
| Mother Education (Year)     | 6 – 19      | 12.1   | 2.08      |
| Father's education (Year)   | 6 – 22      | 12.8   | 2.64      |
| Total Household Income (RM) | 750 – 8500  | 2884.7 | 1383.98   |
| Mother's height (m)         | 1.43 – 1.72 | 1.57   | 7.12      |
| Mother's weight (kg)        | 50.0 – 92.1 | 68.3   | 7.68      |
| Pregnancy duration (weeks)  | 28 – 36     | 31.6   | 2.13      |
| Number of pregnancies       | 1 – 6       | 1.7    | 1.07      |
| Number of children          | 1 – 6       | 1.6    | 1.04      |

$N = 202$

Table 3 shows the scores for neuropsychological test (NCTB) that was carried out on the respondents. It took about 30–45 minutes to complete all the 7 types of NCTB test. Standard Score Index Values

were calculated from the raw scores and used for all statistical analysis.

Figure 2 shows the standardized NCTB score test distributions, which were normal ( $p > 0.05$ ).



**Fig 2.** Distributions of Neuropsychological Scores

**Table 3.** Neuropsychological Scores (NCTB) of Pregnant Mothers


| Neuropsychological scores (NCTB) | Range       | Mean | Std. Dev. |
|----------------------------------|-------------|------|-----------|
| Digit Symbol                     | 29.0 – 78.3 | 50.0 | 10.00     |
| Pursuit Aiming                   | 24.5 – 80.0 | 50.0 | 10.00     |
| Trail Making                     | 26.4 – 73.1 | 50.0 | 10.00     |
| Digit Span                       | 35.8 – 85.2 | 50.0 | 10.00     |
| Benton Visual Retention Test     | 24.8 – 67.3 | 50.0 | 10.00     |
| Santa Ana Manual Dexterity       | 23.1 – 70.8 | 50.0 | 10.00     |
| Reaction Time                    | 23.5 – 72.0 | 50.0 | 10.00     |
| Total Score Total Score          | 36.3 – 67.6 | 50.0 | 5.54      |

*N* = 202

Table 4 shows the correlation between blood lead and each NCTB test. The total NCTB scores also showed inversely significant correlation with the respondents' blood lead except for the Pursuit Aiming and Santa Ana Manual Dexterity tests. The linear model shows that about 21.3% of the variations in blood lead contributed to the

neuropsychological scores. Table 5 shows that the total NCTB scores were influenced by the respondents' blood lead, age, weight and height after all the confounding factors were adjusted. This model shows that these factors contribute 27% of the variations in the NCTB scores.





**Fig 3.** Correlation Between Blood Lead Concentrations and Neuropsychological Scores

16

**Table 4.** Correlation Blood Lead and Neuropsychological Scores

| Neuropsychological test (NCTB) | Blood lead ( $\mu\text{g}/\text{dL}$ ) |           |
|--------------------------------|--|-----------|
|                                | r                                      | p value   |
| Digit Symbol                   | -0.347                                 | < 0.001** |
| Pursuit Aiming                 | -0.021                                 | 0.762     |
| Trail Making                   | -0.254                                 | < 0.001** |
| Digit Span                     | -0.447                                 | < 0.001** |
| Benton Visual Retention Test   | -0.256                                 | < 0.001** |
| Santa Ana Manual Dexterity     | -0.016                                 | 0.817     |
| Reaction Time                  | -0.450                                 | < 0.001** |
| Total score                    | -0.462                                 | < 0.001** |

$N = 202$

\*\* Significant at  $p < 0.01$

**Table 5.** Correlations of Blood Lead on The Neuropsychological Scores After Adjustment of Confounders

| Dependant variable (Mean Neuropsychological score) | Regression coefficient $\beta$ | Statistic | p value   |
|--|--------------------------------|-----------|-----------|
| (Constant) c                                       | 61.74                          | 7.42      | < 0.001** |
| Blood lead ( $\mu\text{g}/\text{dL}$ )             | -0.47                          | -7.51     | < 0.001** |
| Age (years)  | -0.15                          | -2.40     | 0.017*    |
| Weight (kg)  | 0.12                           | 2.67      | 0.008**   |
| Height (m)   | -0.10                          | -1.61     | 0.110     |
| Educational level (years)                          | 0.08                           | 1.18      | 0.239     |
| Household income (RM)                              | -0.13                          | -1.79     | 0.075     |

$N = 202$

\*\* Significant at  $p \leq 0.01$

\* Significant at  $p \leq 0.05$

F value = 12.787,  $p < 0.001$   
 $r = 0.531$ ,  $R^2 = 0.282$

3

## DISCUSSION

<sup>1</sup> The mean blood lead of the respondents was  $7.78 \mu\text{g/dL}$  (Figure 2), which is more than the blood lead of electronic industries soldering workers ( $6.10 \mu\text{g/L}$ ) (4). Pregnant women need an optimum nutrient such as  $\text{Ca}^{2+}$  and  $\text{Fe}^{2+}$  or  $\text{Fe}^{3+}$ , for the growth of the babies (5,6). The body tends to absorb lead if the calcium and iron intake were insufficient in the diet. Lead in the mothers' blood systems has a potential of being transferred to the fetus through the placenta and this process began 2–3 months at an early stage of pregnancies. Blood lead concentrations for pregnant mothers should not exceed  $10 \mu\text{g/dL}$  (7). However, almost 27% of the respondents have <sup>5</sup> blood lead concentrations of more than  $10 \mu\text{g/dL}$ . Studies (8) in African reported that mean blood lead for pregnant mothers who live in the city and the rural <sup>17</sup> ranged from  $0.83$  to  $99 \mu\text{g/dL}$ . The difference of those data was significantly difference. This proves that environment can influence lead concentrations in the blood. <sup>11</sup> There was no significant relationship between blood lead with age and the gestation period. However, the mean blood lead in previous study was higher than this study.

Another study (9), on pregnant women reported that <sup>1</sup> the mean blood lead concentration was  $8.59 \pm 4.45 \mu\text{g/dL}$  and about 27.8% respondents had blood lead <sup>5</sup> concentrations higher than  $10 \mu\text{g/dL}$ . The study found that housewives had higher

blood lead ( $9.55 \pm 5.5 \mu\text{g/dL}$ ) than those working in offices ( $7.44 \pm 2.77 \mu\text{g/dL}$ ), factories ( $8.61 \pm 3.39 \mu\text{g/dL}$ ) and shops ( $7.01 \pm 3.13 \mu\text{g/dL}$ ). <sup>24</sup> However, the difference was not statistically significant. Study by (10) reported that the mean blood lead for women in the Avon are of the UK was range  $0.41$ – $19.14 \mu\text{g/dL}$ . Studies at Port Pirie, Australia (11) reported that 646 women who are 30 to <sup>36</sup> weeks pregnant had an almost similar <sup>31</sup> mean blood lead level of  $7.2 \mu\text{g/dL}$  and 28% <sup>5</sup> had blood lead of more than  $10 \mu\text{g/dL}$ .

In general, neuropsychological scores were calculated from the 7 tests in the NCTB. Correlation tests indicated that blood lead for these pregnant mothers had an inverse significant correlation with Digit Symbol, Digit Span, Trail Making, Benton Visual Retention Test and Reaction Time (Table 4).

Only Pursuit Aiming and Santa Ana Manual Dexterity scores were not significantly correlated with blood lead (Table 4). The NCTB test result shows that pregnant mother with high blood lead have difficulties in concentrating and have short-term hearing and visual abilities (Digit Span Test, Benton Visual Retention Test). They have slow motor speed through vision (Digit Symbol Test) slow reaction toward visual stimulation (Reaction Time Test) and low attention ability, visual scanning and visual motor trailing (Trail Making Test).

In Figure 3 Shown multiple regression test result demonstrated that mean

neuropsychological score for pregnant mothers were influenced by blood lead ( $\beta = -0.56, p < 0.001$ ), age ( $\beta = 0.12, p < 0.017$ ), weight ( $\beta = 0.12, p = 0.08$ ) and height ( $\beta = -0.01, p = 0.050$ ). The model was significant

$$\text{NCTB mean score} = 65.71 - 0.56 (\text{blood lead}) - 0.15 (\text{age}) - 0.10 (\text{height}) + 0.12 (\text{weight})$$

Previous studies showed that blood lead among female workers had the abilities to lower the NCTB score (12,13). Studies (12) showed that 140 female factory operators had a mean blood lead concentration of 30.77  $\mu\text{g/dL}$  and it was inversely correlated only with the Reaction Time test ( $r = -0.201, p = 0.017$ ). Meanwhile (4) found that mean blood lead for female soldering workers at an electronic factory was 6.1  $\mu\text{g/dL}$ , and the comparative group was 4.6  $\mu\text{g/dL}$ . The respondents' blood lead had significant relationships with Digit Span Test ( $p = 0.003$ ), Santa Ana Manual Dexterity Test ( $p = 0.007$ ) and total NCTB scores ( $p = 0.001$ ). The exposed workers had significantly lower mean score compared to unexposed workers for Digit Symbol Test, Trail Making Test and Pursuit Aiming Test.

Pregnant mothers with high blood lead had problems with the ability to concentrate and have poor short-term auditory and visual memory and have the inability to organize geometry patterns in space and memorizing them. They have poor perceptual motor speed and fail to match between symbol and digits.

( $F = 18.23, p < 0.001$ ) with the  $R^2$  value, which shows that 27% of the variations in the NCTB scores were influenced by the variables shown below:

They also had slow reaction to stimulation due to poor ability to pay attention and low attention ability, which cause poor visual motor tracking as well as poor visual scanning.

Blood lead concentrations, age, weight and height influenced 27% of the variations in neuropsychological score after adjusting for confounders. About 1  $\mu\text{g/dL}$  increment in blood lead will reduce 0.4 of the neuropsychological mean score of these pregnant mothers.

## CONCLUSIONS

In conclusion, even though the blood lead concentration is quite low, it affects the neuropsychological ability of the respondents. Digit Symbol, Digit Span, Trail Making, Benton Visual Retention and Reaction Time test scores had significant inverse correlations with the blood lead.

## ACKNOWLEDGEMENTS

Appreciation goes to all respondents and Management of Environmental Health Laboratory, Hospital Universiti Kebangsaan



Malaysia who gave the commitment that this study can be implemented successfully.

## CONFLICT OF INTEREST

3  
There are no conflicts of interest.

## REFERENCES

1. Rubens O, Logina I, Kravale I, Eglite M, Donaghy M. Peripheral neuropathy in chronic occupational inorganic lead exposure: a clinical and electrophysiological study. *J Neurol Neurosurg Psychiatry*. 2009; 71: 200–204.
2. Mahaffey KR. (1995). Nutrition and lead: Strategies for public health. *Env Health Perspect*. 1995; 103(6):191–196.
3. Anger WK. Reconsideration of the WHO NCTB strategy and test selection. 2015; 0: 224–231.
4. Forsyth JE, Islam MS, Parvez SM, Raqib R, Rahman MS, Muehe EM, Fendorf S, Luby SP. Prevalence of elevated blood lead levels among pregnant women and sources of lead exposure in rural Bangladesh: A case control study. *Env Res*. 2018; 166:1–9.
5. Marangoni F, Cetin I, Verduci E, Canzone G, Giovannini M, Scollo P, Corsello G, Poli A. Maternal diet and nutrient requirements in pregnancy and breastfeeding. An Italian consensus documents. *Nutrients*. 2016; 8(629): doi:10.3390/nu8100629.
6. Saunders AV, Craig WJ, Baines SK, Posen JS. Iron and vegetarian diets. 2012; 2: 11–16.
7. León OL, Pacheco JMS, Martínez SE, Rodríguez EE, Juárez FXC, Carrilo AS, Quiñones, Alanís FV, Vargas GG, Hernández MM, Sustaita JD. The relationship between blood lead levels and occupational exposure in a pregnant population. *BMC Pub Health*. 2016; 16(1231): doi:10.1186/s12889-016-3902-3.
8. Bede-Ojimadu O, Amadi CN, Orisakwe OE. Blood levels in women of child-bearing age in Sub-Saharan Africa: A systematic review. 2018; 6 (367): doi:10.3389/fpubh.2018.00367.
9. Moon CS, Zhang ZW, Watanabe T, Shimbo S, Noor Hassim I, Jamal HH, Ikelda M. Malay women in Kuala Lumpur, to cadmium and lead. *Biomarkers*. 1996; 1:81–85.
10. Taylor CM, Golding J, Hibbeln J, Emond AM. Environmental factors predicting blood lead levels in pregnant women in the UK: The ALSPAC study. *PLoS One*. 2013; 8(9): doi.org/10.1371/journal.pone.0072371.
11. Ridzwan SFM, Annual ZF, Sahani M, Ghazali AR. Neurobehavioral performance of estate residents with privately-treated water supply. *Universiti Putra Malaysia, Serdang. Iran J Public Health*. 2013. 42(12): 1374-1386.
12. Zailina H, Nurunniza ZA, Shamsul BS. Comparison of blood lead and the relationship with neuropsychological scores among battery factory workers and administration workers in Selangor, Malaysia. *Sains Malaysiana*. 2004; 33(1):83–100.
13. Shamsul BS, Jamal HH, Nasri NIN, Rahman AJB. The effects of blood lead on the neuropsychology score of 3<sup>rd</sup> trimester pregnant women in the Klang valley, Malaysia. *J Dis Glo Health*. 2016; 11(3): 134–141.

# Blood Lead Concentrations and The Neuropsychology Scores of Pregnant Women in Klang Valley, Malaysia

## ORIGINALITY REPORT

24%

SIMILARITY INDEX

8%

INTERNET SOURCES

15%

PUBLICATIONS

9%

STUDENT PAPERS

## PRIMARY SOURCES

|   |  |    |
|---|--|----|
| 1 | B S Shamsul, H H. Jamal, N I. Nik Nasri, A J. Abdul Rahman, H Zailina. "THE INFLUENCE OF BLOOD LEAD ON THE NEUROPSYCHOLOGY OF PREGNANT WOMEN IN THE KLANG VALLEY, MALAYSIA", <i>Epidemiology</i> , 2005<br>Publication | 8% |
| 2 | Submitted to Universiti Putra Malaysia<br>Student Paper  | 2% |
| 3 | Submitted to Universitas Nahdlatul Ulama Surabaya<br>Student Paper   | 2% |
| 4 | <a href="http://actascientificamalaysia.com">actascientificamalaysia.com</a><br>Internet Source  | 1% |
| 5 | <a href="http://www.who.int">www.who.int</a><br>Internet Source  | 1% |
| 6 | <a href="http://rbafs.org.br">rbafs.org.br</a><br>Internet Source  | 1% |

|    |   |     |
|----|---|-----|
| 7  | <p>Heinrich. "Occupational Exposure of Pesticides on Activities of <i>Alanine aminotransferase</i> and <math>\gamma</math>-Glutamyltranspeptidase in Blood among Vegetables Farmers", <i>American Medical Journal</i>, 2011</p> <p>Publication</p>                          | 1%  |
| 8  | <p><a href="http://www.pinche.hvdgm.nl">www.pinche.hvdgm.nl</a></p> <p>Internet Source</p>  | 1%  |
| 9  | <p>Douglas M. Templeton, Michael Schwenk, John H. Duffus. "IUPAC Glossary of terms used in neurotoxicology (IUPAC Recommendations 2015)", <i>Pure and Applied Chemistry</i>, 2015</p> <p>Publication</p>  | 1%  |
| 10 | <p><a href="http://spaj.ukm.my">spaj.ukm.my</a></p> <p>Internet Source</p>  | 1%  |
| 11 | <p>Kazue Ishitsuka, Kiwako Yamamoto-Hanada, Limin Yang, Hidetoshi Mezawa et al. "Association between blood lead exposure and mental health in pregnant women: results from The Japan Environment and Children's Study", <i>NeuroToxicology</i>, 2020</p> <p>Publication</p> | <1% |
| 12 | <p>Adalberto Luiz Miranda Filho. "Climate Change and Health: An International Panorama :", <i>Epidemiology</i>, 01/2011</p> <p>Publication</p>  | <1% |

13

[www.ajtmh.org](http://www.ajtmh.org)

Internet Source

<1%

---

14

Submitted to University of Ulster

Student Paper

<1%

---

15

[journal2.unusa.ac.id](http://journal2.unusa.ac.id)

Internet Source

<1%

---

16

Veena Kalra, Sheffali Gulati, K. T. Chitrlekha, Pratibha Pande, S. D. Makhijani, C. S. Sharma. "Plumbism—A mimicker of common childhood symptoms", The Indian Journal of Pediatrics, 2000

Publication

<1%

---

17

[www.frontiersin.org](http://www.frontiersin.org)

Internet Source

<1%

---

18

Jin Yan, Zhenyan Gao, Ju Wang, Wenjuan Ma, Xiaolan Ying, Cancan Zhou, Chonghuai Yan. "Family environmental and dietary implications for low-level prenatal lead exposure in Wujiang City, China", Environmental Science and Pollution Research, 2018

Publication

<1%

---

19

[cgspace.cgiar.org](http://cgspace.cgiar.org)

Internet Source

<1%

---

20

Submitted to University of Auckland

Student Paper

<1%

---

- 21 publisher.uthm.edu.my <1 %  
Internet Source
- 
- 22 Chadi Yazbeck, Olivier Thiebaugeorges, Thierry Moreau, Valérie Goua et al. "Maternal Blood Lead Levels and the Risk of Pregnancy-Induced Hypertension: The EDEN Cohort Study", Environmental Health Perspectives, 2009 <1 %  
Publication
- 
- 23 Submitted to Brunel University <1 %  
Student Paper
- 
- 24 Andrzej Furman, Mehmet Laleli. "Analysis of lead body burden in Turkey", Science of The Total Environment, 1999 <1 %  
Publication
- 
- 25 K Fukumoto, I Karai, Y Nishikawa, S Horiguchi. "Erythrocyte arginase activity as an indicator of lead exposure", Occupational and Environmental Medicine, 1983 <1 %  
Publication
- 
- 26 Rahbar, Mohammad, Maureen Samms-Vaughan, Aisha Dickerson, Manouchehr Hessabi, Jan Bressler, Charlene Desai, Sydonnie Shakespeare-Pellington, Jody-Ann Reece, Renee Morgan, Katherine Loveland, Megan Grove, and Eric Boerwinkle. "Concentration of Lead, Mercury, Cadmium,



Aluminum, Arsenic and Manganese in Umbilical Cord Blood of Jamaican Newborns", International Journal of Environmental Research and Public Health, 2015.

Publication

27

Submitted to Dundalk Institute of Technology

Student Paper

<1%

28

[www.medic.upm.edu.my](http://www.medic.upm.edu.my)

Internet Source

<1%

29

Submitted to Oxford Brookes University

Student Paper

<1%

30

[clinicaltrials.gov](http://clinicaltrials.gov)

Internet Source

<1%

31

Submitted to CSU, San Jose State University

Student Paper

<1%

32

Lavanya Vijayasingham, Uma Jogulu, Pascale Allotey. "Chronic illness and sustainable careers: How individuals with multiple sclerosis negotiate work transitions in a middle-income country", Social Science & Medicine, 2020

Publication

<1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On