











Body Temperature, Glasgow Coma Scale (GCS) and Mortality of Patients with Intracerebral Hemorrhage Stroke

Abdul Muhith¹, Teguh Herlambang², Sri Haryuni³, Kun Ika Nur Rahayu⁴, Endang Mei Yunalia⁵, Catur Asmarani⁶, Eko winarti¬, Anwar Mallongi⁶

^{1,2} Nursing Science Department – University of Nahdlatul Ulama, Surabaya (UNUSA) <u>abdulmuhith@unusa.ac.id</u>

3.4.5.6.7 Nursing and Health Sciences Departement – University of Kadiri, Kediri (UNIK) sri.haryuni@unik-kediri.ac.id

⁸Department of Environmental Health, Faculty of Public Health, Hasanuddin University, Makasar, Indonesia

*Corresponding Author Email: abdulmuhith@unusa.ac.id

Author email: abdulmuhith@unusa.ac.id

ABSTRACT

Stroke is the first rank in Asia and the third rank as a cause of death (mortality) in the world after heart disease and cancer. Intracerebral bleeding stroke occurs in almost 52% of deaths from all stroke incidents, resulting in a broader, more fatal and disabling disorders. This study aimed to find out the correlation of the body temperature and Glasgow Coma Scale (GCS) and the incidence of mortality of the stroke patients with intracerebral hemorrhage. This research was a correlational analytic study with a cross-sectional approach. The sample size was 51 samples, selected by the simple random sampling method. Stroke of intracerebral hemorrhage was determined based on clinical and CT scan examination, and the mortality was measured based on the complete medical record data. The data analysis used the Spearman Rank Correlation Test (Rho) with a significance of α = 0.05. The results of the study showed that 44 respondents (86.3%) had body temperatures in the category of hyperthermia. 44 respondents (86.3%) experienced a poor Glasgow Coma Scale (GCS), and 45 respondents (88.2%) experienced mortality. Statistical test results obtained a p value of 0,000 with α of 0.05. Thus, there is a correlation between the body temperature and Glasgow Coma Scale (GCS) and the incidence of mortality of the stroke patients with intracerebral hemorrhage. The coefficient correlation of the body temperature and the mortality of stroke patients with intracerebral $% \left(1\right) =\left\{ 1\right\} =\left\{ 1\right\}$ hemorrhage was r = .739, and that of the Glasgow Coma Scale (GCS) and that of the mortality was r = .911. The study showed a correlation between the body temperature and Glasgow Coma Scale (GCS) and the incidence of mortality of the stroke patients with intracerebral hemorrhage. This study recommended that the respective health workers appropriately make intervention to the body temperature and recognize the awareness status of the stroke patients to reduce intracerebral hemorrhage so as to reduce the patient mortality.

Keywords: Body temperature, GCS, Mortality, Intraserebral Hemorrhage Stroke.

Correspondence:

Abdul Muhith

Nursing Science Department – University of Nahdlatul Ulama, Surabaya (UNUSA)

*Corresponding Author Email: abdulmuhith@unusa.ac.id Author email: abdulmuhith@unusa.ac.id

INTRODUCTION

Stroke is a disease condition due to a sudden cessation of blood flow to the brain. It occurs due to a blockage or rupture of blood vessels, causing a damage to and even death resulted from brain tissues having no exposure to blood flow, lacking oxygen and nutrients1. According to WHO 2010, every year 15 million people around the world experience strokes. About 5 million suffer permanent paralysis and another 5 million die. In Southeast Asia, there are 4.4 million people sufferring from strokes. In 2020 it is estimated that 7.6 million people will die from such strokes2. Stroke cases of intracerebral hemorrhage occur in almost 52% of deaths from all stroke events, resulting in broader, more fatal, and disabling disorders. Mortality cases occuring within 30 days are 32% -50% of patients³. Based on the previous studies, the number of the stroke patients with intracerebral hemorrhage who died while being hospitalized was 63.9%, that of those with intracerebral hemorrhage who lived while being treated and returned home on doctor's indication was 36.1% according to the study conducted4.

According to Basic Health Research data (RISKESDAS, 2013) the incidence of stroke in Indonesia was 12.1 per 1000 population. This data shows an increase compared to the previous year which was around 8.3. This was supported by the statistics of the Republic of Indonesia Health Research and Development Agency in 2014 stating

that with a percentage of 21.1%, the main cause of death in Indonesia was stroke⁵.

Based on the preliminary survey results in November 2018, the number of mortality of intracerebral hemorrhage stroke (PIS) in 2015 at Mardi Waluyo Blitar Regional Hospital was 49.1% and that in 2016 was 57.5%, or there was an increase in the number of mortality due to the intracerebral hemorrhage. This suggested that the high mortality in intracerebral hemorrhage stroke get more attention. An initial survey of 10 respondents taken showed that the body temperature of intracerebral hemorrhagic stroke patients ranged from 37.8° C to 38.9° C. This showed that the body temperature of the stroke patients with intracerebral hemorrhage increased or had hyperthermia, and for GCS patients with intracerebral hemorrhage stroke reached around <8.

The previous research⁶ explained that an increase in the body temperature of $> 37.5\,^{\circ}$ C affected a dangerous outcome of 63.9%. So, the data showed a predictor of body temperature and GCS affected the mortality incidence of intracerebral hemorrhage stroke. Of the intracerebral hemorrhage stroke population of 10-30%, all the strokes treated in hospital ended in disability and death⁷.

According to the previous studies, it was stated that the initial GCS entry of > 8 had a good outcome prediction, whereas GCS of <8 was a predictive of a bad outcome ie mortality within 2 days of treatment⁸. Hemorrhagic stroke

patients on average experienced a high mortality, loss of consciousness upon arrival at the hospital. Another study also stated that of 1175 patients, 87.6% of those with a GCS score of ≤8 had a mortality while the percentage of the mortality of those with a GCS score of ≥9 was 23.3% . Based on the phenomenon above, a research was needed to study the relationship between the body temperature, the Glasgow Coma scale (GCS) and the incidence of mortality in stroke patients with intracerebral hemorrhage.

MATERIALS AND METHODS

The research design was correlational analytic with cross sectional approach using secondary data sources. The population of the study was all the patients at Mardi Waluyo Blitar Hospital in 2018, diagnosed and turned out to have suffered from a stroke of intracerebral haemorrhage based on history taking, physical examination, and CT Head Scan. The sample used in the study was some of the intracerebral haemorrhagee stroke patients at Mardi Waluyo Blitar Regional Hospital in 2018, diagnosed to have experienced strokes based on neurological examination and supported by brain CT-scan aids and they were not those going home on their own request. The sample size of the study was 51 patients.

The sampling technique used is probability sampling with the Simple Random sampling method. The variables observed or examined in the study were body temperature, Glasgow Coma Scale (GCS) and mortality. The parameters used for body temperature were hypothermia if the body temperature reached <35.5° C, normothermic if body temperature reached 36-37.50C, and hypertension if body temperature reached > 37.5° C, using thermometer and an ordinal scale. The Glasgow Coma Scale (GCS) parameters used are eye opening response (E), verbal response (V), motor response (M). GCS is categorized as bad if the score is <8, moderate: 9-12, good: 13-15 and uses an ordinal scale. For mortality, the parameters used are the cessation of vital functions, by measuring medical records and using nominal scales. The data of the study were obtained from the hospital medic records. The data obtained were then analyzed using the Spearman Rank (Rho) Correlation Test.

Α		
Age		
55 – 64 years	12	23.5%
65 – 74 years	23	45.1%
>75 years	16	31.4%
Job history		
Farmer	16	31.4%
Private company	16	31.4%
PNS (Government	5	9.8%
employees)	14	27.5%
IRT		
Stroke history		
Never	7	13.7%
Ever	44	86.3%
Education		
No formal education	9	17.6%
Elementary School	27	52.9%
High School	10	19.6%
College / University	5	9.8%
Blood tension		
Normotensiveon	1	2%
Hypertensionon	50	98%

Based on the data above, almost half of the respondents aged > 75 years old or under the category of the elderly, totaling 23 respondents (45.1%), a small proportion of the respondents had jobs as farmers and the private sector, each with 16 respondents (31.4%) . Nearly all the respondents had a history of stroke, with a total of 44 respondents (86.3%) and most of the respondents had basic education, with a total of 27 respondents (52.9%). The blood pressure history possessed by most respondents was accompanied with a history of hypertension, and the number was 50 respondents (98%).

Table 2. The correlation of GCS and mortality incidents to the stroke patient with intracerebral bleeding

GCS	Mortality				Total	
	Alive		Dead			
	F	%	F	%	f	%
Good	5	9.8	1	2	6	11.8
Average	1	2	0	0	1	2
Poor	0	0	44	86.3	44	86.3
Total	6	11.8	45	88.2	51	100
Correlation coefficient Sig. (2-taile .000	= .911**	α=0,05				

Sources: Secondary Data of Medical Records, 2018

RESULTS

Table 1. Characteristics of Respondent

Characteristics	Freuency	Percentage

Based on table 2 above, it is obtained that almost all the respondents hyaving a poor GCS experienced a mortality, and the number was 44 respondents (86.3%).

Table 3. Correlation of body temperature and mortality of stroke pation with intracerebral bleeding

Mortality	Total

GCS	Alive		Dead			
	F	%	f	%	f	%
Hypotermy	0	0	0	0	0	0
Normotermy	5	9.8	2	3.9	7	13.7
Hypertermy	1	2	43	84.3	44	86.3
Total	6	11.8	45	88.2	51	100
Correlation coefficient = .739 Sig. (2-tailed) = .		α=0,05				

Sources: Secondary Data of Medical Records, 2018

Based on table 3 above, the obtained data indicated that almost all the respondents having hyperthermic body temperature experienced a mortality with a total of 43 respondents (84.3%)

DISCUSSION

Correlation of the Glasgow Coma Scale (GCS) and the mortality of stroke patients with intracerebral hemorrhage

The results of the data analysis using the Spearman rank test showed that almost all the respondents of the stroke patients with intracerebral hemorrhage had a poor Glasgow Coma Scale (GCS) resulting in mortality of 44 (86.3%) of 51 respondents. The results showed that the value of $\rho=0,000<\alpha=0.05$, with a coefficient value of correlation between the Glasgow Coma Scale (GCS) and the mortality of the stroke patients with intracerebral hemorrhage was r = 0.911, the meaning of the correlation direction was positive, and the value of the correlation strength was very strong. Thus, the worse the Glasgow Coma Scale (GCS) of the patient, the higher the mortality of the stroke patients with intracerebral hemorrhage.

The most important estimates for assessing intracerebral hemorrhage outcome (PIS) are intra-cerebral hemorrhage volume, patient's level of consciousness (using the Glasgow Coma Scale score, body temperature, presence of intraventricular blood, age and infratentorial location) 10 . The intracerebral hemorrhage volume and GCS score could be effectively used to predict mortality rates within 30 days with a sensitivity of 96% and specificity of 98%, intracerebral hemorrhage with a volume of > 60 ml, and GCS score of \leq 8 having a mortality rate of 91% in 30 days, intracerebral hemorrhage with a volume <30 ml and GCS score 9- 12 with a mortality rate of 19% and stroke patients with intracerebral hemorrhage with a GCS score of 13-15 having a mortality rate of $1\%^{11}$.

Correlation between body temperature and incidence of mortality of stroke patients with intracerebral haemorrhage

The results of the data analysis using the Spearman rank test found that almost all the respondents of the stroke patients with intracerebral hemorrhage experienced hyperthermy resulting in mortality totaling 43 respondents (84.3%) of 51 respondents. The results showed the value of ρ = 0,000 < α = 0.05, with the coefficient of the correlation of the body temperature and the mortality of the stroke patients with intracerebral hemorrhage reaching r = 0.739, the meaning of the direction was the positive correlation, and the strength value showed strong correlation. Thus, the higher the body temperature of the patient, the higher the mortality of the stroke patients with intracerebral hemorrhage. The high body temperature after intracerebral hemorrhage was associated with hematoma expansion, cerebral edema, increased intracranial pressure, and initial

neurological damage. The higher body temperature of the patients with intracerebral hemorrhage stroke, the higher moortality, and this was due to hematoma expansion, cerebral edema, increased intracranial pressure, and initial neurological damage, resulting in death 12-20. The results of the study conformed to the above theory which states that the higher the body temperature, the higher the incidence of mortality in stroke patients with intracerebral hemorrhage.

CONCLUSION

Some conclusion was drawn; (1) the body temperature of the stroke patients with intracerebral hemorrhage was almost entirely under category of hyperthermy. (2) Stroke patients with intracerebral hemorrhage almost all had a poor Glasgow Coma Scale (GCS). (3) Almost all the respondents with intracerebral hemorrhage experienced a mortality or death incidents. (4) There was a correlation between the body temperature and the mortality of the stroke patients with intracerebral hemorrhage, with strong and positive correlation. (5) There was a correlation between the Glasgow Coma Scale (GCS) and the incidence of mortality of the stroke patients with intracerebral hemorrhage, with a very strong and positive correlation.

ACKNOWLEDMENT

This research was funding support the publication by the LPPM University of Nahdlatul Ulama Surabaya (UNUSA) Indonesia and LPPM University of Kadiri, (UNIK) Kediri, Indonesia in the year of 2019/2020. And thank very much to Patients intracerebral haemorrhagee stroke at Mardi Waluyo Blitar Regional Hospital.

REFERENCES

- Muhith, A dkk (2019). Spatial Pattern and Social Evironmental Risk Factors of Leprosy Occurrence in Barru, Indonesia. Publiser Journal The Lancet, 10 Pages Posted:10 April 2019. THELANCETID-D-19-00461. url: https://papers.ssrn.com/sol3/papers.cfm?abstract_i d=3367052
- 2. Wittenauer BR, Smith L. (2012). Priority Medicines for Europe and the World " A Public Health Approach to Innovation: Ischaemic and Haemorrhagic Stroke. WHO. (December).
- 3. Parry-Jones, A. R., Abid, K. A., Di Napoli, M., Smith, C. J., Vail, A., Patel, H. C., ... & Tyrrell, P. J. (2013). Accuracy and clinical usefulness of intracerebral hemorrhage grading scores: a direct comparison in a UK population. *Stroke*, 44(7), 1840-1845. https://doi.org/10.1161/STROKEAHA.113.001009
- Muhith, A., (2017) Factors Affecting Self-Efficacy on Tuberculosis Patients, Series:Advances in Health Sciences Research, Proceedings of the Health Science International Conference (HSIC 2017), Doi= https://doi.org/10.2991/hsic-17.2017.53, url: https://www.atlantis-press.com/proceedings/hsic-17/25885869.
- Muhith, A dkk (2019). Structural Equation Modeling: Community Empowerment on Health Status and Pregnant Women Nutritional. Journal of Engineering and Applied Sciences Volume 14 Issue: (7): Page No.2385-2392, (2019). ISSN: 1816-949X. DOI:

- 10.3923/jeasci.2019.2385.2392 url: http://medwelljournals.com/abstract/?doi=jeasci.2 019.2385.2392
- Al Mousawi, Z. N., Al Tameemi, K. M., & Haji, G. F. (2012). Predictors of Outcome for Spontaneous Intracerebral Hemorrhage in Iraqi Stroke Patients. *Iraqi Academic Scientific Journal*, 11(4), 503-509.
- Qureshi, A. I., Mendelow, A. D., & Hanley, D. F. (2009). Intracerebral haemorrhage. *The Lancet, 373*(9675), 1632-1644. http://dx.doi.org/10.1016/S0140-6736(09)60371-8
- 8. Go, G. O., Park, H., Lee, C. H., Hwang, S. H., Han, J. W., & Park, I. S. (2013). The outcomes of spontaneous intracerebral hemorrhage in young adults-a clinical study. *Journal of cerebrovascular and endovascular neurosurgery*, 15(3), 214-220.
- Mittal, M. K., & LacKamp, A. (2016). Intracerebral hemorrhage: perihemorrhagic edema and secondary hematoma expansion: from bench work to ongoing controversies. *Frontiers in neurology*, 7, 210. https://doi.org/10.1016/S0140-6736(09)60371-8
- Senapathi, T. G. A., Wiryana, M., Aribawa, I. G. N. M., & Ryalino, C. (2017). Bispectral index value correlates with Glasgow Coma Scale in traumatic brain injury patients. *Open access emergency medicine: OAEM, 9,* 43. https://dx.doi.org/10.2147%2FOAEM.S130643
- Fallenius, M., Skrifvars, M. B., Reinikainen, M., Bendel, S., & Raj, R. (2017). Common intensive care scoring systems do not outperform age and glasgow coma scale score in predicting mid-term mortality in patients with spontaneous intracerebral hemorrhage treated in the intensive care unit. Scandinavian journal of trauma, resuscitation and emergency medicine, 25(1), 102. https://doi.org/10.1186/s13049-017-0448-z
- Kim, J. Y., & Bae, H. J. (2017). Spontaneous intracerebral hemorrhage: management. *Journal of stroke*, 19(1), 28. https://dx.doi.org/10.5853%2Fjos.2016.01935
- 13. The spatial patten and risk factors of leprosy occurence in barru indonesia. Medicine: Public health, Evironmental and Occupational Health. Indian Journal of Public health Research & Development. https://dx.doi.org/10.5958/09765506.2018,009439
- 14. Muhith, A dkk (2018). Analysis of Educational Factors, Interest and Motivation Towards the Behavior of Early Detection of Cancer Cerviks in Women of Fertile Age in the Upt Health Center Kembangbahu Indian Journal of Public Health Research & Development Volume. 9 Number 12. First page 1371-1375. Published: Dec 2018 Print ISSN: 0976-0245 Online ISSN: 0976-5506
 - https://dx.doi.org/10.5958/09765506.2018.02044. 2
- 15. Muhith, A., dkk (2018). The Effectiveness of Health Education on Improving Santris' Knowledge about the Dangers of Scabies at Al Mubtadi'ien Bahrul Ulum Tambak Beras Jombang. Source: INDIAN JOURNAL OF PUBLIC HEALTH RESEARCH & DEVELOPMENT Volume: 9 Issue: 12 Pages: 1421-1425, Published: Dec 2018 Print ISSN: 0976-0245 Online ISSN: 0976-5506 https://dx.doi.org/10.5958/0976-5506.2018.02053.3
- 16. Muhith, A., fardiansyah A., Saputra, M.H., Nurmiyati (2018). Analisis of causes Impacts of Early Marriage on Madurese Sumenep East Java Indonesia. Medicine: Public health, Evironmental and Occupational Health. Indian Journal of Public health Research &

- Development. Volume. 9 Number 8. First page 412-416. (08 Agustus 2018), ISSN 0978-0245 (print), ISSN 0976-5506.
- https://dx.doi.org/10.5958/09765506.2018,00944.
- Ashari, A. E., Muslimin, I., & Mallongi, A. (2020). The Strategy of Directly Observed Treatment Shortcourse (DOTS) Towards Compliance with Treatment of Tb Lung Patients in Tb Lung Mamuju District Hospital, Indonesia. Systematic Reviews in Pharmacy, 11(6), 1185-1187.
 - http://dx.doi.org/10.31838/srp.2020.6.172
- Handayanya, G. N., Mulatb, T. C., & Ekac, A. M. (2020). Resistance Test of Some Bacterial Synthetic Calcular Cycinaris Invitro Amoxicillin and Ciprofloxacin Antibiotics in TK II Pelamonia Hospital Makassar, Indonesia. Systematic Reviews in Pharmacy, 11(6), 1181-1184.
- 19. Muhith, A., Saputra, M.H., fardiansyah A., (2018). Risk factor of rheumatoid arthritis among Eldely in UPT Panti Werdha Mojopahit Mojokerto distric Indonesia. Medicine: Public health, Evironmental and Occupational Health. Indian Journal of Public health Research & Development.Indian Journal of Public health Research & Development, Volume. 9 Number, 6 June 2018, ISSN 0978-0245 (print), ISSN 0976-5506.
- http://dx.doi.org/10.5958/09765506.2018.00589.2.
 20. Mallongi, A., Natsir, M. F., Astuti, R. D. P., Rauf, A. U., Rachmat, M., & Muhith, A. (2020). Potential ecological risks of mercury contamination along community's area in tonasa cement industry Pangkep, Indonesia.
 - *Enfermería Clínica, 30,* 119-122. https://doi.org/10.1016/j.enfcli.2019.10.054