

#### PAPER • OPEN ACCESS

### Preface

To cite this article: 2020 J. Phys.: Conf. Ser. 1469 011001

View the article online for updates and enhancements.



# IOP ebooks<sup>™</sup>

Bringing together innovative digital publishing with leading authors from the global scientific community.

Start exploring the collection-download the first chapter of every title for free.

IOP Conf. Series: Journal of Physics: Conf. Series 1469 (2020) 011001

Preface

The First International Conference on Innovation in Research (ICIIR) is held on 28-29 August 2018 in Bali, Indonesia. The Conference is organized by Universitas Wijaya Kusuma Surabaya in collaboration with Universitas Pendidikan Indonesia, Universitas PGRI Adi Buana, STIE Malangkuçeçwara, Universitas Andalas, Universitas Hindu Indonesia Bali, Universitas Muhammadiyah Gresik, and Stikom Indonesia Bali.

The main theme of the Conference is "Innovation in Research based on Environmental Insight and Entrepreneurship". The objectives of the conference are: (1) to bring together academicians, scientists, researchers, practitioners, and students in international scientific forums; (2) to share and discuss theoretical as well as practical knowledge of innovation in science. This Conference can be a scientific forum to accommodate fruitful discussion among both local and international researchers in various fields of sciences.

We would like to thank all speakers, invited guests, and also other parties and donors for their continuous support to the success of this event. Hopefully, this event provides many banefits for us all.

The Editors

Evangelos Angelou Afendras Achmadi Susilo Hary Sastrya Wanto Muchlis Miftahul Huda

#### LIST OF COMMITTEE

#### **Chairman:**

Dr. Ir. Hary Sastrya Wanto, MS. - Universitas Wijaya Kusuma Surabaya, Indonesia

#### **Co-Chair:**

Ir. Endang Noerhartati, MP., - Universitas Wijaya Kusuma Surabaya, Indonesia

#### **Advisory Board:**

Prof. Bambang Sumintono, Ph.D. Malaya University, Malaysia Evangelos Angelou Afendras, Ph.D. Indian Institute of Management, Indias Dr. Pranav Kumar, Edu Train International, Bahrain Prof. Dr. Nor Hayati Ahmad - Universiti Utara Malaysia, Malaysia Prof. H. Sri Harmadji, dr. Sp. THT-KL (K) - Universitas Wijaya Kusuma Surabaya, Indonesia Prof. dr. Soedarto, DTM&H., Ph.D. Park. - Universitas Wijaya Kusuma Surabaya, Indonesia Prof. Dr. Rochiman Sasmita, drh., MS., MM. - Universitas Wijaya Kusuma Surabaya, Indonesia Prof. Dr. Ismanto Hadi Santoso, MS. Universitas Wijaya Kusuma Surabaya, Indonesia Dr. Ade Gafar Abdullah, M.Si. - Universitas Pendidikan Indonesia, Indonesia Dr. Eng Asep Bayu DN. - Universitas Pendidikan Indonesia, Indonesia Dr. Isma Widiaty, M.Pd. - Universitas Pendidikan Indonesia, Indonesia Prof. Dr. Ratih Hurriyati, MP. - Universitas Pendidikan Indonesia, Indonesia Prof. Dr. H. Muhammad Ali Ramdhani S.TP., MT. - UIN Sunan Gunung Djati, Indonesia Prof. Dr. Tulus Suryanto, AK, CA, CMA. - UIN Sunan Gunung Djati, Indonesia Dra. Erni Haryanti, MA., Ph.D. - UIN Sunan Gunung Djati, Indonesia Yuni Sari Amalia, Ph.D. - Universitas Airlangga, Indonesia Iman Harymawan, Ph.D. - Universitas Airlangga, Indonesia Dr. H. Jarkawi, M.M.Pd. - Universitas Islam Kalimantan Muhammad Arsyad Al Banjari Banjarmasin, Indonesia

Dr. H. Muhammad Alfani, M.Si. - Universitas Islam Kalimantan Muhammad Arsyad Al Banjari Banjarmasin, Indonesia

Prof. Dr. Ermanto, M.Hum. - Universitas Negeri Padang, Indonesia

Prof. Dr. Syahrul R., M.Pd. - Universitas Negeri Padang, Indonesia

Prof. Dr. I Wayan Suastra, M.Pd. - Universitas Pendidikan Ganesha, Indonesia

Prof. Dr. Yupono Bagyo, MS., MM., CHRP., CPTM. - STIE Malangkucecwara, Indonesia

Akhsanul In`am, Ph.D. - Universitas Muhammadiyah Malang, Indonesia

Erfan Rohadi, ST., M.Eng., Ph.D. - Politeknik Negeri Malang, Indonesia

International Conference on Innovation In Research

IOP Conf. Series: Journal of Physics: Conf. Series 1469 (2020) 011001 doi:10.1088/1742-6596/1469/1/011001

Dr. Dyah Nirmala Arum Janie, M.Si. - Universitas Semarang, Indonesia
Dr. Ir. Rudy Laksmono W., MT. - Universitas Pertahanan, Sentul Bogor, Indonesia
Dr. Irma Setyawati, SE., MM. - Universitas Bhayangkara Jakarta Raya, Indonesia
Dr. Lala Palupi Santyaputri, M.Si. - Universitas Pelita Harapan, Indonesia
Dr. Hadi Kuncoro, M.Farm., Apt. - Universitas Mulawarman, Indonesia
Dr. Azizul Kholis, M.Si. - Universitas Negeri Medan, Indonesia
Dr. Maulana, M.M. - Universitas Tamansiswa Palembang, Indonesia
Cynthia Wuisang, ST., Ph.D. - Universitas Sam Ratulangi, Indonesia
Erna Listyaningsih, SE., M.Si., Ph.D. - Universitas Malahayati, Indonesia
Dr. Ir. Pungky Slamet Wisnu Kusuma, M.Si. - Universitas PGRI Adi Buana, Indonesia
Dr. Dra. Sukarjati, M.Kes,- Universitas PGRI Adi Buana, Indonesia

#### **Organizing Committee**

Prof. Bambang Sumintono, Ph.D. Malaya University, Malaysia. Evangelos Angelou Afendras, Ph.D, Indian Institute of Management, India. Dr. Pranav Kumar, Edu Train International, Bahrain. Prof. Dr. Ruswiati Suryasaputra, MS, Universitas Wijaya Kusuma Surabaya, Indonesia Prof. Dr. Ir. Achmadi Susilo, MS., Universitas Wijaya Kusuma Surabaya, Indonesia Prof. Dr. Ismanto Hadi Santoso, MS., Universitas Wijaya Kusuma Surabaya, Indonesia Sarah Yuliarini, SE., M.Ak., Ph.D., Universitas Wijaya Kusuma Surabaya, Indonesia Dr. Umi Enggarsasi, SH., MH., Universitas Wijaya Kusuma Surabaya, Indonesia Dr. Ir. Miftahul Huda, MM., Universitas Wijaya Kusuma Surabaya, Indonesia Dr. Heni Sukrisno, M.Pd., Universitas Wijaya Kusuma Surabaya, Indonesia Dr. Drs. M. Suud, MA., Universitas Wijaya Kusuma Surabaya, Indonesia Dr. Dorta Simamora, M.Si., Universitas Wijaya Kusuma Surabaya, Indonesia Dr. Siti Djamilah, SE., M.Si., Universitas Wijaya Kusuma Surabaya, Indonesia Eva Wany SE. M. Ak., Universitas Wijaya Kusuma Surabaya, Indonesia Diah Yovita Suryarini, S.Sos., M.Pd Universitas Wijaya Kusuma Surabaya, Indonesia Dr. Ade Gafar Abdullah, M.Si. - Universitas Pendidikan Indonesia, Indonesia Dr. Isma Widiaty, M.Pd. - Universitas Pendidikan Indonesia, Indonesia Dr. I Wayan Budi Utama, M.Si., Universitas Hindu Indonesia, Indonesia. Dr. I Gede Ketut Adiputra, Universitas Hindu Indonesia, Indonesia. Dr. I Kadek Budi Sandika, ST., M.Pd., Sekolah Tinggi Ilmu Komunikasi, Indonesia.



## Peer review statement

All papers published in this volume of *Journal of Physics: Conference Series* have been peer reviewed through processes administered by the proceedings Editors. Reviews were conducted by expert referees to the professional and scientific standards expected of a proceedings journal published by IOP Publishing.

## Levels of Cuprum (Cu) and Plumbum (Pb) in Shrimp in the traditional market and their impacts towards the environmental health

#### A Y P Asih\*

Nahdlatul Ulama University, Surabaya, Indonesia

#### \*akasyekti2009@yahoo.com

AbstractAbsorption of heavy metals by the main shrimp Cuprum (Cu) and Plumbum (Pb) will be accumulated in the body and cannot be biodegradable and accumulated in living tissue. Cuprum and plumbum in small amounts are needed in the body's metabolic process but exposure in excess amounts is toxic, so that at a certain level it can increase the risk of death. Heavy metal pollution often comes from human activities, namely industrial waste disposal, domestic waste or pesticide use in agricultural cultivation, plantations, as well as fisheries businesses, especially shrimp farming. This study aims to identify the magnitude of the average levels of cuprum (Cu) and plumbum (Pb) heavy metals in various types of shrimp available in traditional markets, by comparing the average levels of cuprum (Cu) and plumbum (Pb) of various types of shrimp, observation as well done by comparing the physical size of shrimp based on the type of shrimp. and determination of cuprum (Cu) and plumbum (Pb) levels was carried out by atomic absorption spectrometry. The results of research on shrimp sold in the traditional market of Surabaya, showed that the smaller the physical shrimp, the higher levels of cuprum (Cu) and plumbum (PB) will be higher. The highest Cuprum (Cu) and Plumbum (Pb) levels were found in Vannamei (Litopenaeus vannamei) shrimp, Cuprum (Cu) levels of 8,811 mgCu / kg, and Plumbum (Pb) of 1,990 mgPb / kg, on Windu shrimp (Panaeus monodon) Cuprum (Cu) levels of 7,886 mgCu / kg, and Plumbum (Pb) of 0.735 mgPb / kg, whereas for giant prawns (Macrobrachium rosenbergii de Mann) showed levels of Cuprum (Cu) of 5,814 mgCu / kg and Plumbum (Pb) of 0.728 mgPb / kg; this average level is higher than the WHO standard of 0.691 mgCu / kg for Cu, and 0.715 mgPb / kg for Pb.

#### 1. Introduction

Consumer demand for various types of fish and crustaceans, including various types of shrimp, continues to increase every year due to the increasing welfare of the people in Indonesia, so that the quality of consumption shrimp for both exports and local communities is increasing [1]. The quality of shrimp must meet the food safety agreed upon by the world community, one of which is heavy metal content [2,3].

The issue of increasing research on heavy metals in the environment is largely due to the level of damage and the very high level of poisoning both directly and indirectly towards living creatures and their environment. A number of chemicals are known to be carcinogenic, especially in humans and animals, for example heavy metals lead (Pb) and copper (Cu), are inorganic compounds dissolved solids [4], such as other heavy metals known as metals that are dangerous if their presence is in the human body or other animal's body including the shrimp exceeds the threshold.

Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI. Published under licence by IOP Publishing Ltd 1

International Conference on Innovation In Research IOP Publishing IOP Conf. Series: Journal of Physics: Conf. Series **1469** (2020) 012118 doi:10.1088/1742-6596/1469/1/012118

Basically shrimp culture always requires water in a clean state, but in reality river water and coastal waters are often used directly as a source of water for shrimp culture [5,6], but now many aqir sources which have been contaminated with various types of pollutants, including heavy metals, while the need for shrimp production continues to increase throughout the year, so special treatment is needed on the culture, including the use of chemicals in the form of fertilizers and pesticides given that shrimp are very susceptible to pests and diseases [6]. Now many water sources have been contaminated with various types of pollutants including heavy metals that come from various human activities and come from nature [7]. Cuprum (Cu) and plumbum (Pb) are very toxic heavy metals and many are always present in water at various levels of pollution in the environment [8].

The type of shrimp that is widely found in traditional markets is the Windu shrimp (Panaeus monodon); vannamei / white shrimps (Litopenaeus vannamei), freshwater shrimp / river shrimp (Palaemon Sp); giant prawns (Macrobrachium rosenbergii de Mann), grago shrimp or rice shrimp (Caridina Sp), rebon shrimp (Mysis Sp), api2 shrimp (Metapeneoues monoceros). Based on the description above, a research is needed on the content of heavy metals in shrimp that are sold as consumption ingredients in traditional markets considering the effects of heavy metals are very risky to public health. This study aims to compare the average levels of cuprum (Cu) and plumbum (Pb) of various types of shrimp sold in the Surabaya traditional market.

#### 2. Literature review

Heavy metals with a certain amount are needed by living creatures as important micronutrient materials. Some marine biota including shrimp are biota that live in the bottom of the sea and have the ability to accumulate heavy metals in their bodies far beyond the concentration in the surrounding waters and are widely used as bio indicators of the presence of heavy metals in the waters [9].

Some heavy metals are important elements because they are needed as a physiological function, but if they are found in food, water or air, they will have a negative impact on health, including being able to adversely affect human health as the end chain of the food chain in the life process. However, some heavy metals are essential micronutrients that are needed for the metabolism of the shrimp body, while some other types do not have a biological (non-essential) role even at very high levels are very deadly [6,10,11]. Crustaceans include crabs, squid and various types of shrimp sold as supplementary ingredients (side dishes, or even often used in certain cultural processes) or just as flavoring ingredients or ingredients that are consumed. it means that shrimp can be consumed directly as a side dish, a mixture of crackers, crackers, or other preserved ingredients.

Because the habitat in the bottom of the water which is a place of sediment from various types of waste, then this type of crustacean is a good bio indicator to determine the occurrence of environmental pollution, this is because many types of crustaceans eat zooplankton and phytoplankton which are widely found at the bottom of the water while accumulating heavy metals originating from aquatic waste [6]. Shrimp is classified as a booster indicator even though every animal in the water always has the ability to do osmoregulation (as an effort to neutralize toxins around the environment), so the ability of animals that live in the waters of one of the shrimp is very dependent on their physical ability to avoid toxins in the environment (waters) or it can be said that the greater the physical shape of the shrimp, the higher the osmoregulation ability [4]. Shrimp (Crustacean) can live in various types of deep sea waters, shallow sea waters (beaches), brackish water (ponds) or even in certain types can live in fresh water (rivers), for example in types of giant prawns (Macrobrachium rosenbergii de Mann) during its life cycle requires 2 types of water, namely fresh water for the larval period and brackish water for adults.

Based on the amount of shrimp physics, the biggest successive are giant / conggah shrimp (Macrobrachium rosenbergii de Mann), Windu shrimp (Panaeus monodon); vannamei shrimp (Litopenaeus vannamei), freshwater shrimp / river shrimp (Palaemon Sp); grago shrimp or rice shrimp (Caridina Sp), rebon shrimp (Mysis Sp), api2 shrimp (Metapeneoues monoceros).

• The smaller the physical shape of the shrimp, the higher the levels of cuprum (Cu) and plumbum (Pb)

• Vannamei / white shrimps (Litopenaeus vannamei), has the highest levels of Cuprum (Cu) and Plumbum (Pb) when compared

#### 3. Data and methodology

Based on direct observation, the most consumed shrimp in Surabaya in sequence are as follows:

- Vannamei shrimp (Litopenaeus vannamei).
- Tiger Prawn (Panaeus monodon)
- Prawns (Macrobrachium rosenbergii de Mann)

Determination of cuprum (Cu) and plumbum (Pb) heavy metal content in shrimp was carried out using Spectrometry (AAS), with some beker 50 ml glassware, 10 ml volumetric flask, 5 ml polyethylene vial, 10-100  $\mu$ l effendorf micropipette and analytical balance.

The study was conducted in May 2018 in 5 Surabaya traditional markets, the traditional market in question is a market managed by Surabaya regional. Shrimp samples were deliberately chosen by 3 types based on the number of shrimp species available in the market, then the shrimp species were chosen randomly with 150 samples (each market took 10 tails), Sampling is done randomly and intentionally (without distinguishing the condition of the shrimp alive or dead), but it is still chosen that the shrimp can be used as a sample (the shrimp die but not in a rotten state), the sample is taken based on the market area (Surabaya market west, east, south, north and central Surabaya market).

Determination of cuprum (Cu) and plumbum (Pb) heavy metal content in shrimp in Airlangga University Surabaya Biology laboratory. Data calculation is done directly by comparing the average results of cuprum (Cu) and plumbum (Pb) heavy metal levels on various types of shrimp by looking at the small physical body size of each type of shrimp.

#### 4. Results and discussion

Based on direct observation in the field, the shrimp available in the traditional market are often available in a dead state (without knowing the cause). This is likely due to environmental pollution, the use of chemicals during cultivation or mismanagement of harvest or even during the sale process (addition of formalin or ingredients others as preservatives. Shrimp is a food commodity very quickly damaged / rotten in just a matter of hours it is also often based on the physical size of the shrimp, so often traders add preservatives (with the amount of uncertain ingredients) in the merchandise without the buyer's knowledge such as the addition of formaldehyde and certain types widely sold in the traditional market / krempyeng market (in Java).

Osmoregulation (shrimp adaptability to the surrounding environment) in the shrimp occurs at any time if the surrounding waters are contaminated by toxins or heavy metals, the greater the physical shape of the shrimp, the greater the osmoregulation power given the shrimp's ability to avoid its environment will greatly depend on shrimp mobility (the greater the physical shape of the shrimp, the greater the ability to avoid).

Based on the type of shrimp by looking at their physical size, the selling price of shrimp in the market is also very different, the bigger the physical shape of the shrimp, the higher the selling price of shrimp so that it will not be bought by the weak economy. based on the type of shrimp sequentially are:

- Vannamei shrimp (Litopenaeus vannamei)
- Tiger Prawn (Panaeus monodon)
- Prawns (Macrobrachium rosenbergii de Mann)

Considering that the need for vannamei shrimp (Litopenaeus vannamei) in the community is very high, the fish farmers often make efforts to increase shrimp production in various ways to avoid losses due to shrimp death (efforts to avoid crop failure), for example using chemicals (pesticides and chemical fertilizers). in quantity and relatively high / high intensity), although vannamei shrimp can be directly obtained in shallow sea water, this type of shrimp can also be cultivated in brackish water (pond).

Shrimp Type		Average					
	1	2	3	4	5		
Vaname Shrimp	8.377	8.803	8.754	9.053	9.068	8.811	
Windu Shrimp	8.390	8.124	8.512	7.106	7.300	7.886	
Prawn Shrimp	6.423	5.214	5.832	5.737	5.864	5.814	
	Source: Results of laboratory analysis						

Table 1. The average Cu (mg/Kg) based on the type of Shrimp

Based on the results of the study it turned out that the average Cuprum (Cu) level was higher than the Plumbum (Pb) levels in each type of shrimp. Likewise, almost every traditional market in Surabaya, it is known that Cu levels are higher than Pb levels in each type of shrimp based on the physical size of shrimp (table, 1). Cuprum (Cu) is a heavy metal that is always present in the waters. Concentration of Cu in the water is now very high [10], in waters the Cu content should not exceed 0.05 mg / L and natural source waters should not exceed 0.1 mg / L [3]. if the Cu content in the waters is very high, the Cu content in the shrimp body is also high.

Whereas heavy metal Plumbum comes from the air as well as various antropogenik and erosion activities. Cu and Pb at high concentrations are xenobiotic (very dangerous for living creatures) and in the body can cause mutagenic and carcinogenic events [6]. In table 2 it can be seen that the average Pb level is lower when compared to Cu levels in all samples taken from traditional markets in Surabaya, even though the high low levels of heavy metals depend on the high and low pollution in the waters.

Shrimp Type		Average					
	1	2	3	4	5	_ 0	
Vaname Shrimp	0.160	0.271	0.376	0.075	9.068	1.990	
Windu Shrimp	0.542	0.700	0.814	0.519	0.101	0.735	
Prawn Shrimp	0.954	0.660	0.804	0.953	1.227	0.728	
	Source: Results of laboratory analysis						

Table 2. The Average Pb (mg/Kg) based on the Type of Shrimp

The results showed that the levels of cuprum (Cu) and plumbum (Pb) heavy metals in shrimp sold in the Surabaya traditional market were high on average. in giant prawns (Macrobrachium rosenbergii de Mann) showed levels of Cuprum (Cu) of 5,814 mgCu / kg and Plumbum (Pb) of 0.728 mgPb / kg; Windu shrimp (Panaeus monodon) levels of Cuprum (Cu) of 7,886 mgCu / kg, and Plumbum (Pb) of 0.735 mgPb / kg, while for Vannamei (Litopenaeus vannamei) shrimp, Cuprum (Cu) levels were 8,811 mgCu / kg, and Plumbum (Pb) of 1.990 mgPb / kg. The greater the physical shape of the shrimp, the lower the levels of Cuprum (Cu) and Plumbum (Pb).

#### 5. Conclusion

From the results of the study it can be concluded as follows:

- The smaller the physical size of the shrimp, the higher the levels of cuprum (Cu) and plumbum (Pb) metals.
- In the type of shrimp Vannamei (Litopenaeus vannamei), the highest levels of cuprum (Cu) and plumbum (Pb), namely the levels of Cuprum (Cu) of 8.881 mgCu / kg, and Plumbum (Pb) of 1.990 mgPb / kg, when compared with tiger shrimp (Panaeus monodon) and giant prawns (Macrobrachium rosenbergii de Mann) throughout traditional markets in Surabaya.

#### References

[1] Reddy R R, Bindu P and Subendu A 2006 Bioacumulation of Copper in Post Larvae and freshwater Prawn Juvenille (Macrobrachium rosenbergii de mann), *Exposed to Sub-Lethals of Copper Sulfate, Central Institute of Freshwater Aquaqulture, Kausalyaganga, Buhaneswar* 

751002, Orrisa India. Sciendirect. Elsevier Aquaqulture

[2] Palar 2004 Heavy metal pollution and toxicology Jakarta: Rineka Cipta

- [3] 13 World Health Organization (WHO) 1992 Cadmium, Environmental Aspect Environmental Health Criteria
- [4] Soegianto A, Akas Y, Bambang I 2016 Lead Toxicity at Defferent Stage of the Giant Prawn live (Macrobrachium rosenbergii de Mann) considerations of Osmoregulatory capacity and Histological Changes in an Adult Gills Marine and Freshwater Behavior and Physiology. Taylor & Francis
- [5] Soegianto A, Daurews M C, Trilles J P and Charmantier G 1999 Impact of Copper on the Structure of Gills and Epipodites of the Shriems Peneous japonicus (Decapoda) *Journal of Crustacean Biology* 19(2) pp 2009-223
- [6] Akas Y, Bambang I and Agus S 2013 Effect of Copper on Survival, Osmoregulation, and Gill Structure of Freshwater Prawn (Macrobrachium Rosenbergii de Mann) at Defferent Development Stages, Marine and Freshwater Behavior and Physiology *Taylor & Francis*
- [7] Amando M E, Freire C A and Sausa M M 2006 Osmoregulation and Tissue Water Regulation in the Freshwater Red Crab Dilocarcinus pagai (Crustacea, Decapoda) and The Effect of Waterborne Inorganic Lead. Aquatic Toxicology
- [8] Darmono, 2001 Environment and Pollution Relationship with the Toxicology of Metal Compounds Jakarta: University of Indonesia Press
- [9] Saager P M, De Baar H J, de Jong J T, Nolting R F and Schijf J 1997 Hydrography and local sources of dissolved trace metals Mn, Ni, Cu, and Cd in the northeast Atlantic Ocean *Marine Chemistry* 57(3-4) pp 195-216
- [10] Dan'azumi S and Bichi M H 2010 Industrial pollution and implication on source of water supply in Kano, Nigeria *International Journal of Engineering & Technology* **10**(1) pp 101-109
- [11] Martins C D M G, Barcarolli I F, de Menezes E J, Giacomin M M, Wood C M and Bianchini A 2011 Acute toxicity, accumulation and tissue distribution of copper in the blue crab Callinectes sapidus acclimated to different salinities: in vivo and in vitro studies *Aquatic toxicology* 101(1) pp 88-99



# CERTIFICATE



No.: 100/CT/ICIIR/UWKS/VIII/2018 is awarded to

## **Akas Yekti Pulih Asih**

as

## PRESENTER

of a paper entitled

"Concentration of Curcum (Cu) and Plumbum (Pb) Shrimph in Surabaya traditional Markets.

In the International Conference on Innovation in Research Bali, Indonesia, 28<sup>th</sup> – 29<sup>th</sup> August 2018



Prof. H. Sri Harmadji., dr., Sp.THT-KL (K) **Rector of Universitas Wijaya Kusuma Surabaya**  Ir. Endang Noerhartati, MP. **Conference Chair** 

organized by:





