

Self-Management Education for the Quality of Life of Patients with Pulmonary Tuberculosis

Eppy Setiyowati¹, Umi Hanik¹, Erika Martining Wardani¹, Mujad Didien Afandi², Ni Njoman Juliasih³

Abstract---Treatment for pulmonary tuberculosis (TB) may become a problem when the patients show no adherence to the treatment and feel bored with its long duration. This condition may lead to drop out in the treatment. Therefore, this study aimed to analyze the effects of self-management education on the quality of life of patients with pulmonary TB. The design of this study was quasi-experimental, conducted in 3 Puskesmas (Perak Timur, Sawahan, Kebonsari), in which 150 respondents were divided into 2 groups: 75 respondents in the control group and 75 respondents in the treatment group. They were chosen as the sample by using a simple random sampling technique. The data were collected by conducting a pretest to identify the quality of life, interventions through self-management education, and finally providing a posttest. Furthermore, the data were analyzed using the Mann Whitney U test and Wilcoxon signed rank test. Change in the physical health domain, psychological domain, social domain and an environment domain that was effect self-management education the right place for the treatment is puskesmas, and a high hope for a cure needs supports from the family and medical workers. Self-management education increases the quality of life of patients with pulmonary TB. Future research is expected to analyze their life expectancy more deeply.

Keywords--- Quality of Life; Pulmonary TB; Life Expectancy

I. INTRODUCTION

Tuberculosis (TB) is an infectious disease which is a global problem, mostly found in developing countries [1],[2],[3],[4]. Problems related to TB are exacerbated by the presence of comorbidities, which cause inhibiting factors in the treatment in the treatment period [5] [4] [6].

TB control in Indonesia through the national TB program has long been carried out in line with the implementation of short-term treatment strategies and direct monitoring (Directly Observed Treatment Short-course, DOT) carried out in puskesmas (health primary) stage [7] [8] [9].

The elimination tuberculosis is also one of the 3 main focuses of the government in addition, in the health sector, to reducing stunting and increasing the coverage and quality of immunizations [10]; [8]; [5]. The vision relation build to this disease world is free tuberculosis, zero deaths, illnesses, and suffering caused by tuberculosis [11]; [7]; [12].

¹ Faculty of Nursing and Midwifery, Nahdlatul Ulama University of Surabaya, Surabaya, Indonesia

² Faculty of Teacher Training and Education, Nahdlatul Ulama University of Surabaya, Surabaya, Indonesia

³ Institute of Tropical Diseases, Airlangga University, Surabaya, Indonesia

Corresponding Author:

Eppy Setiyowati

E-mail: eppy@umusa.ac.id

The world hopes to be free of tuberculosis, so the quality of life in TB sufferers needs to be investigated so that the treatment process can be run optimally and continuously [13]; [14]; [1]; [15]. Quality of life as an individual's perception is related to their goals, hopes, standards and concerns [14]; [5]; [2]; [16]; [17].

Factors affecting quality of life are physical health, psychological conditions, personal beliefs, social relationships and their relationship with important factors in their environment [18]; [14]; [1]; [17]. Patients with drug resistant TB have different quality of life compared to non-drug resistant TB patients; non-drug resistant TB patients have a better quality of life than drug resistant TB patients [6]; [3]; [19]. The difference in quality of life is the general quality of life (global quality of life), global health, psychological health, and environmental quality of life [3]; [7]; [20]. Aspects of physical health and social relationships do not show any difference between drug resistant TB patients and patients who are non multiple drug resistant [21]; [22]; [7]; [23].

Self management education in pulmonary TB patients involves 3 steps: process creation, peer group discussion, transfer and sharing process [15]; [17]; [19];[3] The purpose of this study was to investigate the relationship of self-management education to quality of life in patients with pulmonary TB.

II METHODS

A quasi-experimental research design was used with one group intervention approach and one control group . The sample came from 3 puskesmas in Surabaya City, and included 150 respondents divided into 2 groups: 75 respondents in the control group and 75 respondents in the treatment group.

Treatment given by researchers involved family empowerment including counseling, guidance / counseling and demonstrations related to TB, how to care, how to prevent transmission and treatment of pulmonary TB involving patients and families for approximately 1 hour for each meeting every week for 6 weeks, in agreement with the respondents. Supporting instruments in the research were booklets, flip sheets and demonstration materials for pulmonary TB independent care [24]; [25]; [26].

The *World Health Organization* (WHO) has developed instruments measuring quality of life consisting of four domains, namely: physical, psychological, social and environmental. The questionnaire was given during the pre-test and post-test. The research process was carried out for 1 month, divided into 2 weeks for the control group and 2 weeks for the treatment group. Analysis was undertaken using the statistical Wilcoxon signed rank test, level of significance $\alpha = 0.05$. That means, if $(p) = 0.05$, then H_1 is accepted; this would means that there is an influence of self-management education on Quality of Life (QoL) among pulmonary TB sufferers. Different test control group and treatment were the Mann Whitney U, test level $\alpha = 0.05$.

III. RESULTS

Respondents' demographic data are the characteristics of respondents which included: 1) gender, 2) age, 3) occupation, 4) education, 5) duration of treatment for pulmonary tuberculosis and the results obtained are shown in Table 1. Quantitative research results (with the Wilcoxon signed rank test and the Mann Whitney U test) obtained the results in Table 2. Statistical test results used the Wilcoxon signed rank test, treatment group (groups that received self-management education treatment) were $p = 0.003$ on the physical health domain means $p = 0.05$, then the hypothesis is accepted meaning there is a significant effect of self-management education on improving quality of life (physical health domain) in patients with pulmonary TB. The statistical test result using the Wilcoxon Signed Rank test, in the control group (groups that didn't get self-management education) was $p = 0.317$ in the physical health domain which means there was no difference in quality of life (physical health domain) in patients with pulmonary TB between *pretest* and *posttest*. Difference in average quality of life (physical health domain) at pretest in the

treatment group and the control group using the *Mann Whitney U Test* was $p = 0,167$; this shows that there was no average difference between the control group and the treatment group. Difference in mean quality of life (physical health domain) at posttest in the treatment group and the control group using the *Mann Whitney U Test* was $p = 0.000$; this shows that there were differences in the mean improvement in quality of life (in the physical health domain) between the treatment group (the group that got SME) and the control group (the group that did not get SME).

For the psychological domain, the statistical test results using *Wilcoxon Signed Rank test*, in the treatment group was $p = 0.003$ means it $p = 0,05$ then the hypothesis is accepted meaning there was a significant effect of self-management education on improving the quality of life (psychological domain) in patients with pulmonary TB. The value control group $p = 0.317$ means it $p = 0.05$ then the hypothesis is rejected meaning that there is no improvement in the quality of life (psychological domain) in patients with pulmonary TB. Difference in average quality of life (psychological domain) at pretest in the treatment group and the control group using *Mann Whitney U test* gave $p = 0.167$; this shows that there was no average difference between the control group and the treatment group. Difference in average quality of life (psychological domain) at posttest in the treatment group and the control group using the *Mann Whitney U Test* gave $p = 0.000$; this shows that there were differences in the mean improvement in quality of life (psychological domain) between the treatment group (the group that received SME) and the control group (the group that did not receive SME).

The social domain, statistical test results used the *Wilcoxon signed rank test*, the treatment group $p = 0.046$ means it $p = 0.05$ then the hypothesis is accepted meaning that there was a significant effect of self-management education on improving the quality of life (social domain) in patients with pulmonary TB. The social domain using *Wilcoxon signed rank test*, in the control group gave $p = 0.157$ meaning there was no difference in quality of life (social domain) in patients with pulmonary tuberculosis. Difference in average quality of life (social domain) at the pretest in the treatment group and the control group using the *Mann Whitney U test* gave $p = 0.061$; this shows that there was no average difference between the control group and the treatment group. Difference in average quality of life (social domain) at posttest in the treatment group and the control group using the *Mann Whitney U test* gave $p = 0.015$; this shows that there were differences in the mean improvement in quality of life (social domain) between the treatment group (the group that received SME) and the group control (groups that did not receive SME).

The environmental domain, statistical test results use *Wilcoxon signed rank test*, in the treatment group is $p = 1.000$ means it $p = 0.05$, the hypothesis is rejected meaning that there was no significant effect of self-management education (SME) on improving quality of life (environmental domain) in patients with pulmonary tuberculosis. The statistical test results using the *Wilcoxon signed rank test*, in the control group was $p = 0.564$ meaning that there was no significant difference in quality of life (environmental domain) between pretest and posttest in patients with pulmonary tuberculosis. Differences in the average quality of life (environmental domain) during the pre-test in the treatment group and the control group using the test the *Mann Whitney U test* didapatkan $p = 0.101$; this shows that there was no average difference between the control group and the treatment group. The average difference in quality of life (physical health domain) at the posttest in the treatment group and the control group using *Mann Whitney U Test* gave $p = 0.167$; this shows that there was no difference in the average improvement in quality of life (environmental domain) between the treatment group (the group that received SME) and the control group (the group that did not receive SME).

Table 1 Table Distribution Responden on Characteristic Demografic

| Respondent Demographics | F | % | F | % |
|---------------------------|-----------------|-----------------|-------------------|-------------------|
| | (Control group) | (Control group) | (Treatment group) | (Treatment group) |
| Gender: | | | | |
| Male | 22 | 29.3 % | 22 | 29.3 % |
| Female | 53 | 70.7 % | 53 | 70.7 % |
| Total Respondent | 75 | 100% | 75 | 100% |
| Age: | | | | |
| >25 - ≤ 35 | 15 | 20 % | 17 | 22.6 % |
| >35 - ≤ 45 | 24 | 32 % | 32 | 42.7 % |
| >45 - ≤ 60 | 36 | 48 % | 26 | 34.7 % |
| Total Respondent | 75 | 100% | 75 | 100 % |
| Profession: | | | | |
| Housewife | 11 | 14.7 % | 15 | 20 % |
| Private sector / Employee | 28 | 73.3 % | 23 | 30.7 % |
| Entrepreneur | 12 | 16 % | 17 | 22.7 % |
| Driver | 27 | 36 % | 10 | 13.3 % |
| Does not work | 3 | 5 % | 10 | 13.3 % |
| Total Respondent | 75 | 100 % | 75 | 100 % |
| Education level: | | | | |
| No school | 11 | 14.7 % | 12 | 16 % |
| Elementary school | 15 | 20 % | 17 | 22.7 % |
| Junior high school | 23 | 30.7 % | 27 | 36 % |
| Senior high School | 19 | 25.3 % | 18 | 24 % |
| College | 7 | 7.3 % | 0 | 0 |
| Total respondent | 75 | 100 % | 75 | 100 % |
| Duration of treatment: | | | | |
| 1-2 months | 13 | 17.3 % | 34 | 45.3 % |
| 2-3 months | 37 | 49.3 % | 17 | 22.7 % |
| 3-4 months | 25 | 33.3 % | 24 | 32 % |
| Total Respondent | 75 | 100 % | 75 | 100 % |

Table 2 Distribution quality of life (physical health domain)

| No | Physical Health Domain | Treatment Group | | | | Control group | | | |
|----------------------------------|------------------------|-----------------|--------|-----------|--------|---|--------|-----------|--------|
| | | Pre test | | Post Test | | Pre test | | Post Test | |
| | | f | % | f | % | f | % | f | % |
| 1 | Had sufficient | 15 | 20 % | 10 | 13.3 % | 18 | 24 % | 17 | 22.7 % |
| 2 | Sufficient | 18 | 24 % | 12 | 16 % | 35 | 46.7 % | 32 | 42.7 % |
| 3 | Good | 23 | 30.7 % | 29 | 38.7 % | 15 | 15 % | 19 | 25.3 % |
| 4 | Very good | 19 | 25.3 % | 24 | 32 % | 7 | 9.3 % | 7 | 9.3 % |
| | Total | 75 | 100 % | 75 | 100 % | 75 | 100 % | 75 | 100 % |
| <i>Wilcoxon Signed Rank Test</i> | | $P = 0.003$ | | | | $P = 0.317$ | | | |
| <i>Mann-Whitney Test</i> | | | | | | Pretest $p = 0.167$ Posttest $p = 0.000$ | | | |

Table 3 Distribution quality of life (psychological domain)

| No | Psychological Domain | Treatment Group | | | | Control group | | | |
|----------------------------------|----------------------|-----------------|--------|-----------|-------|---|--------|-----------|-------|
| | | Pre test | | Post Test | | Pre test | | Post Test | |
| | | f | % | f | % | f | % | f | % |
| 1 | Less stable | 12 | 16 % | 9 | 12 % | 15 | 20 % | 12 | 16 % |
| 2 | Fairly stable | 27 | 36 % | 15 | 20 % | 27 | 36 % | 27 | 36 % |
| 3 | Stable | 25 | 33.3 % | 30 | 40 % | 20 | 26.7 % | 21 | 28 % |
| 4 | Very stable | 16 | 21.3 % | 21 | 28 % | 13 | 17.3 % | 15 | 20 % |
| | Total | 75 | 100 % | 75 | 100 % | 75 | 100 % | 75 | 100 % |
| <i>Wilcoxon Signed Rank Test</i> | | $P = 0.003$ | | | | $P = 0.317$ | | | |
| <i>Mann-Whitney Test</i> | | | | | | Pretest $p = 0.167$ Posttest $p = 0.000$ | | | |

Table 4 Distribution quality of life (social domain)

| No | Social domain | Treatment Group | | | | Control group | | | |
|----------------------------------|-------------------|-----------------|--------|-----------|--------|---|--------|-----------|--------|
| | | Pre test | | Post Test | | Pre test | | Post Test | |
| | | f | % | f | % | f | % | f | % |
| 1 | Not active enough | 12 | 16 % | 5 | 6.6 % | 23 | 30.7 % | 20 | 26.7 % |
| 2 | Quite active | 29 | 38.7 % | 20 | 26.7 % | 30 | 40% | 30 | 40% |
| 3 | Active | 27 | 36 % | 42 | 56% | 17 | 22.7 % | 20 | 26.7 % |
| 4 | Very active | 7 | 9.3 % | 8 | 10.7 % | 5 | 6.6 % | 5 | 6.6 % |
| | Total | 75 | 100 % | 75 | 100 % | 75 | 100 % | 75 | 100 % |
| <i>Wilcoxon Signed Rank Test</i> | | $P = 0.046$ | | | | $P = 0.157$ | | | |
| <i>Mann-Whitney Test</i> | | | | | | Pretest $p = 0.061$ Posttest $p = 0.015$ | | | |

Table 5 Distribution quality of life (environment domain)

| No | Social domain | Treatment Group | | | | Control group | | | |
|----------------------------------|-----------------|-----------------|--------|-----------|--------|---|--------|-----------|-------|
| | | Pre test | | Post Test | | Pre test | | Post Test | |
| | | f | % | f | % | f | % | f | % |
| 1 | Inadequate | 12 | 16 % | 12 | 6.6 % | 17 | 22.7 % | 15 | 20 % |
| 2 | Enough adequate | 32 | 42.7 % | 30 | 36 % | 39 | 52 % | 43 | 56 % |
| 3 | Adequate | 23 | 30.7 % | 25 | 42.6 % | 19 | 25.3 % | 18 | 24 % |
| 4 | Very adequate | 8 | 10.7 % | 8 | 14.7 % | 0 | 0 % | 0 | 0 % |
| | Total | 75 | 100 % | 75 | 100 % | 75 | 100 % | 75 | 100 % |
| <i>Wilcoxon Signed Rank Test</i> | | $P = 1.000$ | | | | $P = 0.564$ | | | |
| <i>Mann-Whitney Test</i> | | | | | | Pretest $p = 0.101$ Posttest $p = 0.167$ | | | |

IV. DISCUSSION

Psychologically, and socially for pulmonary tuberculosis sufferers at the Community Health Center in Surabaya. There was no significant difference in the environmental domain of quality of life in the pretest and posttest treatment groups using the Wilcoxon test and the Mann Whitney test resulting in H1 being rejected. There was no effect of self-management education on improving quality of life. This shows that the intervention given by the researcher had an effect on the improvement of the quality of life of patients with pulmonary tuberculosis. Pulmonary tuberculosis is one of the deadliest diseases in the world caused by *Mycobacterium Tuberculosis* and is contagious [29]; [30]; [31]. The disease causes health problems in millions of people every year and ranks second to infectious diseases that cause death after the Human Immunodeficiency Virus (HIV) [31]; [26]; [32].

Pulmonary tuberculosis is a threat to the Indonesian population. The disease is a chronic disease that can affect the quality of life of sufferers. Pulmonary tuberculosis raises serious problems in the concept of quality of life consisting of aspects of physical, psychological, social, and environmental health [33]; [34]; [35]. The impact of psychological burden on pulmonary tuberculosis patients will worsen physical health so that it will reduce the quality of life patients. The helplessness of pulmonary tuberculosis patients will lead to changes in adaptation to the psychological, social, and spiritual response so that it will affect the quality of life of sufferers.

This is in accordance with the quality of life measurement data conducted by researchers. In this study, out of 75 respondents in the treatment group, 10 people had sufficient physical health status, 12 people sufficient physical health status, 29 people good physical health status and 24 people very good physical health status. In the control group, 17 people had sufficient health status, 32 people sufficient health status, 19 people good health status and 7 people very good physical health status. Items on physical health included pain (coughing and tightness in pulmonary TB, dependence on drug ingredients and medical assistance, energy and fatigue, mobility, sleep satisfaction, ability to move, and ability to work) [7]; [10]; [12].

For the psychological domain, out of 75 respondents in the treatment group, 9 people had a less stable psychological status, 15 people a fairly stable psychological status, 30 people a stable psychological status and 21 people had a very stable psychological status. [8], [17], [5] state that self-management education is a person's ability to manage strengths greater than oneself. For the control group, 12 people had a less stable psychological status, 27 fairly stable, 21 had a stable status and 15 people had a very stable psychological status. Items in psychological terms included positive feelings, appreciation for life, concentration, body appearance, self-esteem, and negative feelings [17]; [2]

For the social domain, out of the 75 respondents in the treatment group, 5, respondents were not active enough, 20 respondents had quite an active social status, 42 respondents had an active status in the social domain and 8 respondents had a very active social status. In the control group, 20 respondents had a less active social status, 30 respondents had a quite active social status, and 5 respondents had an active social status.

In the environmental domain, out of 75 respondents in the treatment group, 12 people had an inadequate environmental status, 30 people had an adequate enough environmental status, 25 people had an adequate environmental status and 8 people had an adequate environmental domain. In the control group out of 75 respondents in the group, 15 people had an inadequate environmental status, 43 people had an adequate enough environmental status, 18 people had an adequate environmental status and 0 people had an adequate environmental domain.

The treatment group in Table 2 showed an increase in quality of life (physical health domain) in almost all respondents. While the control group did not experience a significant difference, because, based on research

conducted by (2), (17) and (38), in the article it has been proven that self-management education affects the improvement of individual health. This supports the result that the control group that was only given standard treatment in the form of OAT alone without accompanied self-management education measures found no significant differences in aspects of physical health.

The treatment group in Table 3 showed an increase in quality of life (psychological domain). Out of 75 respondents in the treatment group, 9 people had a less stable psychological domain status, 15 people had a fairly stable psychological domain status, 30 people had a stable psychological domain status and 21 people had a stable psychological domain status. In the control group, out of 75 respondents, 12 people had a less stable psychological domain status, 27 people had a fairly stable psychological domain status, 21 people had a stable psychological domain status and 15 people had a stable psychological domain status. Self-management education is education to manage the ability of the self effectively to improve the welfare of the respondent's life where this action will be more focused on the ability of the self to understand the illness, transmission, prevention and treatment that must be engaged in by TB lung sufferers so that TB lung sufferers are able to accept conditions of the illness; good conditions and acceptance will improve the quality of life [8]; [36] The control group did not experience a significant difference, because the psychological aspects of pulmonary tuberculosis patients who were only given standard therapy by the puskesmas showed no significant changes. This will cause psychological resistance which will lead to low psychological well-being [32];

The treatment group in Table 4 showed an improvement in quality of life (social domain) for almost all respondents to the active social domain. Social welfare is a system of life and social life, both material and spiritual, which includes a sense of safety, decency, and inner peace to carry out the fulfillment of physical, spiritual, and social needs as well as possible for themselves, families, and society by upholding the rights and human rights obligations [37]; [7]. This supports that education is one of the important aspects in the realization of social welfare. Educational aspects can be built by growing 5 aspects in a person, namely: knowing, understanding, understanding, analysis and synthesis. These five aspects are requirements that must be built when taking self-management education actions.

The treatment group in Table 5 did not show a significant improvement in quality of life (environmental domain). The environmental domain consisted of financial resources, freedom, security, physical comfort, health, and social care (accessibility and quality), home environment, opportunities to obtain new information and skills, participation and opportunities for recreation, and physical environment (pollution and noise) [38]; [39]. Financial difficulties are a component that is not influenced by actions of self-management education. This is based on the understanding that self-management education is education given to pulmonary TB patients so that patients are able to manage their abilities [32]; [5]. The target of self-management education is to increase the intensity of emotions, and to have a dramatic impact on both emotional and physical changes [29]; [8]. Self-management education does not include actions to overcome one's finances [8]; [32]. Self-management education that has been undertaken will cause emotional strength for patients, so patients will positively receive pulmonary tuberculosis through determination, hope of recovery, and be able to make decisions. This will improve the quality of life of tuberculosis patients, with indicators of improvement in physical, social, and psychological aspects.

V. CONCLUSIONS

Self management education can lead to change in the quality of life in the physical health domain, psychological domain, social domain but not in the environmental domain. Self-management education can be used as an alternative method to improve the independent care of people with pulmonary tuberculosis at home.

CONFLICT OF INTEREST

No conflicts of interest have been declared.

ACKNOWLEDGEMENT

The authors would like to thank the Rector of the University Nahdlatul Ulama Surabaya for funding for the research and Puskesmas Perak Timur, Puskesmas Sawahan, Puskesmas Kebonsari in Surabaya City which were the research sites.

REFERENCES

- [1] F. Ambaw, R. Mayston, C. Hanlon, G. Medhin, and A. Alem, 'Untreated depression and tuberculosis treatment outcomes, quality of life and disability, Ethiopia', *Bull. World Health Organ.*, vol. 96, no. 4, pp. 243–255, 2018, doi: 10.2471/BLT.17.192658.
- [2] B. Chang, A. W. Wu, N. N. Hansel, and G. B. Diette, 'Quality of life in tuberculosis: A review of the English language literature', *Qual. Life Res.*, vol. 13, no. 10, pp. 1633–1642, 2004, doi: 10.1007/s11136-004-0374-1.
- [3] N. Guo, F. Marra, and C. A. Marra, 'Measuring health-related quality of life in tuberculosis: A systematic review', *Health Qual. Life Outcomes*, vol. 7, pp. 1–10, 2009, doi: 10.1186/1477-7525-7-14.
- [4] C. F. Medicine, A. H. Services, D. H. Authority, A. H. Services, and A. Dhabi, 'and Quality of Life in Diabetes ? A Systematic Review of Randomised Controlled Trials', vol. 14, no. 2, pp. 31–43, 2016.
- [5] M. Magharei, S. Jaafari, P. Mansouri, A. Safarpour, and S. A. Taghavi, 'Effects of self-management education on self-efficacy and quality of life in patients with ulcerative colitis: A randomized controlled clinical trial', *Int. J. Community Based Nurs. Midwifery*, vol. 7, no. 1, pp. 32–42, 2019, doi: 10.30476/IJCBNM.2019.40844.32.
- [6] M. Shahhamzeh, F. H. T. K. Kabir, A. Montazeri, A. S. M, and V. Saei, 'The Relationship between Self-Management and Quality of Life in Epileptic Patients who referred to Iranian Epilepsy Association', *مراجعه کننده به انجمن صرع ایرنا ارتباط میان خودمدیریتی و کیفیت زندگی رد بیماران مبتلا به صرع*, vol. 19, no. 2, pp. 189–198, 2017.
- [7] A. Sule *et al.*, 'Quality of Life of Patients with Tuberculosis in a Nigerian Teaching Hospital', *Turkish J. Fam. Med. Prim. Care*, vol. 8, no. 2, p. 39, 2014, doi: 10.5455/tjfmpe.46982.
- [8] S. H. Yu, A. M. Guo, and X. J. Zhang, 'Effects of self-management education on quality of life of patients with chronic obstructive pulmonary disease', *Int. J. Nurs. Sci.*, vol. 1, no. 1, pp. 53–57, 2014, doi: 10.1016/j.ijnss.2014.02.014.
- [9] N. Sharma, P. Sharma, N. Sharma, and R. R. Wavare, 'A cross sectional study of knowledge , attitude and practices of menstrual hygiene among medical students in north India', *J. Phytopharm.*, vol. 2, no. 5, pp. 28–37, 2013.
- [10] A. Muhammad *et al.*, 'Impacto del tratamiento de la tuberculosis en la calidad de vida relacionada con la salud de los pacientes con tuberculosis pulmonar: un estudio de seguimiento', *Health Qual. Life Outcomes*, vol. 12, no. 1, p. 19, 2014, doi: 10.1186/1477-7525-12-19.
- [11] R. Navicharern, 'Diabetes self-management, fasting blood sugar and quality of life among type 2 diabetic patients with foot ulcers', *J. Med. Assoc. Thail.*, vol. 95, no. 2, pp. 156–162, 2012.
- [12] S. Xu, Z. Zhang, A. Wang, J. Zhu, H. Tang, and X. Zhu, 'Effect of Self-efficacy Intervention on Quality of Life of Patients with Intestinal Stoma', *Gastroenterol. Nurs.*, vol. 41, no. 4, pp. 341–346, 2018, doi: 10.1097/SGA.0000000000000290.
- [13] C. E. Fryer, J. A. Luker, M. N. McDonnell, and S. L. Hillier, 'Self-management programs for quality of life in people with stroke', *Cochrane Database Syst. Rev.*, vol. 2013, no. 3, pp. 266–267, 2013, doi: 10.1002/14651858.CD010442.
- [14] J. L. Iti, R. Mudaraddi, S. B. Nagaraja, and S. Shastri, 'Deciphering the Quality of Life among Tuberculosis Patients under RNTCP in Karnataka, India', *J. Tuberc. Res.*, vol. 07, no. 02, pp. 45–55, 2019, doi: 10.4236/jtr.2019.72005.
- [15] Y. Estuningsih, T. N. Rochmah, M. Andriani, and T. Mahmudiono, 'Effect of Self-Regulated Learning for Improving Dietary Management and Quality of Life in Patients with Type-2 Diabetes Mellitus at Dr. Ramelan Naval Hospital, Surabaya, Indonesia', *Kesmas Natl. Public Heal. J.*, vol. 14, no. 2, pp. 51–57, 2019, doi: 10.21109/kesmas.v14i2.2257.

- [16] C. T. Li, K. H. Chu, B. Reiher, T. Kienene, and L. Y. Chien, 'Evaluation of health-related quality of life in patients with tuberculosis who completed treatment in Kiribati', *J. Int. Med. Res.*, vol. 45, no. 2, pp. 610–620, 2017, doi: 10.1177/0300060517694491.
- [17] P. A. Grady and L. L. Gough, 'Self-management: A comprehensive approach to management of chronic conditions', *Am. J. Public Health*, vol. 108, no. 8, pp. S430–S436, 2018, doi: 10.2105/AJPH.2014.302041.
- [18] E. Of *et al.*, 'B Elitung N Ursing Education On Self-Efficacy And Effect Of A Workbook In Health Education On Self-', vol. 5, no. 6, 2019.
- [19] H. H. Tung, C. Y. Lin, K. Y. Chen, C. J. Chang, Y. P. Lin, and C. H. Chou, 'Self-management intervention to improve self-care and quality of life in heart failure patients', *Congest. Hear. Fail.*, vol. 19, no. 4, pp. 9–16, 2013, doi: 10.1111/chf.12014.
- [20] P. Nahid *et al.*, *Treatment of drug-resistant tuberculosis an official ATS/CDC/ERS/IDSA clinical practice guideline*, vol. 200, no. 10, 2019.
- [21] I. Mousa, R. Ataba, K. Al-ali, A. Alkaiyat, and S. H. Zyoud, 'Dialysis-related factors affecting self-efficacy and quality of life in patients on haemodialysis: a cross-sectional study from Palestine', *Ren. Replace. Ther.*, vol. 4, no. 1, pp. 1–12, 2018, doi: 10.1186/s41100-018-0162-y.
- [22] M. Malik, R. Nasir, and A. Hussain, 'Health Related Quality of Life among TB Patients: Question Mark on Performance of TB DOTS in Pakistan', *J. Trop. Med.*, vol. 2018, pp. 1–7, 2018, doi: 10.1155/2018/2538532.
- [23] P. Sangprasert, S. Palangrit, N. Tiyoa, and J. Pattaraarchachai, 'Effects of mindfulness-based health education practice on health behaviors and quality of life among hypertensive patients: A quasi-experimental research', *J. Heal. Res.*, vol. 33, no. 3, pp. 186–196, 2019, doi: 10.1108/JHR-07-2018-0059.
- [24] A. A. S. Jaber, A. H. Khan, S. A. S. Sulaiman, N. Ahmad, and M. S. Anaam, 'Evaluation of health-related quality of life among tuberculosis patients in two cities in Yemen', *PLoS One*, vol. 11, no. 6, pp. 1–19, 2016, doi: 10.1371/journal.pone.0156258.
- [25] Y. Sun *et al.*, 'Development and validation of the pulmonary tuberculosis scale of the system of Quality of Life Instruments for Chronic Diseases (QLICD-PT)', *Health Qual. Life Outcomes*, vol. 16, no. 1, pp. 1–10, 2018, doi: 10.1186/s12955-018-0960-5.
- [26] E. W. M. A. Bischoff, R. Akkermans, J. Bourbeau, C. Van Weel, J. H. Vercoulen, and T. R. J. Schermer, 'Comprehensive self management and routine monitoring in chronic obstructive pulmonary disease patients in general practice: Randomised controlled trial', *BMJ*, vol. 345, no. 7885, pp. 1–12, 2012, doi: 10.1136/bmj.e7642.
- [27] M. Bauer *et al.*, 'Health-related quality of life and tuberculosis: A longitudinal cohort study', *Health Qual. Life Outcomes*, vol. 13, no. 1, 2015, doi: 10.1186/s12955-015-0250-4.
- [28] S. Salehitali, K. Noorian, M. Hafizi, and A. H. Dehkordi, 'Quality of life and its effective factors in tuberculosis patients receiving directly observed treatment short-course (DOTS)', *J. Clin. Tuberc. Other Mycobact. Dis.*, vol. 15, p. 100093, 2019, doi: 10.1016/j.jctube.2019.100093.
- [29] A. A. G. C. Jonker, H. C. Comijs, K. C. P. M. Knipscheer, and D. J. H. Deeg, 'Promotion of self-management in vulnerable older people: A narrative literature review of outcomes of the Chronic Disease Self-Management Program (CDSMP)', *Eur. J. Ageing*, vol. 6, no. 4, pp. 303–314, 2009, doi: 10.1007/s10433-009-0131-y.
- [30] V. Voncken-Brewster, A. Moser, T. Van Der Weijden, Z. Nagykaladi, H. De Vries, and H. Tange, 'Usability evaluation of an online, tailored self-management intervention for chronic obstructive pulmonary disease patients incorporating behavior change techniques', *J. Med. Internet Res.*, vol. 15, no. 1, pp. 1–11, 2013, doi: 10.2196/resprot.2246.
- [31] M. S. Awan, M. Waqas, and M. A. Aslam, 'Factors influencing quality of life in patients with active tuberculosis in Pakistan', *World Appl. Sci. J.*, vol. 18, no. 3, pp. 328–331, 2012, doi: 10.5829/idosi.wasj.2012.18.03.3350.
- [32] A. Deribew *et al.*, 'Tuberculosis and HIV co-infection: Its impact on quality of life', *Health Qual. Life Outcomes*, vol. 7, pp. 1–7, 2009, doi: 10.1186/1477-7525-7-105.
- [33] M. A. A. Diaz, T. N. Huff, and C. R. Libertain, 'Nontuberculous mycobacterial infections of the lower extremities: A 15-year experience', *J. Clin. Tuberc. Other Mycobact. Dis.*, vol. 15, p. 100091, 2019, doi: 10.1016/j.jctube.2019.100091.
- [34] M. Borji, M. Otaghi, and S. Kazembeigi, 'The impact of Orem's self-care model on the quality of life in patients with type II diabetes', *Biomed. Pharmacol. J.*, vol. 10, no. 1, pp. 213–220, 2017, doi: 10.13005/bpj/1100.
- [35] G. C. Donaldson, T. A. R. Seemungal, A. Bhowmik, and J. A. Wedzicha, 'Relationship between

- exacerbation frequency and lung function decline in chronic obstructive pulmonary disease (Thorax (2002) 57, (847-852))', *Thorax*, vol. 63, no. 8, p. 753, 2008, doi: 10.1136/thorax.57.10.847corr1.
- [36] E. Monninkhof, P. van der Valk, J. van der Palen, C. van Herwaarden, and G. Zielhuis, 'Effects of a comprehensive self-management programme in patients with chronic obstructive pulmonary disease', *Eur. Respir. J.*, vol. 22, no. 5, pp. 815–820, 2003, doi: 10.1183/09031936.03.00047003.
- [37] N. Guo, F. Marra, and C. A. Marra, 'Measuring health-related quality of life in tuberculosis: A systematic review', *Health Qual. Life Outcomes*, vol. 7, no. March, 2009, doi: 10.1186/1477-7525-7-14.
- [38] J. S. Louw, M. Mabaso, and K. Peltzer, 'Change in health-related quality of life among pulmonary tuberculosis patients at primary health care settings in South Africa: A prospective cohort study', *PLoS One*, vol. 11, no. 5, pp. 1–13, 2016, doi: 10.1371/journal.pone.0151892.
- [39] A. A. S. Jaber, A. H. Khan, and S. A. S. Sulaiman, 'Evaluating Health-Related Quality of Life of Lost to Follow-Up Tuberculosis Patients in Yemen', *World Acad. Sci. Eng. Technol.*, vol. 13, no. 2, pp. 83–87, 2019.