

## **SURAT KETERANGAN**

Nomor:1052/UNUSA-LPPM/Adm.I/VI/2022

Lembaga Penelitian dan Pengabdian Kepada Masyarakat (LPPM) Universitas Nahdlatul Ulama Surabaya menerangkan telah selesai melakukan pemeriksaan duplikasi dengan membandingkan artikel-artikel lain menggunakan perangkat lunak **Turnitin** pada tanggal 15 Juni 2022.

Judul : *Weight Gain During Pregnancy to Childbirth*

Penulis : Uliyatul Laili, Ratna Ariesta Dwi Andriani

No. Pemeriksaan : 2022.06.20.329

Dengan Hasil sebagai Berikut:

**Tingkat Kesamaan diseluruh artikel (*Similarity Index*) yaitu 19%**

Demikian surat keterangan ini dibuat untuk digunakan sebagaimana mestinya.

Surabaya, 20 Juni 2022

Ketua LPPM



UNUSA  
LPPM

Achmad Syafiuddin, Ph.D

NPP: 20071300

**LPPM Universitas Nahdlatul Ulama Surabaya**

Website : [lppm.unusa.ac.id](http://lppm.unusa.ac.id)

Email : [lppm@unusa.ac.id](mailto:lppm@unusa.ac.id)

Hotline : 0838.5706.3867

# Weight Gain During Pregnancy to Childbirth

*by* Ratna Ariesta Dwi Andriani

---

**Submission date:** 15-Jun-2022 01:38PM (UTC+0700)

**Submission ID:** 1857173222

**File name:** Weight\_Gain\_During\_Pregnancy\_toChildbirth.pdf (245.75K)

**Word count:** 1922

**Character count:** 10039

## **Weight Gain During Pregnancy to Childbirth**

**Uliyatul Laili\*, Ratna Ariesta Dwi Andriani\***

\* Fakultas Keperawatan dan Kebidanan, Universitas Nahdlatul Ulama Surabaya, Surabaya, Indonesia

---

### **Article Info**

#### **Article history:**

Received Apr 28<sup>th</sup>, 2020

Revised May 29<sup>th</sup>, 2020

Accepted Jul 29<sup>th</sup>, 2020

#### **Keyword:**

First keyword

Second keyword

Third keyword

Fourth keyword

Fifth keyword

### **ABSTRACT (10 PT)**

Increased energy and nutrients are needed for fetal growth and development, increase in the size of the uterine organs, as well as changes in the composition and metabolism of the mother's body. The nutritional status of pregnant women can be seen through several factors including the adjusted body mass index (BMI) of the mother before pregnancy which will affect how much the ideal increase in pregnancy during pregnancy, including the size of the LLA (Upper Arm Circumference) to achieve an ideal state. It aims to prevent weight gain that is too much or too little for pregnant women who will also affect the birth process and the condition of the fetus being born. The purpose of this study was to analyze the relationship between maternal weight gain during pregnancy and childbirth. The research method used is quantitative, with cross sectional approach. The study was conducted at the Jagir Community Health Center in Surabaya, with a sample of 80 people. Data obtained using SPSS with chi square test. The results showed there was a relationship between maternal weight gain during pregnancy which was seen based on body mass index (BMI) with the labor process that took place. In the implementation of midwifery care for pregnant women nutritional status monitoring and weight gain are monitored since the beginning of pregnancy so that labor takes place smoothly

*Copyright © Jurnal Teknologi Kesehatan (Journal of Health Technology).  
All rights reserved.*

---

### **Corresponding Author:**

Ratna Ariesta Dwi Andriani,  
Fakultas Keperawatan dan Kebidanan,  
Universitas Nahdlatul Ulama Surabaya,  
Surabaya, Indonesia.  
Email: ratnariesta@unusa.ac.id

---

### **1. INTRODUCTION**

Pregnancy is a change in order to continue the offspring that occur naturally, producing a fetus that grows in the mother's womb, and then can be explained by the growth rate and size of the fetus according to gestational age, at every pregnancy examination (N & Safe'i, 2010). Pregnancy causes an increase in energy metabolism. Therefore, the need for energy and other nutrients increases during pregnancy. Recommendations for weight gain for pregnant women based on pre-pregnancy BMI according to IOM are: BMI <18.5 kg / m<sup>2</sup> gain weight 28-40 pounds (equivalent to 12.6-18 kg), BMI 18.5-24.9 kg / m<sup>2</sup> with weight gain 25-35 pounds (equivalent to 11.25-15.75kg), IMT 25-29.9 kg / m<sup>2</sup> with weight gain of 15-25 pounds (equivalent to 6.75-11.25kg) and BMI 30 kg / m<sup>2</sup> with weight gain of 11-20 pounds (equivalent to 4.95-9kg). (Eva et al., 2007) The results of other studies also indicate that prahamil nutritional status is related to infant weight and body length. Women who have a BMI <26.0 kg / m<sup>2</sup> will have a baby with a body length of 1.5 cm more than women who have a BMI <19.8 kg / m<sup>2</sup>, as well as research at Rajavithi Hospital, Thailand obtained data that underweight women group tend to give birth to LBW babies and premature, obese group mothers give birth to macrosomia / large babies (Kliranayungie & Debtarsie, 2012).

### **2. RESEARCH METHOD**

---

**Journal homepage:** <http://e-journal.poltekkesjogja.ac.id/index.php/JTK>

The research method used is quantitative with cross sectional approach. This research was conducted at the Jagir Puskesmas in Surabaya by looking at the Maternal and Child Health (MCH) book and puskesmas medical record data. The population in this study was maternity in the Jagir Health Center in May-June 2019, with a total sample of 80 people. Then the data were analyzed using SPSS with the chi square test.

### 3. RESULTS

The results of this study showed that the majority of mothers during their pregnancy experienced a weight gain that was not in accordance with the Body Mass Index (BMI) calculated at the beginning of pregnancy, totaling 47 (59%) respondents. Respondents in this study, as many as 46 (58%) of mothers whose labor took place spontaneously, while 34 (42%) of the other respondents the delivery process was carried out by caesarean section. After analyzing the data by using the chi square test, the value of  $p = 0.000$  is obtained, which means there is a relationship between maternal weight gain during pregnancy and childbirth.

Table 1. Frequency distribution based on age, weight gain and labor process

| Variable                |                              | f  | %  |
|-------------------------|------------------------------|----|----|
| Age                     | < 20 years old               | 3  | 4  |
|                         | 20 – 35 years old            | 68 | 85 |
|                         | >35 years old                | 9  | 11 |
| Appropriate Weight Gain | Suitable                     | 33 | 41 |
|                         | Not suitable                 | 47 | 59 |
| Delivery process        | Spontaneous delivery process | 46 | 58 |
|                         | Caesarean Sectio             | 34 | 42 |

Based on table 1 shows that of the 80 respondents who participated in this study, most of the 68 respondents (85%) were in the reproductive age range of 20 - 35 years, while for the weight gain some 47 respondents (59%) experienced weight gain which is not in accordance with the ideal weight gain based on the Body Mass Index (BMI) before pregnancy. Research data on the delivery process showed that some 46 (58%) respondents of the birth process took place spontaneously.

Table 2. Cross table of the relationship between weight gain and labor

| Variable                |              | Delivery process             |                  | P value |
|-------------------------|--------------|------------------------------|------------------|---------|
|                         |              | Spontaneous delivery process | Caesarean Sectio |         |
| Appropriate Weight Gain | Suitable     | 19                           | 14               | 0.000   |
|                         | Not suitable | 27                           | 20               |         |
|                         |              | 46                           | 34               | 80      |

Table 2 shows that the majority of respondents who gained weight according to spontaneous labor were 27 respondents. Then performed an analysis using the chi square test obtained  $p$  value  $0.000 > 0.05$  which means there is a relationship between maternal weight gain during pregnancy with childbirth.

### 4. DISCUSSION

Research conducted at the Jagir Community Health Center in Surabaya on 80 respondents showed that there was a relationship between maternal weight gain during pregnancy and childbirth with a  $p$  value of 0,000. Weight gain in pregnant women is measured using the Body Mass Index (BMI) in early pregnancy by looking at body weight and height. Research conducted by Ruth found that mothers with a  $BMI \geq 25 \text{ kg} / \text{m}^2$  are at risk of developing hypertension in pregnancy, tend to have a cesarean section and their babies tend to have low birth weight, death, and asphyxia (Sondakh, Prof, & Manado, 2016).

The results of this study are in accordance with the theory that in pregnant women with excess body weight, it turns out there is an increase and association with outcomes that are not good during perinatal for themselves and the fetus. In women with excess BMI the incidence of disease and other complications increases such as gestational diabetes, macrosomia, and caesarean section delivery. In women with  $BMI > 29 \text{ g} / \text{m}^2$ , there is an increase of 2-4 times the possibility of caesarean section action (Rusli, Ocviyanti, & Mansyur, n.d.). The delivery process is influenced by several factors including the passage, passenger, power, position, psychology. In addition, at the time of data collection most respondents were of reproductive age that is 20-35 years. So that most respondents are not at a high risk score because of the age factor when referring to the Poedji Rahayu Score Card (KSPR). The age factor can be one of the considerations in making decisions to determine the delivery process to be performed. If a pregnant

woman is over 35 years old, it can be an alternative part of caesarean delivery, because the mother is at a high risk score and during labor there can be some complications if labor is done normally / spontaneously such as maternal labor factors that can result in the first stage or the second time elongated, asphyxiated baby.

During the data collection process, the researchers found that some women who gave birth in a caesarean section were those who had an elongated first stage. Research conducted by Purwaningsih and Fatmawati (2010) shows that labor is not progressing labor with a first-phase latent phase that is more than 8 hours, or crosses the alert line on the partograph sheet and the unborn baby (Purwaningsih, Wahyu, & Siti Fatmawati, 2010). In the first stage of labor can cause several complications including infection, dehydration, fetal asphyxia, fatigue and death in the womb. Therefore, the choice of delivery method in an alternative section caesarea choice. These results are in line with research conducted by Aprina and Anita (2016) which states that in labor that is not progressing there are indications for labor in a caesarean section. (Aprina & Putri, n.d.) While one of the predisposing factors of parturition or prolonged labor is education, age, parity and maternal psychology as well as fetal factors (abnormalities in the shape or shape of the fetus ie large fetuses; congenital abnormalities). (Soviyati, 2016)(Ardhiyanti et al., 2016) Other studies have shown that maternal parity also contributes to the incidence of prolonged labor (Ardhiyanti et al., 2016).

So, it is necessary to do better midwifery care by health workers, especially midwives, in terms of monitoring the process of pregnancy. At the beginning of pregnancy it is necessary to measure the Body Mass Index (BMI) of pregnant women with the aim of being able to properly monitor maternal weight gain during pregnancy so as not to be less or more by taking into account the estimated fetal weight that is adjusted to the gestational age.

## 5. CONCLUSION

The conclusion of this study is that the majority of respondents are of reproductive age, that is 20 - 35 years. And from the results of the analysis obtained that there is a relationship between weight gain in pregnant women which is measured based on the Body Mass Index (BMI) at the beginning of pregnancy with the birth process.

## REFERENCES (10 PT)

- [1] N, M., & Safe'i. (2010). *Panduan Lengkap Ibu Hamil, Khusus Ibu Hamil*. Jakarta: Power book.
- [2] Aprina, & Putri, A. (n.d.). Faktor-faktor yang berhubungan dengan persalinan sectio caesarea di rsud dr. h abdul moelock provinsi lampung. *Jurnal Kesehatan*, VII(1), 90–96.
- [3] Ardhiyanti, Y., Susanti, S., Studi, P., Sekolah, K., Ilmu, T., Hang, K., & Pekanbaru, T. (2016). *Faktor Ibu yang Berhubungan dengan Kejadian Persalinan Lama di RSUD Arifin Achmad Pekanbaru Factors of The Mother Related to Obstructed Labour Case at RSUD Arifin Achmad Pekanbaru*. 3(2), 83–87.
- [4] Eva, D., Endang, A., Anies, L. A., Gizi, D., Masyarakat, K., Kesehatan, F., & Universitas, M. (2007). *Kekurangan Energi Kronis pada Wanita Indonesia Validity Mid-Upper Arm Circunference to Detect Chronic Energy Malnutrition Risk of Indonesian Women*. 83–90. Retrieved from <http://journal.fkm.ui.ac.id/kesmas/article/view/67/68>
- [5] Kliranayungie, & Debtarsie, C. (2012). *Universitas indonesia*. (September 2011).
- [6] N, M., & Safe'i. (2010). *Panduan Lengkap Ibu Hamil, Khusus Ibu Hamil*. Jakarta: Power book.
- [7] Purwaningsih, Wahyu, & Siti Fatmawati. (2010). *Asuhan Keperawatan Maternitas*. Yogyakarta: Nuha Medika.
- [8] Rusli, A. I., Oeviyanti, D., & Mansyur, M. (n.d.). *Hubungan Peningkatan Berat Badan Ibu Hamil Dengan Lama Persalinan*
- [9] Sondakh, J. M. M., Prof, R., & Manado, R. D. K. (2016). Hubungan indeks massa tubuh ( IMT ) awal kehamilan dengan luaran maternal neonatal. *Jurnal E-Clinic (ECI)*, 4(2).

**BIOGRAPHY OF AUTHORS (10 PT)**

|                                 |             |
|---------------------------------|-------------|
| First author's<br>Photo (3x4cm) | Xxxx (9 pt) |
| Second author's<br>photo(3x4cm) | Xxxx (9 pt) |
| Thirth author's<br>photo(3x4cm) | Xxxx (9 pt) |

# Weight Gain During Pregnancy to Childbirth

---

ORIGINALITY REPORT

---

19%

SIMILARITY INDEX

16%

INTERNET SOURCES

8%

PUBLICATIONS

6%

STUDENT PAPERS

---

MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

---

4%

★ [www.cldcollege.com](http://www.cldcollege.com)

Internet Source

---

Exclude quotes  On

Exclude matches  < 10 words

Exclude bibliography  On