

SURAT KETERANGAN

Nomor: 2448/UNUSA-LPPM/Adm.I/XI/2022

Lembaga Penelitian dan Pengabdian Kepada Masyarakat (LPPM) Universitas Nahdlatul Ulama Surabaya menerangkan telah selesai melakukan pemeriksaan duplikasi dengan membandingkan artikel-artikel lain menggunakan perangkat lunak **Turnitin** pada tanggal 18 November 2022.

Judul : Effects Of Coffee On Prevention Of Oxidative Stress And
Inflammation Induced By Cigarette Smoke In Male Rats

Penulis : Muhamad Khafid

No. Pemeriksaan : 2022.11.18.1030

Dengan Hasil sebagai Berikut:

Tingkat Kesamaan diseluruh artikel (*Similarity Index*) yaitu 13%

Demikian surat keterangan ini dibuat untuk digunakan sebagaimana mestinya.

Surabaya, 18 November 2022

Ketua LPPM



UNUSA
LPPM

Achmad Syafiuddin, Ph.D

NPP: 20071300

LPPM Universitas Nahdlatul Ulama Surabaya

Website : lppm.unusa.ac.id

Email : lppm@unusa.ac.id

Hotline : 0838.5706.3867

EFFECTS OF COFFEE ON PREVENTION OF OXIDATIVE STRESS AND INFLAMMATION INDUCED BY CIGARETTE SMOKE IN MALE RATS

by Muhammad Khafid

Submission date: 18-Nov-2022 11:00AM (UTC+0700)

Submission ID: 1957436963

File name: EFFECTS_OF_COFFEE_ON_PREVENTION_OF_OXIDATIVE_STRESS.pdf (22.38K)

Word count: 327

Character count: 1756

**EFFECTS OF COFFEE ON PREVENTION OF OXIDATIVE STRESS AND
INFLAMMATION INDUCED BY CIGARETTE SMOKE IN MALE RATS**

(*Rattus norvegicus*)

M. Khafid

**Faculty of Nursing and Midwifery
University of Nahdlatul Ulama Surabaya**

Abstract

Free radicals in cigarette smoke are known to cause oxidative stress. Free radical activity can be inhibited by antioxidants such as chlorogenic acid inside the coffee. The purpose of this study was analyzes whether the administration of coffee may help prevent oxidative stress and inflammation in male rats (*Rattus norvegicus*) exposed by cigarette smoke.

The research design was posttest only control group design. The total sample size of 5 group was 30 male rats (*Rattus norvegicus*) weighing 200-250 grams. The Independent variables were cigarette smoke and coffee, while the dependent variables were MDA and TNF- α . Normality test data by Shapiro Wilks was esed ($\alpha = 0.05$) and homogeneity by Levene test was used($\alpha = 0.05$). Test different between groups by Brown Forsyte ($\alpha = 0.05$) and continued with Games Howell test ($\alpha = 0.05$).

Test results of Brown Forsyte on MDA showed significant difference between groups ($p = 0.001$), and Games Howell test result showed there were significant differences between the levels of MDA between group (non white + coffee) with a control group, a group (cigarette + coffee) with the (non white + coffee). TNF- α showed a significant difference ($p = 0.000$) levels as well. TNF α between groups and by using test Games Howel there are significant differences between the control group and group exposed by mild/white cigarettes, while the (cigarette + coffee) differ significantly from the cigarette group and white/mild cigarettes group.

To sum up, the provision of coffee in a dose of 1.35 mg / g bw / day has not been able to prevent oxidative stress, but it can prevent the inflammation caused by cigarette smoke exposure.

Keywords: smoke, coffee, MDA and TNF- α

EFFECTS OF COFFEE ON PREVENTION OF OXIDATIVE STRESS AND INFLAMMATION INDUCED BY CIGARETE SMOKE IN MALE RATS

ORIGINALITY REPORT

13%

SIMILARITY INDEX

7%

INTERNET SOURCES

0%

PUBLICATIONS

6%

STUDENT PAPERS

MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

6%

★ Submitted to Universitas Negeri Jakarta

Student Paper

Exclude quotes On

Exclude matches < 2%

Exclude bibliography On