

SURAT KETERANGAN

Nomor: 2753/UNUSA-LPPM/Adm.I/XII/2022

Lembaga Penelitian dan Pengabdian Kepada Masyarakat (LPPM) Universitas Nahdlatul Ulama Surabaya menerangkan telah selesai melakukan pemeriksaan duplikasi dengan membandingkan artikel-artikel lain menggunakan perangkat lunak **Turnitin** pada tanggal 27 November 2022.

Judul : Proinflammatorycytokines profile On Skin Aging In Jemursari
Islamic Hospital

Penulis : Winawati Eka P, Meidyta S, Hotimah

No. Pemeriksaan : 2022.12.23.1122

Dengan Hasil sebagai Berikut:

Tingkat Kesamaan diseluruh artikel (*Similarity Index*) yaitu 15%

Demikian surat keterangan ini dibuat untuk digunakan sebagaimana mestinya.

Surabaya, 23 Desember 2022

Ketua LPPM



UNUSA
LPPM

Achmad Syafiuddin, Ph.D

NPP: 20071300

LPPM Universitas Nahdlatul Ulama Surabaya

Website : lppm.unusa.ac.id

Email : lppm@unusa.ac.id

Hotline : 0838.5706.3867

Proinflammatorycytokines profile On Skin Aging In Jemursari Islamic Hospital

by HotimahHotimah

Submission date: 27-Nov-2022 01:37PM (UTC+0700)

Submission ID: 1963932684

File name: Art.Proinflammatory.pdf (214.16K)

Word count: 2125

Character count: 11633

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/352091898>

Proinflammatory cytokines profile On Skin Aging In Jemursari Islamic Hospital

Article · December 2020

CITATIONS
0

READS
33

3 authors, including:



Winawati Eka Putri
Nahdlatul Ulama University of Surabaya Generation prepare Rahmatan lil 'Alamin
9 PUBLICATIONS 3 CITATIONS

[SEE PROFILE](#)



Hotimah Salim
Universitas Nahdlatul Ulama Surabaya
53 PUBLICATIONS 386 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Biomolecular science, Atherosclerosis, Cardio-diabetology, Pharmacology [View project](#)



Herbal Medicine [View project](#)

Proinflammatory cytokines profile On Skin Aging In Jemursari Islamic Hospital

Winawati Eka Putri¹, MeidytaSinantryana², Hotimah³
^{1,2,3}Faculty of Medicine Universitas Nahdlatul Ulama, Surabaya, Indonesia

Date of Submission: 05-12-2020

Date of Acceptance: 20-12-2020

ABSTRACT : Aging is a complex process and make progressive loss of functional capacity of organ such as skin. There is an imbalance of immune system in skin aging. Proinflammatory cytokines such as IL-6 and TNF- α were elevated in skin aging. **Purpose:** This study aims to observe about proinflammatory cytokines on skin aging in Jemursari Islamic Hospital. **Methods:** This research is an observational study with cross sectional design. The IL-6 and TNF- α level was measured by using sandwich ELISA technique. **Results:** The mean level of IL-6 and TNF- α in subjects aged 56-75 years were 19.99 ng/ml and 2.86 ng/ml, respectively. Those were higher than aged 45-55 years. In males group, IL-6 was more elevated than females group. Meanwhile in females group, TNF- α was more elevated than in males group. The mean level of IL-6 and TNF- α in overweight groups were more elevated than normal BMI groups. **Conclusion:** Subjects in Jemursari Islamic hospital had elevated IL-6 and TNF- α level. However, there was needed a much further research in order to find correlation between skin aging and proinflammatory cytokines.

Keywords: Skin aging, IL-6, TNF- α .

I. INTRODUCTION

Aging is an unavoidable and complex process characterized by progressive loss of functional capacity of organ such as skin. This biological phenomenon happens overlapping between internal (genetic, hormonal) and environmental (ultraviolet, pollution) factors (Chung, Cho and Kang, 2004; Assaf, Adly and Hussein, 2010; Minciullo et al., 2016; Garg, Khurana and Garg, 2017). Recently, people in all the world aged 60 years or over are increasing rapidly. This phenomenon will spend high costs of medical care for the elderly (Kovaiou, Weinberger and Grubeck-Loebenstein, 2008).

Manifestation of skin aging both intrinsic and extrinsic include wrinkles, skin atrophy, telangiectasis, solar lentigo, premalignant skin lesions, skin inelastic and rigid, and leathery

appearance (Yaar, 2006; Yaar and Gilchrist, 2012; Tobin, 2017). The immune system of aging or known as immunosenescence undergoes decreases, with major implications for health and survival (Mansouri et al., 2014; Simon, Hollander and McMichael, 2015). 'Inflammaging' has become widely used to describe inflammatory process of aging and is characterized by low-grade, chronic pro-inflammatory state. During this process, the balance of cytokines such as interleukin-6 (IL-6) and tumor necrosis factor- α (TNF- α) are elevated (Brüüngaard and Pedersen, 2003; Milan-Mattos et al., 2019). Interleukin-6 are commonly found in middle aged population than TNF- α which is more found in age 80 years or above (Brüüngaard and Pedersen, 2003). This research was conducted because no study had studied yet about proinflammatory cytokines profile on skin aging patients in Jemursari Islamic Hospital Surabaya.

II. METHODS

This research is an observational study with cross sectional group design. It was conducted at our hospital and was approved by the hospital's ethics committee. The study was comprised of 19 healthy subjects. Anthropometric measurements including weight (in kg) and height (in cm) were measured in all subjects. Body mass index (BMI) was calculated using the formula: weight (kg)/height² (m²). The subjects were eligible for inclusion if they were aged 45 years and above, had skin aging sign in this research we observed any wrinkle both static and dynamic wrinkle. The subjects did not have any past histories of kidney abnormality, hypothyroid, malignancy, hypertension, diabetes mellitus, cardiovascular disease, obstructive biliary and HIV. They also did not have any skin abnormality and did not consume any medication within a month before this research was conducted. IL-6 and TNF- α level measurement were assessed by sandwich ELISA technique in Biochemistry laboratory, Faculty of Medicine, Brawijaya University. The research procedure is as follows:




Figure 2. Research Procedure

Data were collected from the research status, then performed cleaning, editing and coding. The data that was collected were entered in the Statistical Package for the Social Sciences (SPSS) data format version 20.0 (SPSS, Inc., Chicago, Illinois). The main outcome measures in this study were IL-6 and TNF- α level. The statistical test used was the normality test with the Shapiro-Wilk test because the sample number was smaller than 30 per group and the data were normally distributed ($p > 0.05$).

III. RESULTS

Based on this research, the subject distribution based on age was divided into 2 groups, 45-55 years group and 56-65 years group. The mean value of IL-6 in skin aging patients in Jemursari

Islamic Hospital Surabaya were 18,51 ng/ml for ages 45-55 years group and 19,99 ng/ml for ages 56-65 years. The mean value of TNF- α in skin aging patients in Jemursari Islamic Hospital Surabaya were 2,36 ng/ml for ages 45-55 years group and 2,86 ng/ml for ages 56-65 years (Table 1). This study also comprised of 9 (47,4%) males and 10 (52,6%) females. The mean value of IL-6 in males were 19,52 ng/ml and females were 19,09 ng/ml. The mean value of TNF- α for males were 2,48 ng/ml and females were 2,75 ng/ml (Table 2). The BMI of subjects was found in normal value (68,4%) and overweight (31,6%). The mean value of IL-6 in normal group were 18,56 ng/ml and overweight group were 20,89 ng/ml. The mean value of TNF- α in normal group were 2,45 ng/ml and overweight group were 2,99 ng/ml (Table 3).

Table 1. Subject distribution based on age

Age (Year)	N	%	Mean of IL-6 ng/mL	Mean of TNF- α pg/ml
45-55	9	47,4	18,51	2,36
56-65	10	52,6	19,99	2,86
Total	19	100,0		

Tabel 2. Subject distribution based on sex

Sex	N	%	Mean of IL-6 ng/mL	Mean of TNF- α pg/ml
Males	9	47,4	19,52	2,48
Females	10	52,6	19,09	2,75
Total	19	100,0		

Table 3. Subject distribution based on BMI

Body Mass Index	N	%	Mean of IL-6 ng/mL	Mean of TNF- α pg/ml
Underweight	0	0	0	0
Normal	13	68,4	18,56	2,45
Overweight	6	31,6	20,89	2,99
Obese	0	0	0	0
Total	19	100,0		

IV. DISCUSSION

Our results show that the mean value of IL-6 and TNF- α in older group (56-65 years) were found more elevated than younger group (45-55 years). In accordance with ours, Ye et al.(2019) also found that IL-6 and TNF- α were significantly elevated in chronologically aged individuals than young human controls. Lumentut, Marunduh and Rampengan(2015) found that TNF- α level in age 64-74 years and >90 years were more elevated than age 75-90 years. Wu et al.(2007) also found the same results but in mouse adipose tissue not in aging human. The proinflammatory cytokines are regulated by NF- κ B signalling pathway which are activated in aging. This signalling pathway will release proinflammatory cytokines such as IL-6 and TNF- α .

The mean value of IL-6 in males group (56-65 years) in this research were found more elevated than females group (45-55 years). Otherwise mean value of TNF- α in females group were more elevated than males group. These results were not similar with Lumentut, Marunduh and Rampengan (2015) who found that TNF- α level in males group were more elevated than female group. Milan-Mattos et al.(2019) found that IL-6 and TNF- α in 41-70 years females group showed more elevated than males group.

Aging is also related with body composition and it can be measured with BMI. Several studies found that obesity is associated with increased proinflammatory cytokines (Brüünsgaard and Pedersen, 2003). Our resultsshow that the mean value of IL-6 and TNF- α were more elevated than normal BMI group. Azizian et al.(2016) and Ashraf et al.(2018) showed that value of IL-6 and TNF- α were not significantly difference in age 47-49 years old obese and non-obese patients with metabolic syndrome.

V. CONCLUSION

Based on our results, we have shown proinflammatory cytokines changes in skin aging patients in Jemursari Islamic Hospital include IL-6 and TNF- α level. Those cytokines were elevated in older group, males group and overweight group. Our limitation that we did not measure grading of skin aging and next research will find correlation between variables.

REFERENCES

- [1]. Ashraf, H. et al. (2018) 'Evaluation of proinflammatory cytokines in obese vs non-obese patients with metabolic syndrome', Indian journal of endocrinology and metabolism. Wolters Kluwer--Medknow Publications, 22(6), p. 751.
- [2]. Assaf, H., Adly, M. and Hussein, M. (2010) 'Aging and Intrinsic Aging: Pathogenesis and Manifestations', in Farage, M., Miler, K., and Maibach, H. (eds) Textbook of Aging Skin. Berlin: Springer.
- [3]. Azizian, M. et al. (2016) 'Cytokine profiles in overweight and obese subjects and normal weight individuals matched for age and gender', Annals of Clinical Biochemistry. SAGE Publications Sage UK: London, England, 53(6), pp. 663-668.
- [4]. Brüünsgaard, H. and Pedersen, B. K. (2003) 'Age-related inflammatory cytokines and disease', Immunology and Allergy Clinics of North America, 23(1), pp. 15-39. doi: 10.1016/S0889-8561(02)00056-5.
- [5]. Chung, J., Cho, S. and Kang, S. (2004) 'Why Does the Skin Age? Intrinsic Aging, Photoaging, and Their Pathophysiology', in Rigel, D. et al. (eds). New York: Marcel Dekker, Inc.

- [6]. Garg, C., Khurana, P. and Garg, M. (2017) 'Molecular mechanisms of skin photoaging and plant inhibitors', *International Journal of Green Pharmacy (IJGP)*, 11(02), pp. 217–232.
- [7]. Kovaïou, R. D., Weinberger, B. and Grubeck-Loebenstien, B. (2008) 'Aging and the immune system', in *Clinical Immunology*. Elsevier, pp. 503–511.
- [8]. Lumentut, A. R., Marunduh, S. and Rampengan, J. J. V (2015) 'Profil TNF- α pada orang lanjut usia di Panti Wredha Bethania Lembean', 3, pp. 1–4.
- [9]. Mansouri, P. et al. (2014) 'Skin aging and immune system', in Massoud, A. and Rezaei, N. (eds) *Immunology of Aging*. New York: Springer, pp. 339–368.
- [10]. Milan-Mattos, J. C. et al. (2019) 'Effects of natural aging and gender on pro-inflammatory markers', *Brazilian Journal of Medical and Biological Research*, 52(9), pp. 1–10. doi: 10.1590/1414-431x20198392.
- [11]. Minciullo, P. L. et al. (2016) 'Inflammaging and Anti-Inflammaging: The Role of Cytokines in Extreme Longevity', *Archivum Immunologiae et Therapiae Experimentalis*. Springer International Publishing, 64(2), pp. 111–126. doi: 10.1007/s00005-015-0377-3.
- [12]. Simon, A. K., Hollander, G. A. and McMichael, A. (2015) 'Evolution of the immune system in humans from infancy to old age', *Proceedings of the Royal Society B: Biological Sciences*, 282(1821). doi: 10.1098/rspb.2014.3085.
- [13]. Tobin, D. J. (2017) 'Introduction to skin aging', *Journal of Tissue Viability*. Elsevier Ltd, 26(1), pp. 37–46. doi: 10.1016/j.jtv.2016.03.002.
- [14]. Wu, D. et al. (2007) 'Aging Up-Regulates Expression of Inflammatory Mediators in Mouse Adipose Tissue', *The Journal of Immunology*, 179(7), pp. 4829–4839. doi: 10.4049/jimmunol.179.7.4829.
- [15]. Yaar, M. (2006) 'Clinical and Histological Features of Intrinsic versus Extrinsic Skin Aging', in Gilchrist, B. and Krutmann, J. (eds) *Skin Aging*. Berlin: Springer, pp. 9–21.
- [16]. Yaar, M. and Gilchrist, B. (2012) 'Aging of skin', in Goldsmith, L. et al. (eds) *Fitzpatrick's dermatology in general medicine*. 8th ed. New York, NY: McGraw-Hill Professional. New York: McGraw-Hill Companies Inc.
- [17]. Ye, L. et al. (2019) 'Topical applications of an emollient reduce circulating pro-inflammatory cytokine levels in chronically aged humans: a pilot clinical study', *Journal of the European Academy of Dermatology and Venereology*, 33(11), pp. 2197–2201. doi: 10.1111/jdv.15540.

Proinflammatorycytokines profile On Skin Aging In Jemursari Islamic Hospital

ORIGINALITY REPORT

15%

SIMILARITY INDEX

12%

INTERNET SOURCES

12%

PUBLICATIONS

8%

STUDENT PAPERS

MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

3%

★ hull-repository.worktribe.com

Internet Source

Exclude quotes On

Exclude matches < 1%

Exclude bibliography On