

SURAT KETERANGAN

Nomor: 1291/UNUSA-LPPM/Adm-I/VII/2023

Lembaga Penelitian dan Pengabdian Kepada Masyarakat (LPPM) Universitas Nahdlatul Ulama Surabaya menerangkan telah selesai melakukan pemeriksaan duplikasi dengan membandingkan artikel-artikel lain menggunakan perangkat lunak **Turnitin** pada tanggal 13 Juli 2023

Judul : *Observation of children's age to predict maximum height*
Penulis : Sapta Kunta Purnama, Muhammad Furqon Hidayatullah, Fadilah Umar, Slamet Riyadi, Siti Nurunnayah, Eppy Setiyowati, Hilwah Nora, Agus Mukholid
No. Pemeriksaan : 2023.07.13.447

Dengan Hasil sebagai Berikut:

Tingkat Kesamaan diseluruh artikel (*Similarity Index*) yaitu 17%

Demikian surat keterangan ini dibuat untuk digunakan sebagaimana mestinya

Surabaya, 13 Juli 2023

Ketua LPPM,



Achmad Syafiuddin, Ph.D.

NPP. 20071300

LPPM Universitas Nahdlatul Ulama Surabaya

Website : lppm.unusa.ac.id

Email : lppm@unusa.ac.id

Hotline : 0838.5706.3867

Observation of children's age to predict maximum height

by Eppy Setiyowati

Submission date: 13-Jul-2023 07:35AM (UTC+0700)

Submission ID: 2130317165

File name: 2_Artikel_Jurnal_Observation_of_children.pdf (238.16K)

Word count: 4432

Character count: 23101

Observation of children's age to predict maximum height

SAPTA KUNTA PURNAMA¹, MUHAMMAD FURQON HIDAYATULLAH¹, FADILAH UMAR¹ ✉, SLAMET RIYADI¹, SITI NURUNNIYAH², EPPY SETIYOWATI³, HILWAH NORA⁴, AGUS MUKHOLID⁵

¹Department Sports Coaching Education, University of Sebelas Maret, Surakarta, Indonesia

²Midwifery Program, Alma Ata University, Yogyakarta, Indonesia

³Faculty of Nursing and Midwifery, Nahdlatul Ulama University Surabaya, Indonesia

⁴Faculty of Medicine, Syiah Kuala University, Banda Aceh, Indonesia

⁵Department Physical Education, Health, and Recreation, University of Sebelas Maret, Surakarta, Indonesia

ABSTRACT

The ideal height is one of the things that people in Indonesia and around the world want. This study aims to formulate guidelines for predicting the maximum height of men and women. The method used in this research is descriptive. This study observed the growth and development of height for girls aged 0-18 years and boys aged 0-19 years. This research was conducted in the city of Surakarta, Central Java, Indonesia. The research sample was obtained from Posyandu, PAUD, SD, SMP, and SMA. Data were collected by observing the respondent's height. Data analysis used the Miles and Huberman model, namely data reduction, data tabulation, data presentation, and concluding. Measurements for predicting a child's height can be done by considering weight and age percentage. The results showed that predicting a child's height could be done by considering the percentage of weight and age. Every year the growth of the child's height will increase according to the age of growth. Therefore, child height growth data can predict the maximum height of children at the age of 18 years for girls and 19 years for boys. Furthermore, the child height prediction method can be applied to determine the child's height in the future.

Keywords: Physical education, Observation model, Height-for-age, Maximum height, Growth, Development.

Cite this article as:

Purnama, S. K., Hidayatullah, M. F., Umar, F., Riyadi, S., Nurunniyah, S., Setiyowati, E., Nora, H., & Mukholid, A. (2023). Observation of children's age to predict maximum height. *Journal of Human Sport and Exercise*, 18(2), 366-374. <https://doi.org/10.14198/jhse.2023.182.08>

✉ **Corresponding author.** Department Sports Coaching Education, Universitas Sebelas Maret, Surakarta, Indonesia.

<https://orcid.org/0000-0003-3371-2613>

E-mail: hus.dam@yahoo.com

Submitted for publication May 31, 2022.

Accepted for publication June 13, 2022.

Published April 01, 2023 (in press August 05, 2022).

JOURNAL OF HUMAN SPORT & EXERCISE ISSN 1988-5202.

© Faculty of Education, University of Alicante.

doi:10.14198/jhse.2023.182.08

INTRODUCTION

Children should grow and develop naturally following the stages of growth and development of the child. Growth and development are unique physiological phenomena in infants and adolescents (Mascherini et al., 2019). Height and weight are important growth indicators and have attracted considerable attention from the family and society (Reinehr et al., 2020; Wang et al., 2018). Adolescence is related to Puberty (Feka et al., 2019; Silva et al., 2021). Puberty is the final stage of children's growth and development in which adult height, reproductive and psychosocial maturity is reached (Abe et al., 2017; Pei et al., 2020).

Height growth is one of the important things in life for certain people. The body needs to be measured to determine height; weight is used to determine a person's health (Li et al., 2017). Anthropometry represents the measurement of the human body (Goss et al., 2019). Anthropometric measurements usually include height, weight, and various body and limb measurements (Agata & Monyeki, 2018).

It is very important to measure age-related height. Child height is an important indicator of human capital and human development (Alves Junior et al., 2021). Health is the beginning of life for humans (Phukan et al., 2021; Sulaiman et al., 2020). Nutrition received by the body can shape the child's height and provide health (Vyas et al., 2016). Therefore, height measurement can be valuable data shaping a child's height growth model (Haile et al., 2016; Socha & Tymińska-Czabańska, 2019). Human physical form is an important measure and is the main capital of health (Spears, 2020).

Children must be able to grow and develop properly. Good growth will affect the movement carried out, such as exercising (Faria et al., 2020; Kim et al., 2021). The growth and development of children are expected to run simultaneously, both cognitive, affective, psychomotor, physiological, psychological, and social (Karra & Fink, 2019). The height achievement of each child is different because each child has a high, medium, and low body (Heshin-Bekenstein et al., 2018). The growth and development of children are influenced, among others, by factors such as conditions before birth (Pomeroy et al., 2021). Prenatal exposure to polychlorinated biphenyls (PCBs) may long-term affect growth in early adulthood among girls and identifies the peri-pubertal period as another sensitivity window for such measures (Bhowmik & Das, 2017; Manzoni et al., 2019; Soekatri et al., 2020). PCB exposure and body size are documented in several periods, from infancy to adulthood (Tahir et al., 2020). Prenatal conditions, reflected in birth weight, are more important in determining the height trajectory than postnatal factors, which do not help children fully recover from early growth deficits (Krishna et al., 2016).

Adolescence will increase children's activity in sports. Children who practice sports are not required to have a proportional height (Mackala et al., 2020). On the other hand, some sports require height, such as volleyball and basketball (S. Shen et al., 2021). In addition, height is one of the components recommended for growth in elementary school to high school (Junior et al., 2019).

Children's height is a concern for several parties in maintaining the health of their children. In addition, nutrition and lifestyle must be maintained for children's health in the future (Priymak et al., 2021; Wijaya-Erhardt, 2019). Height for age reflects the accumulation of malnutrition over the years, affecting children's educational achievement (Adami et al., 2021; A. Shen et al., 2020). To estimate a child's height when growing up, we need a method that can be used to predict a child's height (Ruchman et al., 2020). So that children will be taught to exercise to maintain their physical fitness, this study aims to formulate guidelines for predicting the maximum height of men and women.

MATERIALS AND METHODS

The method used in this research is descriptive development with a cross-sectional technique. The study observed the growth and development of height for girls from 0-18 years old and boys from 0-19 years old. The research was conducted in the city of Surakarta, Central Java, Indonesia. The research sample was obtained from Posyandu, early childhood schools, elementary schools, junior high schools, and high schools. Data were collected by observing the respondents' height (Mustafa et al., 2019). Height should be measured with the child standing straight, without shoes, with his back against the wall. The data were analysed using the Miles and Huberman model, namely data reduction, tabulation, data presentation, and concluding.

RESULTS AND DISCUSSION

The growth and development of female

The results of measuring the growth and development of female height in the study can be seen in Table 1.

Table 1. Age, mean, and percentage of females aged 0-18 years.

No.	Age (Year)	Average (cm)	Percentage (%)
1.	0	47.95	30.73
2.	1	71.25	45.66
3.	2	87.25	55.91
4.	3	95.18	60.99
5.	4	99.17	63.55
6.	5	104.30	66.20
7.	6	113.09	66.54
8.	7	117.09	75.04
9.	8	123.64	79.24
10.	9	127.28	81.57
11.	10	137.89	88.36
12.	11	142.21	91.14
13.	12	150.43	96.40
14.	13	152.38	97.65
15.	14	154.04	98.78
16.	15	154.66	99.11
17.	16	155.52	99.66
18.	17	155.61	99.72
19.	18	156.04	100.00

Table 1 shows that the growth and development of women's height from 13 years shows a stable change. However, growth in height is slow at the age of 13 to 18 years (Nezami et al., 2016). At the age of 16 years, the child did not show changes in height growth. Finally, 17-18 years old showed stagnant changes, meaning that there was no longer an increase in height (Kokun et al., 2021).

Growth and development of male

The results of measuring the growth and development of height for males in the study can be seen in Table 2.

Table 2. Age, mean, and percentage of males aged 0-19 years.

No.	Age (Year)	Average (cm)	Percentage (%)
1.	0	48.22	28.87
2.	1	72.33	43.30
3.	2	88.73	53.13
4.	3	96.46	57.75
5.	4	101.85	60.98
6.	5	106.86	63.98
7.	6	113.44	67.92
8.	7	117.19	70.16
9.	8	123.74	74.09
10.	9	129.21	77.36
11.	10	136.64	81.81
12.	11	140.72	84.25
13.	12	148.00	88.61
14.	13	155.39	93.04
15.	14	160.00	95.80
16.	15	163.41	97.64
17.	16	165.64	99.17
18.	17	166.21	99.41
19.	18	166.71	99.81
20.	19	167.02	100.00

Table 2 shows the growth and development of both male's heights from 0 to 15 years of age. Age 17 years, there is no significant increase in size. Ages 18-19 in males experience stagnant growth (A. Shen et al., 2019). Stagnant growth means that at the age of 18-19 years, males do not experience height growth.

Comparison of the percentage of growth and development of height for women and men

The results of measuring the growth and development of height for women and men in the study can be seen in Table 3.

Measurements for predicting a child's height can be done by considering weight and age percentage. The formula for calculating maximum height can be seen below

$$\text{Maximum Height} = 100\% : \text{Total Percentage } X - \text{height at age}$$

Assume that child A is a boy who is 6 years old and 114 cm tall. Then it can be obtained by 100% divided by 63.98 and multiplied by 114 cm then the result is 178.68 cm. Then the child, based on the prediction, will have a maximum height of 178.68 cm when he is 19 years old. Thus, heritability for human height, which measures the relative importance of additive genetic factors and environmental factors in expressing a trait, has been consistently estimated (Subekti et al., 2020). First, of course, from the nutritional factors needed, protein for cell growth and repair, calcium for bones, followed by balanced nutrition from other sources (Acharya et al., 2020; Dogbe & Revoredo-Giha, 2021; Sinharoy et al., 2021). While the environment, in this case, means exercise to circulate blood and help the metabolism run properly (Valencia et al., 2021). However, it should be noted that height gain can be made if the growth plates in the long bones are still open (Saadat Mostafavi et al., 2021). This is because the plate opens at puberty and will generally close at 20-21

years (Agostinete et al., 2017). Therefore, in theory, your growth plate is likely to be/already closed, and if it has closed, then your growth potential.

Table 3. Age and percentage of female aged 0-18 years and male old 0-19 years.

No.	Age (Year)	Percentage	
		Female	Male
1.	0	30.73	28.87
2.	1	45.66	43.30
3.	2	55.91	53.13
4.	3	60.99	57.75
5.	4	63.55	60.98
6.	5	66.84	63.98
7.	6	72.47	67.92
8.	7	75.04	70.16
9.	8	79.24	74.09
10.	9	81.57	77.36
11.	10	88.36	81.81
12.	11	91.14	84.25
13.	12	96.40	88.61
14.	13	97.65	93.04
15.	14	98.78	95.80
16.	15	99.11	97.64
17.	16	99.66	99.17
18.	17	99.72	99.41
19.	18	100.00	99.81
20.	19		100.00

CONCLUSION

Every year the growth of the child's height will increase according to the age of growth. Data on the growth of children's height can predict the maximum height of children at the age of 18 years for girls and 19 years for boys. The results of this study can be used to predict the development of children's height. This research is expected to provide a way to determine children's height in the future. The method of predicting a child's height can be applied to determine a child's height in the future. This method provides an overview of the child's height as an adult. This research is expected to provide information for parents, health workers, and children to maintain physical fitness. It is recommended to do sports that burden the long bones of the legs, such as athletics, leisurely running, skipping, basketball, badminton and other similar sports. In this way, the bones are stimulated to grow a little more due to the pounding of the weight. Swimming can also increase a person's height.

AUTHORS CONTRIBUTIONS

Sapta Kunta Purnama conceived and designed the analysis. Muhammad Furqon Hidayatullah designed the data analysis. Fadilah Umar collected the data. Lamet Riyadi contributed data. Siti Nurunnayah performed analysis tools. Eppy Setiyowati performed the analysis. Hilwah Nora wrote the paper. Agus Mukholid wrote the paper.

SUPPORTING AGENCIES

No funding agencies were reported by the authors.

DISCLOSURE STATEMENT

No potential conflict of interest were reported by the authors.

ACKNOWLEDGMENT

The authors would like to thank the Universitas Sebelas Maret, Surakarta, for the support to complete this research.

REFERENCES

- Abe, T., Kamada, M., Kitayuguchi, J., Okada, S., Mutoh, Y., & Uchio, Y. (2017). Is being a regular player with fewer teammates associated with musculoskeletal pain in youth team sports? A cross-sectional study. *BMC Musculoskeletal Disorders*, 18(1), 1–8. <https://doi.org/10.1186/s12891-017-1470-z>
- Acharya, Y., Luke, N., Naz, S., & Sharma, D. (2020). Exposure to conflict-related violence and nutritional status of children in Iraq. *SSM - Population Health*, 11, 100585. <https://doi.org/10.1016/j.ssmph.2020.100585>
- Adami, P. E., Rocchi, J. E., Melke, N., & Macaluso, A. (2021). Physiological Profile of High Intensity Functional Training Athletes. *Journal of Human Sport and Exercise*, 16(3), 675–688. <https://doi.org/10.14198/jhse.2021.163.16>
- Agata, K., & Monyeki, M. A. (2018). Association between sport participation, body composition, physical fitness, and social correlates among adolescents: The PAHL study. *International Journal of Environmental Research and Public Health*, 15(12), 1–16. <https://doi.org/10.3390/ijerph15122793>
- Agostinete, R. R., Ito, I. H., Kemper, H., Pastre, C. M., Rodrigues-Júnior, M. A., Luiz-De-Marco, R., & Fernandes, R. A. (2017). Maturação somática e o relacionamento entre variáveis minerais ósseas e modalidades esportivas em adolescentes: Estudo transversal. *Sao Paulo Medical Journal*, 135(3), 253–259. <https://doi.org/10.1590/1516-3180.2016.0270210217>
- Alves Junior, C. A. S., Moraes, M. S., de Souza, C. S., Costa, G., & Silva, D. A. S. (2021). Body composition among university female athletes of team sports. *Revista Brasileira de Medicina Do Esporte*, 27(2), 156–160. https://doi.org/10.1590/1517-8692202127022020_0046
- Bhowmik, K. R., & Das, S. (2017). On exploring and ranking risk factors of child malnutrition in Bangladesh using multiple classification analysis. *BMC Nutrition*, 3(1), 1–10. <https://doi.org/10.1186/s40795-017-0194-7>
- Dogbe, W., & Revoredo-Giha, C. (2021). Nutritional Implications of Trade-Offs Between Fresh and Processed Potato Products in the United Kingdom (UK). *Frontiers in Nutrition*, 7(11). <https://doi.org/10.3389/fnut.2020.614176>
- Faria, F., Howe, C., Faria, R., Andaki, A., Marins, J. C., & Amorim, P. R. (2020). Impact of recreational sports activities on metabolic syndrome components in adolescents. *International Journal of Environmental Research and Public Health*, 17(1). <https://doi.org/10.3390/ijerph17010143>
- Feka, K., Pomara, F., Russo, G., Piccione, M. C., Petrucci, M., Giustino, V., Messina, G., Iovane, A., Palma, A., & Bianco, A. (2019). How do sports affect static baropodometry? An observational study among women living in southern Italy. *Human Movement*, 20(1), 9–16. <https://doi.org/10.5114/hm.2019.78091>

- Goss, C. W., O'Brian, K., Dubray, C., Fischer, P. U., Hardy, M., Jambulingam, P., King, C. L., Laman, M., Lemoine, J. F., Robinson, L. J., Samuela, J., Subramanian, S., Supali, T., Weil, G. J., & Schechtman, K. B. (2019). Dosing pole recommendations for lymphatic filariasis elimination: A height-weight quantile regression modeling approach. *PLoS Neglected Tropical Diseases*, 13(7), 1–20. <https://doi.org/10.1371/journal.pntd.0007541>
- Haile, D., Nigatu, D., Gashaw, K., & Demelash, H. (2016). Height for age z score and cognitive function are associated with academic performance among school children aged 8-11 years old. *Archives of Public Health*, 74(1), 1–7. <https://doi.org/10.1186/s13690-016-0129-9>
- Heshin-Bekenstein, M., Perl, L., Hersh, A. O., von Scheven, E., Yelin, E., Trupin, L., Yazdany, J., & Lawson, E. F. (2018). Final adult height of patients with childhood-onset systemic lupus erythematosus: A cross sectional analysis. *Pediatric Rheumatology*, 16(1), 1–9. <https://doi.org/10.1186/s12969-018-0239-8>
- Junior, D. B. R., Vianna, J. M., Lauria, A. de A., Coelho, E. F., & Werneck, F. Z. (2019). Sports potential modeling of young basketball players: A preliminary analysis. *Revista Brasileira de Cineantropometria e Desempenho Humano*, 21. <https://doi.org/10.1590/1980-0037.2019v21e59832>
- Karra, M., & Fink, G. (2019). Long run height and education implications of early life growth faltering: A synthetic panel analysis of 425 birth cohorts in 21 low- and middle-income countries. *BMC Public Health*, 19(1), 1–8. <https://doi.org/10.1186/s12889-019-7203-5>
- Kim, K., Melough, M. M., Kim, D., Sakaki, J. R., Lee, J., Choi, K., & Chun, O. K. (2021). Nutritional adequacy and diet quality are associated with standardized height-for-age among U.S. children. *Nutrients*, 13(5), 1–10. <https://doi.org/10.3390/nu13051689>
- Kokun, O., Serdiuk, L., & Shamysh, O. (2021). Personal characteristics supporting Paralympic athletes' self-realization in sports. *Journal of Human Sport and Exercise*, 16(2), 435–444. <https://doi.org/10.14198/jhse.2021.162.17>
- Krishna, A., Fink, G., Berkman, L. F., & Subramanian, S. V. (2016). Short- and long-run associations between birth weight and children's height. *Economics and Human Biology*, 21, 156–166. <https://doi.org/10.1016/j.ehb.2016.02.004>
- Li, X. Y., Jia, C., & Zhang, Z. C. (2017). The normal range of maximum mouth opening and its correlation with height or weight in the young adult Chinese population. *Journal of Dental Sciences*, 12(1), 56–59. <https://doi.org/10.1016/j.jds.2016.09.002>
- Mackala, K., Rauter, S., Simenko, J., Kreft, R., Stodolka, J., Krizaj, J., Coh, M., & Vodicar, J. (2020). The effect of height on drop jumps in relation to somatic parameters and landing kinetics. *International Journal of Environmental Research and Public Health*, 17(16), 1–11. <https://doi.org/10.3390/ijerph17165886>
- Manzoni, G., Laillou, A., Samnang, C., Hong, R., Wieringa, F. T., Berger, J., Poirot, E., & Checchi, F. (2019). Child-sensitive WASH composite score and the nutritional status in Cambodian children. *Nutrients*, 11(9). <https://doi.org/10.3390/nu11092142>
- Mascherini, G., Galanti, G., Massetti, L., Calà, P., & Modesti, P. A. (2019). Growth charts for height, weight, and bmi (6–18 y) for the tuscan youth sports population. *International Journal of Environmental Research and Public Health*, 16(24). <https://doi.org/10.3390/ijerph16244975>
- Mustafa, M. Z., Buntat, Y., Omar, M. A., Razzaq, A. R. A., & Ahad, R. (2019). Organizational commitment survey: A confirmatory factorial analysis based on vocational colleges teachers sample. *International Journal of Engineering and Advanced Technology*, 8(5C), 279–288. <https://doi.org/10.35940/ijeat.E1041.0585C19>
- Nezami, M., Segovia-Siapco, G., Beeson, W. L., & Sabaté, J. (2016). Associations between consumption of dairy foods and anthropometric indicators of health in adolescents. *Nutrients*, 8(7). <https://doi.org/10.3390/nu8070427>

- Pei, W., Jain, A., Zhao, G., Feng, B., Xu, D., & Wang, X. (2020). Knockdown of OsSAE1a affects the growth and development and phosphate homeostasis in rice. *Journal of Plant Physiology*, 255, 153275. <https://doi.org/10.1016/j.jplph.2020.153275>
- Phukan, M. I., Thapa, R. K., Kumar, G., Bishop, C., Chaabene, H., & Ramirez-Campillo, R. (2021). Inter-limb jump asymmetries and their association with sport-specific performance in young male and female swimmers. *International Journal of Environmental Research and Public Health*, 18(14). <https://doi.org/10.3390/ijerph18147324>
- Pomeroy, E., Stock, J. T., & Wells, J. C. K. (2021). Population history and ecology, in addition to climate, influence human stature and body proportions. *Scientific Reports*, 11(1), 1–10. <https://doi.org/10.1038/s41598-020-79501-w>
- Priymak, S., Krutsevich, T., Pangelova, N., Trachuk, S., Kravchenko, T., Stepanenko, V., & Ruban, V. (2021). Modeling of Functional Support of Sports Activities of Biathletes of Different Qualifications. *Journal of Human Sport and Exercise*, 16(1), 136–146. <https://doi.org/10.14198/jhse.2021.161.12>
- Reinehr, T., Carlsson, M., Chrysis, D., & Camacho-Hübner, C. (2020). Adult height prediction by bone age determination in children with isolated growth hormone deficiency. *Endocrine Connections*, 9(5), 370–378. <https://doi.org/10.1530/EC-20-0090>
- Ruchman, S. G., Green, A. S., Schonholz, S., Hersh, E., Sikka, N., Baranowski, K. A., & Singer, E. K. (2020). A toolkit for building medical programs for asylum seekers: Resources from the Mount Sinai Human Rights Program. *Journal of Forensic and Legal Medicine*, 75. <https://doi.org/10.1016/j.jflm.2020.102037>
- Saadat Mostafavi, S. R., Memarian, A., Motamedi, O., Mohamadi nejad khanamani, M., Khaleghi, M., & Habibi, S. (2021). Fourth lumbar vertebral parameters in predicting the gender, height and age in Iranian population. *Forensic Science International: Reports*, 3(January), 100175. <https://doi.org/10.1016/j.fsir.2021.100175>
- Shen, A., Bernabé, E., & Sabbah, W. (2019). The bidirectional relationship between weight, height and dental caries among preschool children in China. *PLoS ONE*, 14(4), 1–13. <https://doi.org/10.1371/journal.pone.0216227>
- Shen, A., Bernabé, E., & Sabbah, W. (2020). The socioeconomic inequality in increment of caries and growth among Chinese children. *International Journal of Environmental Research and Public Health*, 17(12), 1–9. <https://doi.org/10.3390/ijerph17124234>
- Shen, S., Suzuki, K., Kohmura, Y., Fuku, N., Someya, Y., & Naito, H. (2021). Engagement in different sport disciplines during university years and risk of locomotive syndrome in older age: J-Fit+Study. *Environmental Health and Preventive Medicine*, 26(1), 1–11. <https://doi.org/10.1186/s12199-021-00958-w>
- Silva, B., Cruz, G., Rocha-Rodrigues, S., & Clemente, F. M. (2021). Monitoring physical performance and training load in young surf athletes. *Journal of Human Sport and Exercise*, 16(2), 261–272. <https://doi.org/10.14198/jhse.2021.162.03>
- Sinharoy, S. S., Reese, H. E., Prahara, I., Chang, H. H., & Clasen, T. (2021). Effects of a combined water and sanitation intervention on biomarkers of child environmental enteric dysfunction and associations with height-for-age Z-score: A matched Cohort study in rural Odisha, India. *PLoS Neglected Tropical Diseases*, 15(3), 1–13. <https://doi.org/10.1371/journal.pntd.0009198>
- Socha, J., & Tymińska-Czabańska, L. (2019). A method for the development of dynamic site index models using height-age data from temporal sample plots. *Forests*, 10(7). <https://doi.org/10.3390/f10070542>
- Soekatri, M. Y. E., Sandjaja, S., & Syauby, A. (2020). Stunting was associated with reported morbidity, parental education and socioeconomic status in 0.5–12-year-old Indonesian children. *International*

- Journal of Environmental Research and Public Health, 17(17), 1–9. <https://doi.org/10.3390/ijerph17176204>
- Spears, D. (2020). Exposure to open defecation can account for the Indian enigma of child height. *Journal of Development Economics*, 146. <https://doi.org/10.1016/j.jdeveco.2018.08.003>
- Subekti, N., Sistiasih, V. S., Syaokani, A. A., & Fatoni, M. (2020). Kicking ability in pencak silat, reviewed from eye-foot coordination, speed, and ratio of limb length-body height. *Journal of Human Sport and Exercise*, 15(Proc2), 453–461. <https://doi.org/10.14198/jhse.2020.15.Proc2.36>
- Sulaiman, W. M. H. Bin, Jumintono, Wahyuniati, C. F. S., Suryandartiwi, W., Sunardi, J., Umar, F., Manoppo, F. K., Yudanto, & Hos, J. (2020). Quickly understanding on progressive muscle relaxation with video-based learning in secondary school students. *International Journal of Human Movement and Sports Sciences*, 8(5), 181–185. <https://doi.org/10.13189/saj.2020.080504>
- Tahir, E., Cordier, S., Courtemanche, Y., Forget-Dubois, N., Desrochers-Couture, M., Bélanger, R. E., Ayotte, P., Jacobson, J. L., Jacobson, S. W., & Muckle, G. (2020). Effects of polychlorinated biphenyls exposure on physical growth from birth to childhood and adolescence: A prospective cohort study. *Environmental Research*, 189(July). <https://doi.org/10.1016/j.envres.2020.109924>
- Valencia, O., Saka, C., Ramos, C., Caparrós-manosalva, C., & Guzmán-venegas, R. (2021). Kinetic Effect of the Taping on the Ankle During a Change of Direction in Basketball Players. *Journal of Human Sport and Exercise*, 16(3), 711–720. <https://doi.org/10.14198/jhse.2021.163.19>
- Vyas, S., Kov, P., Smets, S., & Spears, D. (2016). Disease externalities and net nutrition: Evidence from changes in sanitation and child height in Cambodia, 2005–2010. *Economics and Human Biology*, 23, 235–245. <https://doi.org/10.1016/j.ehb.2016.10.002>
- Wang, L., Shi, C., & Zhao, Z. (2018). The Research and Development of Growth Curve for Children's Height and Weight on Android Platform. 2017 10th International Congress on Image and Signal Processing, BioMedical Engineering and Informatics (CISP-BMEI 2017), 1–5. <https://doi.org/10.1109/CISP-BMEI.2017.8302293>
- Wijaya-Erhardt, M. (2019). Nutritional status of Indonesian children in low-income households with fathers that smoke. *Osong Public Health and Research Perspectives*, 10(2), 64–71. <https://doi.org/10.24171/j.phrp.2019.10.2.04>



This work is licensed under a [Attribution-NonCommercial-NoDerivatives 4.0 International](https://creativecommons.org/licenses/by-nc-nd/4.0/) (CC BY-NC-ND 4.0).

Observation of children's age to predict maximum height

ORIGINALITY REPORT

17 %

SIMILARITY INDEX

16 %

INTERNET SOURCES

10 %

PUBLICATIONS

2 %

STUDENT PAPERS

MATCH ALL SOURCES (ONLY SELECTED SOURCE PRINTED)

2%

★ www.readkong.com

Internet Source

Exclude quotes On

Exclude matches Off

Exclude bibliography On